

PHY250: Review and Introduction

Anabela R. Turlione

Digipen Institute

Bilbao, Fall 2021

Introduction

Review of PHY200
PHY250

Until now, you have been describing the motion of a particle, using the equations of Newton:

Until now, you have been describing the motion of a particle, using the equations of Newton:

$$\sum_i \vec{F}_i = m\vec{a} \quad (1)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

And you have proved that:

$$W = \Delta E_k \quad (3)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

And you have proved that:

$$W = \Delta E_k \quad (3)$$

Where

$$E_k = \frac{1}{2}mv^2 \quad (4)$$

is the kinetic energy.

And, there are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

And, there are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell}$$

And, there are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = W \quad (6)$$

And, there are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = W \quad (6)$$

In this case, the total energy of the system is conserved:

And, there are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = W \quad (6)$$

In this case, the total energy of the system is conserved:


$$E = U + K = \text{constant} \quad (7)$$

So, when a force is conservative, that is, there is a function $U(x)$ such that

$$F = -\frac{U(x)}{dx} \quad (8)$$

then, the energy of the particle is constant.

You also studied systems of N particles,:



And you defined the following vector quantities:

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

$$\vec{\tau}_i = \vec{r}_i \times \vec{F}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

$$\vec{\tau}_i = \vec{r}_i \times \vec{F}_i \rightarrow \vec{\tau} = \sum_i^N \vec{\tau}_i \quad (\text{Torque}) \quad (11)$$

And found some interesting relations:

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F}$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (*Conservation of Linear Momentum*)} \quad (12)$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (Conservation of Linear Momentum)} \quad (12)$$

$$\frac{d\vec{L}}{dt} = \vec{\tau}$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (Conservation of Linear Momentum)} \quad (12)$$

$$\frac{d\vec{L}}{dt} = \vec{\tau} \text{ (Conservation of Angular Momentum)} \quad (13)$$

Finally, you have defined the center of mass of the system:

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

And you found the following expressions that relates the motion of the center of mass with the total external force:

$$\vec{P} = M \frac{d}{dt}(\vec{R}_{CM}) = M \vec{V}_{CM} \quad (15)$$

$$\vec{F}^{ext} = M \frac{d^2}{dt^2}(\vec{R}_{CM}) = M \vec{a}_{CM} \quad (16)$$

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

And you found the following expressions that relates the motion of the center of mass with the total external force:

$$\vec{P} = M \frac{d}{dt}(\vec{R}_{CM}) = M \vec{V}_{CM} \quad (15)$$

$$\vec{F}^{ext} = M \frac{d^2}{dt^2}(\vec{R}_{CM}) = M \vec{a}_{CM} \quad (16)$$

So the center of mass moves as a particle acted by the total external force.

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

$$\vec{L} = I\vec{\omega} \quad (17)$$

$$\vec{\tau} = I\vec{\alpha} \quad (18)$$

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

$$\vec{L} = I\vec{\omega} \quad (17)$$

$$\vec{\tau} = I\vec{\alpha} \quad (18)$$

where **I** is the **Moment of Inertia**

$$I = \sum_i^N m_i r_i^2 \quad (19)$$

The total kinetic energy is:

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i}$$

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

Then, the motion of a system can be decomposed in:

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

Then, the motion of a system can be decomposed in:

Translation of CM + Rotation Around CM

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM}$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

$$\vec{F} = M\vec{a}_{CM}$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

$$\vec{F} = M\vec{a}_{CM} \leftrightarrow \vec{\tau} = I\vec{\alpha} \quad (22)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2 \rightarrow \int_V \rho(r) r^2 dv \quad (24)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2 \rightarrow \int_V \rho(r) r^2 dv \quad (24)$$

In PHY250 we are going to extend the study of systems:

In PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary

In PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary
- ▶ we are going to study how the energy is propagated

In PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary
- ▶ we are going to study how the energy is propagated
- ▶ and we are going to study the thermal properties of matter

In PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary
- ▶ we are going to study how the energy is propagated
- ▶ and we are going to study the thermal properties of matter
- ▶ finally, we are going to study the nature of light and how it interacts with different mediums