

PHY250: Review and Introduction

Anabela R. Turlione

Digipen Institute


Bilbao, Fall 2022

Introduction

PHY250 in a nutshell


Review of PHY200

In PHY 200 you studied:



In PHY 200 you studied:

System of particles →



→ CM

→ Torque


→ Angular
Momentum

→ Inertia Moment

In PHY 200 you studied:

Continuous systems
constant distances

↓
Rigid bodies



CM
Particle Motion

↓
Rot + Trans

↑
Torque
Angular Momentum
Inertia

In PHY250 we are going to study fluids, we can think of them as systems of particles.



We also are going to study mechanics waves, energy traveling in a medium. We can think of them as the collective motion of the particles in a continuous system.



Sumarizing...in PHY250 we are going to extend the study of systems:

Sumarizing...in PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary

Sumarizing...in PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary
- ▶ we are going to study how the energy is propagated

Sumarizing...in PHY250 we are going to extend the study of systems:

- ▶ this time, the distance between their particles may vary
- ▶ we are going to study how the energy is propagated
- ▶ finally, we are going to study the nature of light and how it interacts with different mediums

So far, you have been describing the motion of a particle, using the equations of Newton:

So far, you have been describing the motion of a particle, using the equations of Newton:

$$\sum_i \vec{F}_i = m\vec{a} \quad (1)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

And you have proved that:

$$W = \Delta E_k \quad (3)$$

Then, you have defined the work done by a force,

$$W = \int_1^2 \vec{F} \cdot d\vec{\ell} \quad (2)$$

And you have proved that:

$$W = \Delta E_k \quad (3)$$

Where

$$E_k = \frac{1}{2}mv^2 \quad (4)$$

is the kinetic energy.

There are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

There are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell}$$

There are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = -W \quad (6)$$

There are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = -W \quad (6)$$

In this case, the total energy of the system is conserved:

There are some kind of forces that have the following property:

$$\oint \vec{F} \cdot d\vec{\ell} = 0 \quad (5)$$

In this case, we can define the potential energy, and we say that the force is **conservative**

$$\Delta U(\vec{r}) = - \int_{\vec{r}_0}^{\vec{r}} \vec{F} \cdot d\vec{\ell} \rightarrow \Delta U = -W \quad (6)$$

In this case, the total energy of the system is conserved:


$$E = U + K = \text{constant} \quad (7)$$

So, when a force is conservative, that is, there is a function $U(x)$ such that

$$F = -\frac{U(x)}{dx} \quad (8)$$

then, the energy of the particle is constant.

You also studied systems of N particles,:



And you defined the following vector quantities:

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

$$\vec{\tau}_i = \vec{r}_i \times \vec{F}_i$$

And you defined the following vector quantities:

$$\vec{p}_i = m_i \vec{v}_i \rightarrow \vec{P} = \sum_i^N \vec{p}_i \quad (\text{Linear Momentum}) \quad (9)$$

$$\vec{L}_i = \vec{r}_i \times \vec{p}_i \rightarrow \vec{L} = \sum_i^N \vec{L}_i \quad (\text{Angular Momentum}) \quad (10)$$

$$\vec{\tau}_i = \vec{r}_i \times \vec{F}_i \rightarrow \vec{\tau} = \sum_i^N \vec{\tau}_i \quad (\text{Torque}) \quad (11)$$

And found some interesting relations:

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F}$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (Conservation of Linear Momentum)} \quad (12)$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (Conservation of Linear Momentum)} \quad (12)$$

$$\frac{d\vec{L}}{dt} = \vec{\tau}$$

And found some interesting relations:

$$\frac{d\vec{P}}{dt} = \vec{F} \text{ (Conservation of Linear Momentum)} \quad (12)$$

$$\frac{d\vec{L}}{dt} = \vec{\tau} \text{ (Conservation of Angular Momentum)} \quad (13)$$

Finally, you have defined the center of mass of the system:

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

And you found the following expressions that relates the motion of the center of mass with the total external force:

$$\vec{P} = M \frac{d}{dt}(\vec{R}_{CM}) = M \vec{V}_{CM} \quad (15)$$

$$\vec{F}^{ext} = M \frac{d^2}{dt^2}(\vec{R}_{CM}) = M \vec{a}_{CM} \quad (16)$$

Finally, you have defined the center of mass of the system:

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N = m_i \vec{r}_i \quad (14)$$

And you found the following expressions that relates the motion of the center of mass with the total external force:

$$\vec{P} = M \frac{d}{dt}(\vec{R}_{CM}) = M \vec{V}_{CM} \quad (15)$$

$$\vec{F}^{ext} = M \frac{d^2}{dt^2}(\vec{R}_{CM}) = M \vec{a}_{CM} \quad (16)$$

So the center of mass moves as a particle acted by the total external force.

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

$$\vec{L} = I\vec{\omega} \quad (17)$$

$$\vec{\tau} = I\vec{\alpha} \quad (18)$$

In particular, for rotations of **Rigid bodies** (the distances between particles are constant) in 2D:

$$\vec{L} = I\vec{\omega} \quad (17)$$

$$\vec{\tau} = I\vec{\alpha} \quad (18)$$

where **I** is the **Moment of Inertia**

$$I = \sum_i^N m_i r_i^2 \quad (19)$$

The total kinetic energy is:

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i}$$

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

Then, the motion of a system can be decomposed in:

The total kinetic energy is:

$$E_k = \sum_i^N E_{k,i} = \underbrace{\frac{1}{2} M V_{CM}^2}_{\text{Translational } E_k} + \underbrace{\frac{1}{2} I \omega^2}_{\text{Rotational } E_k} \quad (20)$$

Then, the motion of a system can be decomposed in:

Translation of CM + Rotation Around CM

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM}$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

$$\vec{F} = M\vec{a}_{CM}$$

We can make a parallelism between the equations for cartesian coordinates and the equations for angular coordinates:

$$\vec{P} = M\vec{V}_{CM} \leftrightarrow \vec{L} = I\vec{\omega} \quad (21)$$

$$\vec{F} = M\vec{a}_{CM} \leftrightarrow \vec{\tau} = I\vec{\alpha} \quad (22)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2 \rightarrow \int_V \rho(r) r^2 dv \quad (24)$$

Finally, for continuous bodies, all the sums turns into integrals

$$\vec{R}_{CM} = \frac{1}{M} \sum_i^N m_i \vec{r}_i \rightarrow \int_V \rho(r) \vec{r} dv \quad (23)$$

$$I = \sum_i^N m_i r_i^2 \rightarrow \int_V \rho(r) r^2 dv \quad (24)$$