

PHY115: Light

Anabela R. Turlione

Digipen

Spring 2021

Light

Introduction

Waves, Wave Fronts, and Rays

Interaction of Light with different materials

Introduction

What is light?

Introduction

What is light?

Light is Electromagnetic radiation

Nature of light

Is it an electromagnetic wave or particles?

Nature of light

Is it an electromagnetic wave or particles?

Both!

Nature of light

Is it an electromagnetic wave or particles?

Both!

Some effects can be better explained if we consider the light as a wave (interference, diffraction).

Nature of light

Is it an electromagnetic wave or particles?

Both!

Some effects can be better explained if we consider the light as a wave (interference, diffraction).

Some other can be better explained if we consider it as particles (emission and absorption).

Sources of light

The fundamental sources of all electromagnetic radiation are electric charges in accelerated motion.

Sources of light

The fundamental sources of all electromagnetic radiation are electric charges in accelerated motion.

- ▶ Thermal radiation: all bodies emit electromagnetic radiation as a result of thermal motion of their molecules.

Sources of light

The fundamental sources of all electromagnetic radiation are electric charges in accelerated motion.

- ▶ Thermal radiation: all bodies emit electromagnetic radiation as a result of thermal motion of their molecules.
- ▶ Light is also produced during electrical discharges through ionized gases.

Sources of light

The fundamental sources of all electromagnetic radiation are electric charges in accelerated motion.

- ▶ Thermal radiation: all bodies emit electromagnetic radiation as a result of thermal motion of their molecules.
- ▶ Light is also produced during electrical discharges through ionized gases.
- ▶ In most light sources, light is emitted independently by different atoms within the source. Mixture of different wavelengths.

Sources of light

The fundamental sources of all electromagnetic radiation are electric charges in accelerated motion.

- ▶ Thermal radiation: all bodies emit electromagnetic radiation as a result of thermal motion of their molecules.
- ▶ Light is also produced during electrical discharges through ionized gases.
- ▶ In most light sources, light is emitted independently by different atoms within the source. Mixture of different wavelengths.
- ▶ In a laser, atoms are induced to emit light in a cooperative, coherent fashion. Narrow and intense monochromatic beam of radiation

Sources of light

No matter what its source, electromagnetic radiation travels in vacuum at the same speed. The speed of light in vacuum is

Sources of light

No matter what its source, electromagnetic radiation travels in vacuum at the same speed. The speed of light in vacuum is


$$c = 3 \times 10^8 \text{ m/s} \quad (1)$$

Waves, Wave Fronts, and Rays

We define a wave front as the locus of all adjacent points at which the phase of vibration of a physical quantity associated with the wave is the same.


Waves, Wave Fronts, and Rays

Electromagnetic waves in vacuum spread out as spherical wave: waves fronts spread out uniformly in all directions from a point source.



Waves, Wave Fronts, and Rays


Electromagnetic waves in vacuum spread out as spherical wave: waves fronts spread out uniformly in all directions from a point source.



Far from the source: Plane waves

Waves, Wave Fronts, and Rays


Electromagnetic waves in vacuum spread out as spherical wave: waves fronts spread out uniformly in all directions from a point source.



Far from the source: Plane waves

Waves, Wave Fronts, and Rays

Electromagnetic waves in vacuum spread out as spherical wave: waves fronts spread out uniformly in all directions from a point source.



Far from the source: Plane waves

Waves, Wave Fronts, and Rays

To describe the directions in which light propagates, we represent a light wave by rays.


Waves, Wave Fronts, and Rays

To describe the directions in which light propagates, we represent a light wave by rays.

A ray is an imaginary line along the direction of travel of the wave.


(a)

When wave fronts are spherical, the rays radiate from the center of the sphere.



(b)

When wave fronts are planar, the rays are perpendicular to the wave fronts and parallel to each other.



Waves, Wave Fronts, and Rays

The branch of optics for which the ray description is adequate is called geometric optics; the branch dealing specifically with wave behavior is called physical optics.

Test Your Understanding of Section:

Some crystals are not isotropic: Light travels through the crystal at a higher speed in some directions than in others. In a crystal in which light travels at the same speed in the x and z -directions but at a faster speed in the y direction, what would be the shape of the wave fronts produced by a light source at the origin?


Test Your Understanding of Section:

Some crystals are not isotropic: Light travels through the crystal at a higher speed in some directions than in others. In a crystal in which light travels at the same speed in the x and z -directions but at a faster speed in the y direction, what would be the shape of the wave fronts produced by a light source at the origin?


1. spherical,
2. ellipsoidal, flattened along the y -axis
3. ellipsoidal, stretched out along the y -axis.

Reflection and refraction.


(a) Plane waves reflected and refracted from a window




(b) The waves in the outside air and glass represented by rays



(c) The representation simplified to show just one set of rays




Two type of reflections: Diffuse and Specular




Two type of reflections: Diffuse and Specular

(a) Specular reflection



(b) Diffuse reflection




Refraction


What happens when a waves front changes the medium?

Refraction


What happens when a waves front changes the medium?



Refraction




Refraction



$$\sin\theta_1 = \frac{v_1 t}{AD}, \quad \sin\theta_2 = \frac{v_2 t}{AD}$$

Refraction



$$\sin \theta_1 = \frac{v_1 t}{AD}, \quad \sin \theta_2 = \frac{v_2 t}{AD}$$

$$\rightarrow \frac{\sin \theta_1}{\sin \theta_2} = \frac{v_1}{v_2}$$

Refraction

The frequency does not change, but the wavelength does,

Refraction

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

Refraction

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

where,

$$n = \frac{c}{v} \quad (2)$$

Refraction

The frequency does not change, but the wavelength does,

$$\frac{\lambda_2}{\lambda_1} = \frac{v_2}{v_1} = \frac{n_1}{n_2}$$

where,

$$n = \frac{c}{v} \quad (2)$$

is the **refraction index**.

Refraction

Substance	Index of Refraction, n
Solids	
Ice (H_2O)	1.309
Fluorite (CaF_2)	1.434
Polystyrene	1.49
Rock salt (NaCl)	1.544
Quartz (SiO_2)	1.544
Zircon ($\text{ZrO}_2 \cdot \text{SiO}_2$)	1.923
Diamond (C)	2.417
Fabulite (SrTiO_3)	2.409
Rutile (TiO_2)	2.62
Glasses (typical values)	
Crown	1.52
Light flint	1.58
Medium flint	1.62
Dense flint	1.66
Lanthanum flint	1.80
Liquids at 20°C	
Methanol (CH_3OH)	1.329
Water (H_2O)	1.333
Ethanol ($\text{C}_2\text{H}_5\text{OH}$)	1.36
Carbon tetrachloride (CCl_4)	1.460

The Laws of Reflection and Refraction


1. The incident, reflected, and refracted rays and the normal to the surface all lie in the same plane (plane of incidence).
2. The angle of reflection θ_r is equal to the angle of incidence θ_a for all wavelengths and for any pair of materials.

$$\theta_r = \theta_a \text{ (law of reflection)} \quad (3)$$

3. For monochromatic light and for a given pair of materials,

$$n_a \sin \theta_a = n_b \sin \theta_b \text{ (law of refraction, Snell)} \quad (4)$$

1. The incident, reflected, and refracted rays and the normal to the surface all lie in the same plane.



3. When a monochromatic light ray crosses the interface between two given materials a and b , the angles θ_a and θ_b are related to the indexes of refraction of a and b by

$$\frac{\sin \theta_a}{\sin \theta_b} = \frac{n_b}{n_a}$$


Refraction

Example: a straight ruler half-immersed in water

(a) A straight ruler half-immersed in water




(b) Why the ruler appears bent



Refraction

Example: Mirages



Index of Refraction and the Wave Aspects of Light

- ▶ The frequency f of the wave does not change when passing from one material to another.


Index of Refraction and the Wave Aspects of Light

- ▶ The frequency f of the wave does not change when passing from one material to another.
- ▶ the wavelength of the wave is different in general in different materials. This is because in any material $v = \lambda f$

$$\lambda = \frac{\lambda_0}{n} \quad (5)$$

Huygens' principle

Every point on a wave front can be considered as a source of tiny wavelets that spread out in the forward direction at the speed of the wave itself. The new wave front is the envelope of all the wavelets—that is, the tangent to all of them.



Diffraction

What happens when waves impinge on an obstacle?

Diffraction

What happens when waves impinge on an obstacle?

→ waves bend in behind an obstacle

Diffraction

What happens when waves impinge on an obstacle?

→ waves bend in behind an obstacle

→ The bending into the "shadow region" is known as diffraction

Diffraction

What happens when waves impinge on an obstacle?

→ waves bend in behind an obstacle

→ The bending into the "shadow region" is known as diffraction

→ If the opening is much larger than the wavelength, diffraction goes unnoticed.


Diffraction

What happens when waves impinge on an obstacle?




→ waves bend in behind an obstacle

→ The bending into the "shadow region" is known as diffraction

→ If the opening is much larger than the wavelength, diffraction goes unnoticed.



Interference: Young's double-slit Experiment



If light consists of tiny particles, we might expect to see two bright lines on a screen placed behind the slits as in (b). But instead a series of bright lines are seen, as in (c). Young was able to explain this result as a wave-interference phenomenon.

Dispersion

- ▶ White Light \rightarrow superposition of waves with wavelengths extending throughout the visible spectrum

Dispersion

- ▶ White Light \rightarrow superposition of waves with wavelengths extending throughout the visible spectrum
- ▶ The speed of light in vacuum is the same for all wavelengths

Dispersion

- ▶ White Light \rightarrow superposition of waves with wavelengths extending throughout the visible spectrum
- ▶ The speed of light in vacuum is the same for all wavelengths
- ▶ The speed in a material substance is different for different wavelengths.

Dispersion

- ▶ White Light → superposition of waves with wavelengths extending throughout the visible spectrum
- ▶ The speed of light in vacuum is the same for all wavelengths
- ▶ The speed in a material substance is different for different wavelengths.
- ▶ **the index of refraction of a material depends on the wavelength.**

$$\lambda = \frac{\lambda_0}{n} \quad (6)$$


Dispersion

- ▶ White Light \rightarrow superposition of waves with wavelengths extending throughout the visible spectrum
- ▶ The speed of light in vacuum is the same for all wavelengths
- ▶ The speed in a material substance is different for different wavelengths.
- ▶ **the index of refraction of a material depends on the wavelength.**


$$\lambda = \frac{\lambda_0}{n} \quad (6)$$

- ▶ The dependence of wave speed and index of refraction on wavelength is called **dispersion**.

How much the light ray is bent, when it is diffracted depends on n (Snell), then, the bending is different for different wavelength.



How much the light ray is bent, when it is diffracted depends on n (Snell), then, the bending is different for different wavelength.



- ▶ So far, we have defined 3 different phenomenons: reflection, refraction, diffraction

- ▶ So far, we have defined 3 different phenomenons: reflection, refraction, diffraction
- ▶ Elastic collision of photons with surface \rightarrow reflection

- ▶ So far, we have defined 3 different phenomenons: reflection, refraction, diffraction
- ▶ Elastic collision of photons with surface → reflection
- ▶ The speed of light changes when it is transmitted in a medium → the ray is bent: refraction

- ▶ So far, we have defined 3 different phenomenons: reflection, refraction, diffraction
- ▶ Elastic collision of photons with surface → reflection
- ▶ The speed of light changes when it is transmitted in a medium → the ray is bent: refraction
- ▶ Huygen's principle → diffraction

How can we explain those phenomenons from a microscopical point of view?

- ▶ When a light ray hits a material, it interacts with charged particles in the material

How can we explain those phenomenons from a microscopical point of view?

- ▶ When a light ray hits a material, it interacts with charged particles in the material
- ▶ The interaction depends on the internal structure of the material.

How can we explain those phenomenons from a microscopical point of view?

- ▶ When a light ray hits a material, it interacts with charged particles in the material
- ▶ The interaction depends on the internal structure of the material.
- ▶ The EM field accelerates the charges, and they generate electromagnetic radiation

When a photon interacts with an atom inside the material 2 things can happen:

1. The photon is scattered, that is, its trajectory is deviated

When a photon interacts with an atom inside the material 2 things can happen:


1. The photon is scattered, that is, its trajectory is deviated
2. The photon is absorbed by the material

Scattering

Deviation of the trajectory of a photon by a charged particle.

Scattering

Deviation fo the trajectory of a photon by a charged particle.



Scattering

Deviation fo the trajectory of a photon by a charged particle.

Scattering

Deviation fo the trajectory of a photon by a charged particle.

- ▶ The EM field of the light accelerates the charge, and it emits a new EM field.

Scattering

Deviation of the trajectory of a photon by a charged particle.

- ▶ The EM field of the light accelerates the charge, and it emits a new EM field.
- ▶ We can have 2 kind of scattering:

Scattering

Deviation of the trajectory of a photon by a charged particle.

- ▶ The EM field of the light accelerates the charge, and it emits a new EM field.
- ▶ We can have 2 kind of scattering:
 1. Elastic: The energy is conserved, the incident angle is equal to the reflected.

Scattering

Deviation of the trajectory of a photon by a charged particle.


- ▶ The EM field of the light accelerates the charge, and it emits a new EM field.
- ▶ We can have 2 kind of scattering:
 1. Elastic: The energy is conserved, the incident angle is equal to the reflected.
 2. Inelastic: The energy is not conserved, the reflected angle changes. This happens when the light ray penetrates into the material and interacts with the atoms nuclei. The EM field makes the atoms to vibrate and is transformed into thermal energy.

Absorption

The energy of the photon is equal to the gap between electronic levels inside the atom, so electrons move to a level with a higher energy.


Absorption

The energy of the photon is equal to the gap between electronic levels inside the atom, so electrons move to a level with a higher energy.



Absorption

The energy of the photon is equal to the gap between electronic levels inside the atom, so electrons move to a level with a higher energy.



A photon can only be absorbed if its energy is equal to the gap.

Then, when light interacts with a material

Then, when light interacts with a material

- ▶ A part is reflected

Then, when light interacts with a material

- ▶ A part is reflected
- ▶ A part is transmitted

Then, when light interacts with a material

- ▶ A part is reflected
- ▶ A part is transmitted
- ▶ A part is absorbed

Then, when light interacts with a material

- ▶ A part is reflected
- ▶ A part is transmitted
- ▶ A part is absorbed

Then, when light interacts with a material

Then, when light interacts with a material

- ▶ A part is reflected: Elastic Scattering → Specular Reflection.
Inelastic Scattering → Diffuse Reflection.

Then, when light interacts with a material

- ▶ A part is reflected: Elastic Scattering → Specular Reflection.
Inelastic Scattering → Diffuse Reflection.
- ▶ A part is transmitted: The EM radiation accelerates charge particles inside, particles generate an EM radiation. The resulting EM radiation is the superposition of both → the velocity inside changes.

Then, when light interacts with a material

- ▶ A part is reflected: Elastic Scattering → Specular Reflection.
Inelastic Scattering → Diffuse Reflection.
- ▶ A part is transmitted: The EM radiation accelerates charge particles inside, particles generate an EM radiation. The resulting EM radiation is the superposition of both → the velocity inside changes.
- ▶ A part is absorbed: The frequencies that are absorbed are not transmitted or reflected.

Then, when light interacts with a material

- ▶ A part is reflected: Elastic Scattering → Specular Reflection.
Inelastic Scattering → Diffuse Reflection.
- ▶ A part is transmitted: The EM radiation accelerates charge particles inside, particles generate an EM radiation. The resulting EM radiation is the superposition of both → the velocity inside changes.
- ▶ A part is absorbed: The frequencies that are absorbed are not transmitted or reflected.

We only can observe Reflected or Transmitted frequencies. If some frequencies are absorbed, the Reflected or Transmitted light does not have all the frequencies, so it is not white anymore. → Color

Fresnel Reflectance


Fresnel Reflectance

The Fresnel coefficients describe the reflection and transmission of light. It gives us an idea of the ratio between the light reflected and the light transmitted.

- ▶ transparent materials have low Fresnel coefficients
- ▶ opaques materials have high Fresnel coefficients
- ▶ it depends on the angle of incidence
- ▶ it also depends on the Index Of Reflection (IOR)

Fresnel Reflectance

Fresnel Reflectance



So, what happens in different materials?

So, what happens in different materials?


METALS

So, what happens in different materials?

METALS

Metallic bonding can be thought as a cloud of positive charge ions immersed in a cloud of valence electrons.

metallic bond




METALS


METALS

The positive ions, form a net:


Common metallic crystal structures



body-centred cubic (bcc)



face-centred cubic (fcc)



hexagonal close-packed (hcp)

© 2012 Encyclopædia Britannica, Inc.

In metals:

In metals:

- ▶ Atoms have metallic bonds.

In metals:

- ▶ Atoms have metallic bonds.
- ▶ Electrons in higher levels are not constrained to a single atom, but are free to move around the material.

In metals:

- ▶ Atoms have metallic bonds.
- ▶ Electrons in higher levels are not constrained to a single atom, but are free to move around the material.
- ▶ → High electric conductivity, heat conductivity.

In metals:

- ▶ Atoms have metallic bonds.
- ▶ Electrons in higher levels are not constrained to a single atom, but are free to move around the material.
- ▶ → High electric conductivity, heat conductivity.
- ▶ When the metal is exposed to an electric field, the electrons are redistributed around the material so that the electric field inside is 0.

When a photon interacts with an atom in the metal, 3 things can happen:

When a photon interacts with an atom in the metal, 3 things can happen:

1. Elastic Scattering

When a photon interacts with an atom in the metal, 3 things can happen:

1. Elastic Scattering
2. Inelastic Scattering

When a photon interacts with an atom in the metal, 3 things can happen:

1. Elastic Scattering
2. Inelastic Scattering
3. Absorption

When a photon interacts with an atom in the metal, 3 things can happen:

1. Elastic Scattering
2. Inelastic Scattering: Low ratio, light intensity is highly dumped inside the metal. So it does not interact with the net.
3. Absorption

When a photon interacts with an atom in the metal, 3 things can happen:


1. Elastic Scattering
2. Inelastic Scattering: Low ratio, light intensity is highly dumped inside the metal. So it does not interact with the net.
3. Absorption: Low ratio, electrons energy gap is too big for visible light.

When a photon interacts with an atom in the metal, 3 things can happen:

1. Elastic Scattering: High ratio, specular reflection
2. Inelastic Scattering: Low ratio, light intensity is highly dumped inside the metal. So it does not interact with the net.
3. Absorption: Low ratio, electrons energy gap is too big for visible light.

1. Specular reflection for all the visible light frequencies \rightarrow not colored
2. They are opaques.


1. Specular reflection for all the visible light frequencies → not colored
2. They are opaques.



Structure of non-metallic solids


Structure of non-metallic solids

ionic bond



An idealized ionic (or electrovalent) bonding of oppositely charged ions.


covalent bond




Covalent bonds involve electron sharing, such as between these carbon atoms when they form a diamond.

Structure of non-metallic solids

SALT



DIAMOND



In non-metallic solids:

In non-metallic solids:

- ▶ Atoms have covalent bonds or ionic bonds.

In non-metallic solids:

- ▶ Atoms have covalent bonds or ionic bonds.
- ▶ electrons can not move freely through the material.

In non-metallic solids:

- ▶ Atoms have covalent bonds or ionic bonds.
- ▶ electrons can not move freely through the material.
- ▶ → low electric conductivity, low heat conductivity.

In non-metallic solids:

- ▶ Atoms have covalent bonds or ionic bonds.
- ▶ electrons can not move freely through the material.
- ▶ → low electric conductivity, low heat conductivity.
- ▶ When the material is exposed to an electric field, it interacts with electrons and the ions in the net, they emit EM radiation. The resultant EM radiation is a superposition of both, it is a wave with a slower velocity (refraction law).

In non-metallic solids:

In non-metallic solids:

1. Elastic Scattering: specular reflection

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.
 - ▶ Ionic bonds: The energy needed to heat the net is lower → higher ratio of inelastic scattering.

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.
 - ▶ Ionic bonds: The energy needed to heat the net is lower → higher ratio of inelastic scattering.
 - ▶ covalent bonds: The energy needed to heat the net is higher → lower ratio of inelastic scattering.

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.
 - ▶ Ionic bonds: The energy needed to heat the net is lower → higher ratio of inelastic scattering.
 - ▶ covalent bonds: The energy needed to heat the net is higher → lower ratio of inelastic scattering.
3. Absorption: The ratio depends on the electron energy gap.

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.
 - ▶ Ionic bonds: The energy needed to heat the net is lower → higher ratio of inelastic scattering.
 - ▶ covalent bonds: The energy needed to heat the net is higher → lower ratio of inelastic scattering.
3. Absorption: The ratio depends on the electron energy gap.
 - ▶ High gaps, less absorption → the material is more transparent, no -colored.

In non-metallic solids:

1. Elastic Scattering: specular reflection
2. Inelastic Scattering: The light is transmitted into the material, so it is scattered by the ions on the net. It is diffused.
 - ▶ Ionic bonds: The energy needed to heat the net is lower → higher ratio of inelastic scattering.
 - ▶ covalent bonds: The energy needed to heat the net is higher → lower ratio of inelastic scattering.
3. Absorption: The ratio depends on the electron energy gap.
 - ▶ High gaps, less absorption → the material is more transparent, no -colored.
 - ▶ Lower gaps, more absorption → the material is less transparent, colored.

1. Specular reflection and Diffuse reflection
2. Absorption: color
3. Absorption + Diffusion \rightarrow Low Fresnel coefficient

1. Specular reflection and Diffuse reflection
2. Absorption: color
3. Absorption + Diffusion \rightarrow Low Fresnel coefficient

