

PHY115

Free Falling Motion

Digipen

Spring 2023

Free fall

Free falling bodies

Free fall:

1. Between 1589 and 1592 → Galileo's experiment:
 - ▶ He dropped two spheres of different masses from the Tower of Pisa and demonstrated that their falling time was independent of their mass

Free fall:

1. Between 1589 and 1592 → Galileo's experiment:
 - ▶ He dropped two spheres of different masses from the Tower of Pisa and demonstrated that their falling time was independent of their mass
2. In 1687 → Newton's Laws
 - ▶ He published "*Philosophiæ naturalis principia mathematica*", the laws that describe this experiment.

Video: https://www.youtube.com/watch?v=QyeF-_QPSbk

Free fall: Newton's Law:

The magnitude of the acceleration that the Earth makes on a body is:

$$g = G \frac{M}{d^2}$$

d is the distance between the body and the center of the Earth.

► $G = 6,6710^{-11} \text{ Nm}^2/\text{kg}^2$


Free fall: Newton's Law:

The magnitude of the acceleration that the Earth makes on a body is:

$$g = G \frac{M}{d^2}$$

d is the distance between the body and the center of the Earth.

- ▶ $G = 6,6710^{-11} \text{ Nm}^2/\text{kg}^2$
- ▶ $M = 5,972 \times 10^{24} \text{ kg}$




What happens when the bodies near the Earth surface?

What happens when the bodies near the Earth surface?

$$g = \frac{GM}{(h+R)^2} = \frac{GM}{\underbrace{\left[R\left(\frac{h}{R} + 1\right)\right]^2}_{\approx 0}}$$

$$\rightarrow g \approx \frac{GM}{R} \approx 9.8 \text{ m/s}^2$$



Free fall:

- ▶ Body falling under the influence of the earth's gravitational attraction.

Free fall:

- ▶ Body falling under the influence of the earth's gravitational attraction.
- ▶ If the effects of the air can be neglected, all bodies fall with the same downward acceleration, regardless of their size or weight.

Free fall:

- ▶ Body falling under the influence of the earth's gravitational attraction.
- ▶ If the effects of the air can be neglected, all bodies fall with the same downward acceleration, regardless of their size or weight.
- ▶ If the fall is small compared with the radius of the earth, and if we ignore small effects due to the earth's rotation, the acceleration is constant.

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

$$g = 9.8 \text{ m/s}^2 \quad (1)$$

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

$$g = 9.8 \text{ m/s}^2 \quad (1)$$

Near the surface of the moon:

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

$$g = 9.8 \text{ m/s}^2 \quad (1)$$

Near the surface of the moon:

$$g = 1.6 \text{ m/s}^2 \quad (2)$$

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

$$g = 9.8 \text{ m/s}^2 \quad (1)$$

Near the surface of the moon:

$$g = 1.6 \text{ m/s}^2 \quad (2)$$

Near the surface of the sun:

The constant acceleration of a freely falling body is called the **acceleration due to gravity**, and we denote its magnitude with the letter g .

Near the surface of the earth.

$$g = 9.8 \text{ m/s}^2 \quad (1)$$

Near the surface of the moon:

$$g = 1.6 \text{ m/s}^2 \quad (2)$$

Near the surface of the sun:


$$g = 270 \text{ m/s}^2 \quad (3)$$

g is always a positive number

Because g is the magnitude of a vector quantity, it is always a positive number.


EXAMPLE 3.1

$$x(t) = -\frac{1}{2}gt^2 + v_0t + x_0 \quad (4)$$



EXAMPLE 3.2

$$x(t) = \frac{1}{2}gt^2 + v_0t + x_0 \quad (5)$$



EXERCISE 3.1

EXERCISE 3.1

A one-euro coin is dropped from the Leaning Tower of Pisa and falls freely from rest. What are its position and velocity after 1.0 s, 2.0 s, and 3.0 s?

EXERCISE 3.2

You throw a ball vertically upward from the roof of a building. The ball leaves your hand with an upward speed of 15 m/s ; the ball is then in free fall. Find

EXERCISE 3.2

You throw a ball vertically upward from the roof of a building. The ball leaves your hand with an upward speed of 15 m/s ; the ball is then in free fall. Find

1. the ball's position and velocity 1.00 s and 4.00 s after leaving your hand;

EXERCISE 3.2

You throw a ball vertically upward from the roof of a building. The ball leaves your hand with an upward speed of 15 m/s ; the ball is then in free fall. Find

1. the ball's position and velocity 1.00 s and 4.00 s after leaving your hand;
2. the ball's velocity when it is 5.00 m above the railing;

EXERCISE 3.2

You throw a ball vertically upward from the roof of a building. The ball leaves your hand with an upward speed of 15 m/s ; the ball is then in free fall. Find

1. the ball's position and velocity 1.00 s and 4.00 s after leaving your hand;
2. the ball's velocity when it is 5.00 m above the railing;
3. the maximum height reached;

EXERCISE 3.2

You throw a ball vertically upward from the roof of a building. The ball leaves your hand with an upward speed of 15 m/s ; the ball is then in free fall. Find

1. the ball's position and velocity 1.00 s and 4.00 s after leaving your hand;
2. the ball's velocity when it is 5.00 m above the railing;
3. the maximum height reached;
4. the ball's acceleration when it is at its maximum height.

EXERCISE 3.2






Figure: Figure from Sears and Zemansky's University Physics with Modern Physics, 13th Edition.

Test Your Understanding

Test Your Understanding

If you toss a ball upward with a certain initial speed, it falls freely and reaches a maximum height h a time t after it leaves your hand.

Test Your Understanding

If you toss a ball upward with a certain initial speed, it falls freely and reaches a maximum height h a time t after it leaves your hand.

- ▶ If you throw the ball upward with double the initial speed, what new maximum height does the ball reach?

Test Your Understanding

If you toss a ball upward with a certain initial speed, it falls freely and reaches a maximum height h a time t after it leaves your hand.

- If you throw the ball upward with double the initial speed, what new maximum height does the ball reach?

1. $h\sqrt{2}$
2. $2h$
3. $4h$
4. $8h$
5. $16h$

Test Your Understanding

If you toss a ball upward with a certain initial speed, it falls freely and reaches a maximum height h a time t after it leaves your hand.

Test Your Understanding

If you toss a ball upward with a certain initial speed, it falls freely and reaches a maximum height h a time t after it leaves your hand.

- ▶ If you throw the ball upward with double the initial speed, how long does it take to reach its new maximum height?
 1. $t/2$
 2. $t/\sqrt{2}$
 3. t
 4. $t\sqrt{2}$
 5. $2t$