

Ore's Theorem

CS 111 Presentation

Youssef Adam Thomas Kang
Youssef Koreatam Michael Wong Gabriel Serrano

Department of Computer Science, University of California, Riverside

March 19, 2024

Questions

Introduction: Background Knowledge

To understand **Ore's theorem**, we must briefly recapitulate what it means for a graph to be **Hamiltonian**.

Introduction: Background Knowledge

To understand **Ore's theorem**, we must briefly recapitulate what it means for a graph to be **Hamiltonian**.

Let $G(V, E)$ be an undirected graph with $|V| \geq 3$ vertices.

Introduction: Background Knowledge

To understand **Ore's theorem**, we must briefly recapitulate what it means for a graph to be **Hamiltonian**.


Let $G(V, E)$ be an undirected graph with $|V| \geq 3$ vertices. If we are able to create a cycle T that meets each and every vertex strictly once, then G is considered **Hamiltonian**.

Introduction: Background Knowledge

To understand **Ore's theorem**, we must briefly recapitulate what it means for a graph to be **Hamiltonian**.

Let $G(V, E)$ be an undirected graph with $|V| \geq 3$ vertices. If we are able to create a cycle T that meets each and every vertex strictly once, then G is considered **Hamiltonian**.

Example:





Introduction: Background Knowledge

To understand **Ore's theorem**, we must briefly recapitulate what it means for a graph to be **Hamiltonian**.

Let $G(V, E)$ be an undirected graph with $|V| \geq 3$ vertices. If we are able to create a cycle T that meets each and every vertex strictly once, then G is considered **Hamiltonian**.

Example:




$$T = \{a, b, c, e, d, a\}$$

Introduction: Ore's Theorem Statement

Example I

First Case


$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .



Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .




$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .




$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$

Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$


$$\deg(a) + \deg(e) = 6$$

$$\deg(a) + \deg(f) = 6$$

Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$


$$\deg(a) + \deg(f) = 6$$

$$\deg(b) + \deg(c) = 6$$

Example I

First Case

$G_1(V, E)$ **satisfies** Ore's theorem Ω and **has** a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$

$$\deg(a) + \deg(f) = 6$$


$$\deg(b) + \deg(c) = 6$$

$$\deg(b) + \deg(f) = 6$$

Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$

$$\deg(a) + \deg(f) = 6$$

$$\deg(b) + \deg(c) = 6$$


$$\deg(b) + \deg(f) = 6$$

$$\deg(c) + \deg(e) = 6$$

Example I

First Case

$G_1(V, E)$ **satisfies** Ore's theorem Ω and **has** a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$

$$\deg(a) + \deg(f) = 6$$

$$\deg(b) + \deg(c) = 6$$

$$\deg(b) + \deg(f) = 6$$


$$\deg(c) + \deg(e) = 6$$

$$\vdash \Omega_2$$

Example I

First Case

$G_1(V, E)$ satisfies Ore's theorem Ω and has a Hamiltonian cycle T .



$$|V| = 6 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(e) = 6$$

$$\deg(a) + \deg(f) = 6$$

$$\deg(b) + \deg(c) = 6$$

$$\deg(b) + \deg(f) = 6$$

$$\deg(c) + \deg(e) = 6$$


$$\vdash \Omega_2$$

This graph satisfies Ω and has a Hamiltonian cycle $T = \{d, a, b, e, f, c, d\}$.

Example II

Second Case


$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .



Example II

Second Case

$G_2(V, E)$ **does not satisfy** Ω and **does not have** a Hamiltonian cycle T .




$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

Example II

Second Case

$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .




$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(c) = 4$$

Example II

Second Case

$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .



$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$


$$\deg(a) + \deg(c) = 4$$

$$\deg(a) + \deg(e) = 4$$

Example II

Second Case

$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .



$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(c) = 4$$


$$\deg(a) + \deg(e) = 4$$

$$\deg(b) + \deg(d) = 6$$

Example II

Second Case

$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .



$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(c) = 4$$

$$\deg(a) + \deg(e) = 4$$


$$\deg(b) + \deg(d) = 6$$

$$\deg(c) + \deg(e) = 4$$

Example II

Second Case

$G_2(V, E)$ does not satisfy Ω and does not have a Hamiltonian cycle T .



$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(c) = 4$$

$$\deg(a) + \deg(e) = 4$$

$$\deg(b) + \deg(d) = 6$$


$$\deg(c) + \deg(e) = 4$$

$$\not\vdash \Omega_2$$

Example II

Second Case

$G_2(V, E)$ **does not satisfy** Ω and **does not have** a Hamiltonian cycle T .



$$|V| = 5 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(c) = 4$$

$$\deg(a) + \deg(e) = 4$$

$$\deg(b) + \deg(d) = 6$$

$$\deg(c) + \deg(e) = 4$$


$$\not\vdash \Omega_2$$

This graph **does not satisfy** Ω and **does not have** a Hamiltonian cycle.

Example III

Third Case


$G_3(V, E)$ **does not satisfy** Ω and **has** a Hamiltonian cycle T .



Example III

Third Case

$G_3(V, E)$ **does not satisfy** Ω and **has** a Hamiltonian cycle T .




$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

Example III

Third Case

$G_3(V, E)$ **does not satisfy** Ω and **has** a Hamiltonian cycle T .



$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(d) = 6$$


$$\deg(a) + \deg(e) = 8$$

$$\deg(a) + \deg(f) = 8$$

Example III

Third Case

$G_3(V, E)$ does not satisfy Ω and has a Hamiltonian cycle T .



$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(d) = 6$$

$$\deg(a) + \deg(g) = 4$$

$$\deg(a) + \deg(e) = 8$$

$$\deg(b) + \deg(f) = 8$$


$$\deg(a) + \deg(f) = 8$$

$$\deg(b) + \deg(g) = 8$$

Example III

Third Case

$G_3(V, E)$ does not satisfy Ω and has a Hamiltonian cycle T .



$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(d) = 6$$

$$\deg(a) + \deg(g) = 4$$

$$\deg(c) + \deg(e) = 8$$

$$\deg(a) + \deg(e) = 8$$

$$\deg(b) + \deg(f) = 8$$

$$\deg(c) + \deg(g) = 8$$

$$\deg(a) + \deg(f) = 8$$


$$\deg(b) + \deg(g) = 8$$

$$\deg(d) + \deg(g) = 6$$

Example III

Third Case

$G_3(V, E)$ does not satisfy Ω and has a Hamiltonian cycle T .



$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(d) = 6$$

$$\deg(a) + \deg(g) = 4$$

$$\deg(c) + \deg(e) = 8$$

$$\deg(a) + \deg(e) = 8$$

$$\deg(b) + \deg(f) = 8$$

$$\deg(c) + \deg(g) = 8$$

$$\not\vdash \Omega_2$$

$$\deg(a) + \deg(f) = 8$$


$$\deg(b) + \deg(g) = 8$$

$$\deg(d) + \deg(g) = 6$$

Example III

Third Case

$G_3(V, E)$ does not satisfy Ω and has a Hamiltonian cycle T .



$$|V| = 7 \geq 3 \quad \vdash \quad \Omega_1$$

$$\deg(a) + \deg(d) = 6$$

$$\deg(a) + \deg(e) = 8$$

$$\deg(a) + \deg(f) = 8$$

$$\deg(a) + \deg(g) = 4$$

$$\deg(b) + \deg(f) = 8$$

$$\deg(b) + \deg(g) = 8$$

$$\deg(c) + \deg(e) = 8$$

$$\deg(c) + \deg(g) = 8$$

$$\deg(d) + \deg(g) = 6$$

$\not\vdash \Omega_2$

This graph **does not satisfy** Ω but still **has**
a Hamiltonian cycle $T = \{a, b, d, e, g, f, c, a\}$.

Applications

Conclusion: Summary

Conclusion: Questions