

## Week 49

### Table of contents

|                                                                                   |   |
|-----------------------------------------------------------------------------------|---|
| 1 Difference between $Q$ and $K_{eq}$ .....                                       | 1 |
| 2 Determination of $\Delta H^\circ$ for a reaction from the van't Hoff plot. .... | 2 |
| 3 ATP hydrolysis .....                                                            | 5 |


```
try:
    import fysisk_biokemi
    print("Already installed")
except ImportError:
    %pip install -q "fysisk_biokemi[pyodide] @ https://au-mbg.github.io/fysisk-biokemi/
    wheels/fysisk_biokemi-0.2.2-py3-none-any.whl"
```

---

### 1 Difference between $Q$ and $K_{eq}$

```
import matplotlib.pyplot as plt
import pandas as pd
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

Consider the reaction:



The time course of the reaction was followed monitoring formation of the product  $B$ . The starting concentration of  $A$  was 50  $\mu\text{M}$ , and as you will learn later in the course, the reversible reaction going from  $A$  to  $B$  follows first order kinetics.

#### (a) Load & convert

The data is contained in the `diff_q_keq_data.xlsx` file.

```
data_file_path = ... # Write the path to the dataset file
df = pd.read_excel(data_file_path)
display(df)
```

Convert the measured concentrations of  $B$ ,  $[B]$ , to concentrations in the SI-unit M (mol/L). Repeat for the starting concentration of  $A$ ,  $[A_0]$ .

```
## Your task: Convert the concentration of B to M and set in new column ##
df['[B]_(M)'] = ...

## Your task: Set the initial concentration of A, in the variable A0.
```

```
## Remember to convert to M.
A0 = ...
```

### (b) Law of mass balance

Use the laws of mass balance to calculate  $[A]$  as a function of time in the unit M, add it as a column to the DataFrame

```
# Your task: Calculate the concentration of A for each time-point
## Add it as a column to the dataframe called '[A]_(M)'.
... # Your code here.
```

Make a plot of the time vs. the concentration of  $A$  and of time versus the concentration of  $B$ .

```
fig, ax = plt.subplots()
## Your task: Plot time vs. concentrations. ##
ax.plot(..., ..., label='[A]') # Time vs. concentration of A
ax.plot(..., ..., label='[B]') # Time vs. concentration of B

## This shows the labels. ##
ax.legend()
```

### (c) Mass action ratio

Calculate and plot the mass action ratio  $Q$  as a function of time.

```
## Your task: Calculate the mass action ratio
Q = ... # calculate Q
```

Now plot the mass action ratio versus time.

```
fig, ax = plt.subplots()

## Your task: Make a plot of the axis from the dataframe df['Time_(min)'] vs Q
...
```

### (d) $Q$ vs $K_{eq}$

What is the difference between  $Q$  and  $K_{eq}$ ?

### (e) Favoured compound

Which of the two compounds  $A$  and  $B$  is favored at equilibrium? (Favored: highest concentration)

### (f) Equilibrium constant

What is the equilibrium constant of this reaction?

## 2 Determination of $\Delta H^\circ$ for a reaction from the van't Hoff plot.

```
import pandas as pd
import matplotlib.pyplot as plt
from scipy.optimize import curve_fit
import numpy as np
from IPython.display import display
```

The equilibrium constant for a reversible reaction going from  $A$  to  $B$  was measured as a function of temperature. The data obtained is given in the Excel document `deter_delta_h_data.xlsx`. Load the data with pandas.

```
data_file_path = ... # Write the path to the dataset file
df = pd.read_excel(data_file_path)
display(df)
```

### (a) Transform & plot the data



Tip

In NumPy the `np.log` function calculates the natural logarithm that we often write as  $\ln$  and `np.log10` calculates the base 10 logarithm.

Add new columns to the DataFrame containing  $\ln(K_{eq})$  and  $1/T$  and make a plot with  $1/T$  on the x-axis and  $\ln(K_{eq})$  on the y-axis.

```
## Your task: Add a column containing 1/T and one containing ln(Keq) ##
df['1/T'] = ...
df['ln_Keq'] = ...
```

Now you can make the plot in the cell below

```
fig, ax = plt.subplots()

## Your task: Make a plot of 1/T vs ln(Keq)
...

## EXTRA: Customization you don't need to edit or understand this
## It adds axis labels to the plot you made.
ax.set_xlabel(r'1/T [K]')
ax.set_ylabel(r'$\ln(K_{eq})$');
```

### (b) Fit trendline

Make a linear trendline by fitting a linear function to the data.

Start by defining a linear function

```
def vant_hoff(T, a, b):
    ## Your task: Implement a linear function
```

```

result = ...
return result

```

Now we can use regression to find the parameters

```

## Your task: Make a fit with curve_fit to find the parameters ##
fitted_parameters, trash = ...
a_fit, b_fit = fitted_parameters

# Print
print(a_fit)
print(b_fit)

```

Then make a plot with the trendline added

```

fig, ax = plt.subplots()

## Your task: Choose points to evaluate the fit
## Look at the x-axis above and estimate a value for the minimum and maximum.
## Then calculate the fit at those points
max_x = ...
min_x = ...
T_trend = np.array([min_x, max_x]) # Select or make some appropriate points for the
plotting evaluating the trendline.
y_fit = vant_hoff(T_trend, ..., ...)

## Your task: Plot the fit
...

## Plots the data
ax.plot(df['1/T'], df['ln(Keq)'], 'o')

## EXTRA: You do not need to edit or understand this.
## It adds axis labels to the plot you made.
ax.set_xlabel('1/T [K]')
ax.set_ylabel(r'$\ln(K_{eq})$');

```

### (c) Reaction type

Calculate the reaction enthalpy and entropy, at  $T = 298$  K, assuming that  $R = 8.314472 \times 10^{-3} \frac{\text{kJ}}{\text{mol} \cdot \text{K}}$ . Is the reaction endo- or exothermic?

```

## Your task: Set known values and add units as comments e.g. # kJ/(mol K) ##
R = ...
T = ...

## Your task: Calculate ΔH and ΔS ##
## Consider the units!
dH = ...
dS = ...

```

```
print(dH)
print(dS)
```

#### 💡 Tip

The units of a slope is the units of the y-axis divided by the units of the x-axis.

In the van't Hoff plot the y-axis is unitless and the x-axis is in 1/K.

#### (d) Free energy change

Calculate the free energy change,  $\Delta G^\circ$ , at  $T = 298 \text{ K}$ , assuming that  $R = 8.314472 \times 10^{-3} \frac{\text{kJ}}{\text{mol}\cdot\text{K}}$

```
## Your task: Calculate ΔG
dG = ...


print(dG)
```

### 3 ATP hydrolysis

```
import numpy as np
```

Below is shown the structure of ATP.

$\Delta G^\circ$  for the hydrolysis of ATP to ADP and Pi is  $-30.5 \text{ kJ/mol}$ .



(a)  $\Delta G$

In cells at 298 K the steady state concentrations of ATP, ADP og Pi resulted in a value for the reaction quotient  $Q = \frac{[ADP][Pi]}{[ATP]} = 0.002$ . Calculate  $\Delta G'$  under these conditions.

```
## Your task: Define known values
## Add units as comments! E.g. # kJ / mol
dG_std = ... # kJ / mol
Q = ...
T = ...
... # One more might be useful.

## Your task: Calculate dG
dG = dG_std + R*T*np.log(Q)
print(dG)
```

### (b) Equilibrium or not?

Assuming that the temperature within these cells is 298 K, are these cells in equilibrium with respect to the hydrolysis of ATP?

```
## Your task: Calculate K_eq.
...
```

### (c) ATP synthesized

Calculate the number of moles of ATP synthesized by a resting person in a period of 24 hours, assuming that the energy consumption by this person is 8368 kJ per 24 hours and that 40% of this energy is converted directly into ATP without loss of energy. Use the  $\Delta G'$  value calculated in question a, and assume that 298 K.

```
# Your task: Set the known value of the energy spent for ATP.
energy_atp = ...

# Your task: Calculate the number of moles of ATP synthezied
synth_atp = ...
print(synth_atp)
```



Tip

You can use `np.absolute` to calculate the absolute value of a number.

### (d) Turnover

Next, calculate the average number of times each ATP molecule is turned over (synthesis followed by hydrolysis) in 24 hours when the total amount of ATP in the body equals 0.2 mol.

```
## Your task: Set the known total_atp value
## Remember unit
...

## Your task: Calculate the turnover number.
```

```
... # Your calculation here
```

```
## Your task: Print the calculated turnover number.
```