

Week 44

Table of contents

1 Sekvens egenskaber	1
2 Fasta til DataFrame	1
3 Self-study: Calculations in python	1

```
try:
    import fysisk_biokemi
    print("Already installed")
except ImportError:
    %pip install -q "fysisk_biokemi[pyodide] @ https://au-mbg.github.io/fysisk-biokemi/
    wheels/fysisk_biokemi-0.2.2-py3-none-any.whl"
```

```
import fysisk_biokemi.widgets as fsb_widgets
```

1 Sekvens egenskaber

```
fsb_widgets.sequence_properties()
```

2 Fasta til DataFrame

```
from fysisk_biokemi.widgets.sequence_properties import sequence_dataframe

widget = sequence_dataframe()
```

DataFrame'en ved den valgte pH kan trækkes ud

```
df = widget.get_dataframe()
```

3 Self-study: Calculations in python

```
import numpy as np
```

In the following, we will go through the some common mathematical manipulations of number, arrays and data frames using python. You should have covered most of these in "Introduction to the Molecules of Life", so we will not go through them in class.

(a) Simple calculations

Evaluate the following expressions using Python

- 10.5×4
- $5 + 10 + 15 + 12$

- $70 - 30 + 2$
- $420/10$

```
... # Replace ... with your code
...
...
...
```

💡 Tip

Recall that the basic arithmetic operations are, in Python, represented as follows

- $+$ $\Rightarrow$ `+`
- $-$ $\Rightarrow$ `-`
- $/$ $\Rightarrow$ `/`
- $\times$ $\Rightarrow$ `*`
- x^y $\Rightarrow$ `x**y`

(b) Array calculations

The ability to perform operations on all elements (or data points) is incredibly useful when doing data analysis, for example when converting units. This is enabled by array calculations.

The cell below defines an array a with all the numbers from 0 to 9.

For each entry a_i in the array calculate the expression $b_i = a_i/3 + a_i^2$ and assign that to a new array b . Use the fact that you can do elementwise operations with an array - do not calculate it separately for every element of the array.

```
a = np.arange(0, 10)
```

```
b = ...
print(b)
```

💡 Tip

With an array all the basic arithmetic operations can be done elementwise, so the expression $x^2 + 1$ can be done for all elements in an array as such

```
x = np.arange(0, 10)
y = x**2 + 1
```

Just like if x was just a single number. Similarly, expressions can involve adding arrays together like

```
y = x**2 + x
```

(c) Variables

Variables are an essential part of a program. Calculate the following where each intermediate is assigned to a variable

$$\begin{aligned}a &= 5 \times 3 \\b &= a + 9 \\c &= \frac{4}{3}a + b - 2\end{aligned}$$

```
a = ...  
b = ...  
...  
print(a, b, c)
```

Bonus question: Why does c print as a decimal number when the others print without decimals?

(d) Parentheses

Parentheses are important, a misplaced or lacking parentheses can drastically change a calculation.

Using the variables a, b and c you calculate in the previous exercise, evaluate the following expressions

- $a \times (b + c)$
- $a \times b + c$
- $\frac{a+b+c}{b+c}$
- $(b - a)\left(\frac{1}{c} + 1\right)$
- $(b - a)\left(\frac{1}{c+1} + 1\right)$

```
q1 = ...  
q2 = ...  
q3 = ...  
q4 = ...  
q5 = ...  
print(q1, q2, q3, q4, q5)
```



Tip

If you're in doubt having an extra set of unnecessary parentheses is better than missing a necessary set of parentheses. Just be careful to place them correctly.

(f) Functions

For repeated computations it's good practice to define a function.

Define a function that calculates the following expression

$$f(x) = \frac{Ax + B + C}{A + B}$$

Where $A = 10$, $B = 5$, $C = 2.5$

```
def fun_function(x):  
    result = ...  
    return result
```

Now evaluate the function for $x = 62.5$, $x = 629.25$, $x = 42$ and $x = 2025$

```
fun_output_1 = fun_function(...) # Your code replaces ...  
fun_output_2 = ... # Same here  
fun_output_3 = ... # And here  
fun_output_4 = ... # And finally here.  
  
print(f"{fun_output_1:.1f}")  
print(f"{fun_output_2:.1f}")  
print(f"{fun_output_3:.1f}")  
print(f"{fun_output_4:.1f}")
```

Tip

The input of a function will often be the independent variable and the functions calculates the independent variable.

Later when we get to regression/fitting this will be the case.
