

Week 47

v1.0.1

Table of contents

1 A tricky protein binding.	2
2 Determination of reaction orders.	9
3 Analysis of a reversible reaction	15
4 Determination of reaction order and activation energy	19
5 The pH effect on an active site histidine residue	27
6 Qualitative description of Michaelis-Menten kinetics	30
7 Design of an enzyme kinetics experiment	31
8 Lab Exercise Analysis: Michaelis Menten	40

```
try:
    import fysisk_biokemi
    print("Already installed")
except ImportError:
    %pip install -q "fysisk_biokemi[colab] @ git+https://github.com/au-mbg/fysisk-biokemi.
    git#subdirectory=course-utils"
```

1 A tricky protein binding.

```
import numpy as np
import matplotlib.pyplot as plt
import pandas as pd
from scipy.optimize import curve_fit
from fysisk_biokemi.widgets import DataUploader
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

Here, we will analyse the results from an experiment used to quantify affinity of a protein:lignd interaction. The data in `binding_data_sq.xlsx` is in the form of saturation (theta) as a function of the total ligand concentration `L_tot`. This is the type of data you typically get from e.g. a fluorescence titration experiment. Each measurement has been repeated three times.

Load the dataset as usual using the widget below

Run the next cell **after** uploading the file



Tip

The command `display(df)` shows the dataframe that you've loaded.

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('binding_data_sq') # Load from package for the solution so it doesn't
require to interact.
display(df)
```

	L_(uM)	theta
0	0.0	0.000000
1	0.0	0.005258
2	0.0	0.000000
...	...	...
72	500.0	0.875167
73	500.0	0.873289
74	500.0	0.854871

(a) Convert to SI

Convert the ligand concentration to M and add it to the dataframe as a new column named '`L_(m)`'.

You can use `display(df)` to check if you have been successful.

```
df['L_(M)'] = df['L_(uM)'] * 10**(-6)
display(df)
```

	L_(uM)	theta	L_(M)
0	0.0	0.000000	0.0000
1	0.0	0.005258	0.0000
2	0.0	0.000000	0.0000
...	...	...	...
72	500.0	0.875167	0.0005
73	500.0	0.873289	0.0005
74	500.0	0.854871	0.0005


(b) Plot

Make a scatter plot of the dataset and use the plot to estimate K_D .

```
fig, ax = plt.subplots()

## Your task: Make a scatter plot (just points, no lines) of the data. ##
ax.plot(df['L_(M)'], df['theta'], 'o')

## EXTRA: ##
## This adds x and y-axis labels. ##
ax.set_xlabel('L (M)')
ax.set_ylabel(r'$\theta$')
plt.show()
```



Put your estimate of K_D in the cell below

```
## Your task: Assign your estimate of K_D to the variable below ##
K_D_estimate = 0.0001
```

(c) Fit and analyze.

Fit the data using the simple binding expression

$$\theta = \frac{[L]}{[L] + K_D}$$

Start by defining a function, called `simple_binding`, that calculates this expression

```
## Your task: Define the function simple_binding that evaluates the equation ##
## from above. ##
def simple_binding(L, K_D):
    theta = L / (L + K_D)
    return theta
```

Then use that function to fit the data using `curve_fit`

```
## Your task: Set your estimate of K_D as the initial guess.
initial_guess = [K_D_estimate]

## Your task: Use curve_fit to find the fitted value of K_D
fitted_parameters, trash = curve_fit(simple_binding, df['L_(M)'], df['theta'],
initial_guess)

## This extracts K_D and prints it ##
K_D_fit_simple = fitted_parameters[0]
print(K_D_fit_simple)
```


```
0.00013628483915769865
```

And evaluate the fit and plot it together with the dataset

```
## Your task: Evaluate simple_binding using L_smooth and the value of K_D from your fit ##
L_smooth = np.linspace(0, 0.0005, 100)
theta_fit_simple = simple_binding(L_smooth, K_D_fit_simple)

fig, ax = plt.subplots()
## Your task: Plot the fit as line plot and the data as a scatter plot. ##
ax.plot(df['L_(M)'], df['theta'], 'o')
ax.plot(L_smooth, theta_fit_simple)

## EXTRA:
## Labels for the axis.
ax.set_xlabel('L (M)')
ax.set_ylabel(r'$\theta$')
plt.show()
```



Is it a good fit?

(d) Residuals

To get a more quantitative view of the quality of the fit calculate residuals

```
theta_model = simple_binding(df['L_(M)'], K_D_fit_simple)
residuals_simple = df['theta'] - theta_model
```

And plot them versus the measured [L]


```
fig, ax = plt.subplots()

## Your task: Make a plot with residuals on the y-axis and L on the x-axis ##
ax.plot(df['L_(M)'], residuals_simple, 'o')

## This makes a horizontal line through 0 ##
ax.axhline(0, color='black', linestyle='--')

## EXTRA
## Adds axis labels.
ax.set_xlabel('[L] (M)')
ax.set_ylabel('Residual')
```

```
Text(0, 0.5, 'Residual')
```



Are the residuals randomly distributed around zero? What does this tell you about the quality of the fit?

(e) Formulate a hypothesis

The protein concentration in the above titrations was 250 μM . Formulate a hypothesis for why the fit fails.

! Answer

The assumption made to derive the simple binding model is that the ligand concentration is much larger than the protein concentration such that the $[L_{\text{free}}] \approx [L_{\text{total}}]$ - or in other words that the number of ligand molecules bound to the proteins are not a significant proportion of the total number of ligand molecules. With $[P] = 250 \mu\text{M}$ and ligand concentrations in the range of 0 to 500 μM this assumption is not valid.

(f) Choosing a better model

Next, we will try a fitting model that explicitly considers the concentration of the protein being titrated. This is known as the quadratic binding equation

$$\theta = \frac{K_D + [P_{\text{tot}}] + [L_{\text{tot}}]}{2[P_{\text{tot}}]} - \sqrt{\left(\frac{K_D + [P_{\text{tot}}] + [L_{\text{tot}}]}{2[P_{\text{tot}}]}\right)^2 - \frac{[L_{\text{tot}}]}{[P_{\text{tot}}]}}$$

The value of $[P_{\text{tot}}]$ is 250 μM .

Start by implementing the expression as a Python function - remember that you need to be careful with parentheses with a function like this.

```
def quadratic_binding(L, K_D):
    ## Your task: Set the protein concentration in appropriate units. ##
    P_tot = 250 * 10**(-6)
```

```

## Your task: Implement the quadratic binding equation ##
## Be careful with parentheses! ##
theta = (K_D + P_tot + L) / (2 * P_tot) - np.sqrt(((K_D + P_tot + L) / (2 * P_tot))**2
- L / P_tot)
return theta

print(f"{quadratic_binding(100, 138.09) = :.3f}") # Should give 0.420

```

```
quadratic_binding(100, 138.09) = 0.420
```

And now make a fit using this expression

```

## Your task: Make a fit using the quadratic binding equation. ##
fitted_parameters, trash = curve_fit(quadratic_binding, df['L_(M)'], df['theta'])

## Extratcs and prints K_D ##
K_D_fit_quad = fitted_parameters[0]
print(K_D_fit_quad)

```

```
4.214239381928562e-05
```

The cell below makes a figure of the data and the residuals for each fit. You are not expected to understand every line of code – the figure is mainly to aid you interpretation of the data and show case how you can make beautiful figures from you own data later on.

```

theta_fit_quad = quadratic_binding(L_smooth, K_D_fit_quad)
residuals_quad = df['theta'] - quadratic_binding(df['L_(M)'], K_D_fit_quad)

fig, axes = plt.subplots(1, 2, figsize=(9, 4))


# Data & Fit
ax = axes[0]
ax.plot(df['L_(M)'], df['theta'], 'o', label='Data')
ax.plot(L_smooth, theta_fit_simple, label='Simple binding', linewidth=3)
ax.plot(L_smooth, theta_fit_quad, label='Quadratic binding', linewidth=3)
ax.set_xlabel('L (M)')
ax.set_ylabel(r'$\theta$')
ax.set_title('Data & Fits')
ax.legend()

# Residuals
ax = axes[1]
ax.plot(df['L_(M)'], residuals_simple, 'o', color='C1', label='Simple binding')
ax.plot(df['L_(M)'], residuals_quad, 'o', color='C2', label='Quadratic binding')
ax.axhline(0, color='black', linestyle='--')
ax.set_title('Residuals')
ax.set_xlabel('L (M)')

```

```
ax.legend()
```

```
plt.show()
```



(g) Analyze

Consider the following questions:

- How does the quadratic binding expression fit the data?
- What's the difference in how the residuals are distributed?
- What is the biochemical situation that makes quadratic expression necessary?

! Answer

- The quadratic binding equation fits the data very well!
- For the quadratic binding equation the residuals are randomly distributed around zero, whereas for the simple binding model there is a trend as a function of the ligand concentration with negative residuals at low concentrations and positive residuals at higher concentrations.
- That the protein and ligand concentrations are comparable - making the assumptions that the simple binding model is based on invalid.

2 Determination of reaction orders.

```
import matplotlib.pyplot as plt
import numpy as np
import pandas as pd
from scipy.optimize import curve_fit
from fysisk_biokemi.widgets import DataUploader
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

You have carried out a kinetic experiment investigating the turnover of two metabolites, A1 and A2, in a cell lysate. For each metabolite, you have measured the concentration of the compound as a function of time using UV-Vis spectroscopy aiming to determine the reaction order, and thus deduce something about the reaction mechanism by which they are processed. The data from this experiment is found in `reaction-orders.xlsx`.

(a) Load the dataset

Load the dataset `reaction-orders.xlsx` using the widget below

Run the next cell **after** uploading the file

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('reaction_order_week48') # Load from package for the solution so it
doesn't require to interact.
display(df)
```

	Time_s	A1_uM	A2_uM
0	0.0	2.000000	4.500000
1	1.0	1.833259	4.212598
2	2.0	1.823775	3.939486
...	...	...	...
48	48.0	0.136936	0.854009
49	49.0	0.132587	0.885128
50	50.0	0.212870	0.752202

(b) SI units.

Add two new columns with the concentrations given in M.

```
df['A1_M'] = df['A1_uM'] * 10**(-6)
df['A2_M'] = df['A2_uM'] * 10**(-6)
display(df)
```

	Time_s	A1_uM	A2_uM	A1_M	A2_M
0	0.0	2.000000	4.500000	2.000000e-06	4.500000e-06

	Time_s	A1_uM	A2_uM	A1_M	A2_M
1	1.0	1.833259	4.212598	1.833259e-06	4.212598e-06
2	2.0	1.823775	3.939486	1.823775e-06	3.939486e-06
...	...	...	...	...	...
48	48.0	0.136936	0.854009	1.369359e-07	8.540093e-07
49	49.0	0.132587	0.885128	1.325872e-07	8.851275e-07
50	50.0	0.212870	0.752202	2.128700e-07	7.522017e-07

(c) Plotting


For each reactant make a plot of the concentration versus time.

```
fig, ax = plt.subplots()

## Your task: Make plots of concentration (y) versus time (x). ##
## Hint: ax.plot(x_data, y_data, 'o') to make a scatter plot. ##
ax.plot(df['Time_s'], df['A1_M'], 'o')
ax.plot(df['Time_s'], df['A2_M'], 'o')

## EXTRA: Customization of axis labels ##
ax.set_xlabel('Time [s]')
ax.set_ylabel('Concentration')
```

```
Text(0, 0.5, 'Concentration')
```



(d) Model as N'th-order reaction

We will model the reactions as zeroth, first and second order. We will consider both the rate constant k and the concentration at time $t = 0$ parameters of the fit.

```
def zeroth_order(t, k, A0):
    ## Your task: Implement the zeroth order rate equation ##
    result = A0 - k*t
    return result

def first_order(t, k, A0):
    ## Your task: Implement the first order rate equation ##
    result = A0 * np.exp(-k*t)
    return result

def second_order(t, k, A0):
    ## Your task: Implement the second order rate equation ##
    result = A0 / (1 + 2*k*t*A0)
    return result

## Use these to check that your functions are correct!
print('zeroth_order', zeroth_order(1, 1, 2)) # Should give 1
print('first_order', first_order(1, 1, 2)) # Should give 0.7357...
print('second_order', second_order(1, 1, 2)) # Should give 0.4
```

```
zeroth_order 1
first_order 0.7357588823428847
second_order 0.4
```

Then we can make a fit to measurements of each reactant.

```
## Your task: Pick a rate equation and assign it to the variable rate_equation
## You can change this to first_order or second_order
## Do not call the function - so no (...) just assign it to the variable.
rate_equation = zeroth_order

## Your task: Make a fit of the A1_M data with 'rate_equation'.
fitted_parameters_A1, trash = curve_fit(rate_equation, df['Time_s'], df['A1_M'])
k_A1_fit = fitted_parameters_A1[0]
A1_0_fit = fitted_parameters_A1[1]

## Your task: Make a fit of the A2_M with 'rate_equation'.
fitted_parameters_A2, trash = curve_fit(rate_equation, df['Time_s'], df['A2_M'])
k_A2_fit = fitted_parameters_A2[0]
A2_0_fit = fitted_parameters_A2[1]

## Prints the found parameters
print(k_A1_fit)
print(A1_0_fit)

print(k_A2_fit)
print(A2_0_fit)
```

```
3.390184293164445e-08
1.5887604659704164e-06
5.786947687397643e-08
3.148604267116672e-06
```

Now that we fitted the data using one of the rate equations, we can plot to see how well it fits

```
fig, ax = plt.subplots()


## Your task: Finish the code to calculate the fit for A2.
t_smooth = np.linspace(0, 55)
A1_fit = rate_equation(t_smooth, k_A1_fit, A1_0_fit)
A2_fit = rate_equation(t_smooth, k_A2_fit, A2_0_fit)

## Your task: Plot the fits
ax.plot(t_smooth, A1_fit)
ax.plot(t_smooth, A2_fit)

## Plots the data ##
ax.plot(df['Time_s'], df['A1_M'], 'o')
ax.plot(df['Time_s'], df['A2_M'], 'o')

## EXTRA: Customization of axis labels ##
ax.set_xlabel('Time [s]')
ax.set_ylabel('Concentration')
```

```
Text(0, 0.5, 'Concentration')
```




Once you've finished the fitting with one of the rate equations you can go back and change the `rate_equation` variable to try one of the others.

(e) Determining reaction orders

The cell below makes a plot for each rate law in the same panel using your defined rate equations.

```
from fysisk_biokemi.utils.deter_reacti_orders import make_plot
rate_laws = {0: zeroth_order, 1: first_order, 2: second_order}
make_plot(df, rate_laws)
```



Based on this plot;

1. What is the reaction order for [A1]?
2. What is the unit of the rate constant for [A1]?
3. What is the reaction order for [A2]?
4. What is the unit of the rate constant for [A2]?

! Answer

1. First order
2. The unit of the rate constant for a first order reaction is $\frac{1}{s}$

The first order rate equation is

$$A_0 \cdot \exp(-k \cdot t)$$

The term in the exponential has to be unitless, so we need k to have the inverse units of t

$$k \rightarrow \frac{1}{t} \rightarrow \frac{1}{s}$$

3. Second order
4. The unit of the rate constant for a second order reaction is $\frac{1}{s \cdot M}$

The integrated second order rate is

$$\frac{A_0}{1 + 2 \cdot k \cdot t \cdot A_0}$$

Looking at the denominator it includes the term

$$1 + 2 \cdot k \cdot t \cdot A_0$$

For this addition to make sense the second term needs to be unitless, so we can conclude that the unit of k must cancel those of $t \cdot A_0$, so

$$k \rightarrow \frac{1}{t \cdot A_0} \rightarrow \frac{1}{s \cdot M}$$

(f) Reaction mechanism


What does the above tell you about the reaction mechanism whereby the two compounds disappear from the lysate?

! Answer

3 Analysis of a reversible reaction

```
import matplotlib.pyplot as plt
import numpy as np
```

Consider the following reaction



The magnitudes of the rate constants are $k_1 = 10 \text{ s}^{-1}$ and $k_2 = 1 \text{ s}^{-1}$.

(a) Reaction order

What is the reaction order in each direction?

! Answer

Two perspectives

From the chemical equation: Each elementary step involves only one reactant molecule:

- Forward reaction: $A \rightarrow B$ (one molecule of A)
- Reverse reaction: $B \rightarrow A$ (one molecule of B)

Since each step involves a single reactant, both are first-order reactions.

From the units: The rate constants have units of s^{-1} , which is the unit of first order reactions.

(b) Derive equilibrium constant

Show mathematically how the equilibrium constant K_{eq} is given by the ratio between the two rate constants.

! Answer

At equilibrium, the forward and reverse reaction rates are equal:

$$r_{\text{forward}} = r_{\text{reverse}}$$

For the reaction $A \rightleftharpoons B$, both are first order as discussed above:

- Forward rate: $r_{\text{forward}} = k_1[A]_{\text{eq}}$
- Reverse rate: $r_{\text{reverse}} = k_2[B]_{\text{eq}}$

Setting them equal:

$$k_1[A]_{\text{eq}} = k_2[B]_{\text{eq}}$$

Rearranging to solve for the ratio of concentrations:

$$\frac{[B]_{\text{eq}}}{[A]_{\text{eq}}} = \frac{k_1}{k_2}$$

By definition, the equilibrium constant is:

$$K_{\text{eq}} = \frac{[B]_{\text{eq}}}{[A]_{\text{eq}}}$$

Therefore:

$$K_{\text{eq}} = \frac{k_1}{k_2}$$

(c) Calculate K_{eq}

```
## Your task: Assign known values & calculate the equilibrium constant.  
k1 = 10  
k2 = 1  
K_eq = k1 / k2  
print(K_eq)
```

10.0

(d) Equilibrium concentrations

Calculate the concentrations of A and B at equilibrium, $[A]_{\text{eq}}$ and $[B]_{\text{eq}}$, if $[A]_0 = 10^{-3}\text{M}$

```
A0 = 10**(-3)  
A_eq = (k2 * A0) / (k1 + k2)  
B_eq = (k1 * A0) / (k1 + k2)
```



```
print(A_eq)
print(B_eq)
```

```
9.090909090909092e-05
0.0009090909090909091
```

(e) Initial rate of formation


If $[A]_0 = 10^{-3}$ M and $[B]_0 = 0$ M, calculate the initial rate of formation of B.

```
## Your task: Calculate initial rate of formation of B
r_fwd = A0 * k1
print(r_fwd)
```

```
0.01
```

(f) Time-dependence

$$\begin{matrix} k_f = k_1 \\ k_b = k_{-1} \end{matrix}$$



$$[B]_{eq} = \frac{k_1}{k_1 + k_{-1}} [A]_0$$

$$[A]_{eq} = \frac{k_{-1}}{k_1 + k_{-1}} [A]_0$$

$$[A]_0 = [A] + [B] \quad [B]_0 = 0$$

$$[A] = \frac{k_b [A]_0}{k_f + k_b} \left[1 + \frac{k_f}{k_b} e^{-(k_f + k_b)t} \right] = \frac{k_b + k_f e^{-(k_f + k_b)t}}{k_f + k_b} [A]_0$$

We now want to calculate and plot the time-dependent concentrations using the above equations.

In the cell below finish the implementation of the function `A_time` that calculates the concentration $[A]$ as a function of time.

```
## Your task: Implement the function to calculate the time dependent concentration.
## Be careful with parentheses.
def A_time(t, A0, k_f, k_b):
    A_t = (k_b + k_f * np.exp(-(k_f+k_b)*t)) / (k_f + k_b) * A0
    return A_t
```

```
test = A_time(1, 1, 1, 1)
print(test) # This should give 0.567...
```

```
0.5676676416183064
```


And then we can use that function to calculate and plot the concentrations as a function of time. As you have seen above, the reaction is pretty fast, so we will just focus on the first second

```
fig, ax = plt.subplots()
t = np.linspace(0, 1, 100)

## Your task: Calculate [A](t) and [B](t) ##
At = A_time(t, A0, k1, k2)
Bt = A0 - At

## Plots the concentrations versus time.
ax.plot(t, At, label='[A](t)')
ax.plot(t, Bt, label='[B](t)')

## EXTRA: Sets x and y-axis labels.
ax.set_ylabel('Concentrations (M)', fontsize=14)
ax.set_xlabel('Time (s)', fontsize=14)
ax.legend()
```



4 Determination of reaction order and activation energy

```
import numpy as np
import matplotlib.pyplot as plt
import pandas as pd
from fysisk_biokemi.widgets import DataUploader
from IPython.display import display
from scipy.optimize import curve_fit
pd.set_option('display.max_rows', 6)
```

The irreversible isomerization of compound A to compound B results in a decreasing absorbance. The isomerization was followed in a time course at two different temperatures ($T_1 = 25\text{ °C}$ and $T_2 = 40\text{ °C}$). The absorbance ($\epsilon = 16700\text{ cm}^{-1}\text{M}^{-1}$) was used to calculate the concentration of compound A in a spectrophotometer with a pathlength of 1 cm. The obtained dataset is given in the file `deter-reacti-order-activ.csv`.

(a) Temperatures

What are the two temperatures in Kelvin? Set them as variables in the cell below.

```
## Your task: Convert the temperatures to K and assign to the variables:
T1 = 25 + 273.15
T2 = 40 + 273.15
```

(b) Load the dataset

Load the dataset using the widget below

```
uploader = DataUploader()
uploader.display()
```

Run the next cell **after** uploading the file

```
df = uploader.get_dataframe()
display(df)
```

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('reaction_order_activation_week48') # Load from package for the solution
so it doesn't require to interact.
display(df)
```

	t_(s)	Abs(t)_25C	Abs(t)_40C
0	0.000	0.835	0.835
1	0.002	0.801	0.806
2	0.004	0.787	0.731
...	...	...	...
23	0.046	0.327	0.207

	t_(s)	Abs(t)_25C	Abs(t)_40C
24	0.048	0.321	0.201
25	0.050	0.306	0.185

(c) SI Units

Calculate the concentration of A at each timepoint in SI units, by adding new columns to the DataFrame.

```
## Your task: Assign known values ##
extinction_coeff = 16700
L = 1 # Path length

## Calculate concentrations using the variables above and the data in the dataframe ##
df['[A]_(M)_25C'] = df['Abs(t)_25C'] / (extinction_coeff * L)
df['[A]_(M)_40C'] = df['Abs(t)_40C'] / (extinction_coeff * L)
display(df)
```

	t_(s)	Abs(t)_25C	Abs(t)_40C	[A]_(M)_25C	[A]_(M)_40C
0	0.000	0.835	0.835	0.000050	0.000050
1	0.002	0.801	0.806	0.000048	0.000048
2	0.004	0.787	0.731	0.000047	0.000044
...	...	...	...	...	...
23	0.046	0.327	0.207	0.000020	0.000012
24	0.048	0.321	0.201	0.000019	0.000012
25	0.050	0.306	0.185	0.000018	0.000011


(d) Plot the data

We will be resuing the plot, so we will put the code for it in a function.

```
fig, ax = plt.subplots()

# Task: Plot the two data series in a scatter plot
# That is: t_(s) vs [A]_(M)_25C
# and: t_(s) vs [A]_(M)_40C
ax.plot(df['t_(s)'], df['[A]_(M)_25C'], 'o', label='[A]_25C')
ax.plot(df['t_(s)'], df['[A]_(M)_40C'], 'o', label='[A]_40C')

## EXTRA: This sets x and y-axis labels and shows the legends.
ax.set_xlabel('t (s)')
ax.set_ylabel('[A] (M)')
ax.legend()
```



(e) Fitting with rate equations

Now want to fit the data using integrated rate laws - in order to ultimately determine the reaction orders of the two datasets.

To do so we need functions for zeroth, first and second order integrated rate laws.

```
def zeroth_order(t, k, A0):
    ## Your task: Implement the zeroth order rate equation ##
    result = A0 - k*t
    return result

def first_order(t, k, A0):
    ## Your task: Implement the first order rate equation ##
    result = A0 * np.exp(-k*t)
    return result

def second_order(t, k, A0):
    ## Your task: Implement the second order rate equation ##
    result = A0 / (1 + 2*k*t*A0)
    return result

## Use these to check that your functions are correct!
print('zeroth_order', zeroth_order(1, 1, 2)) # Should give 1
print('first_order', first_order(1, 1, 2)) # Should give 0.7357...
print('second_order', second_order(1, 1, 2)) # Should give 0.4
```

```
zeroth_order 1
first_order 0.7357588823428847
second_order 0.4
```

Having defined these functions we can use them for fitting

```
## Task: Choose one of the rate equations and assign to the variable
rate_equation = first_order

## Task: Fit to the data at 25 C
fitted_parameters_25C, trash = curve_fit(rate_equation, df['t_(s)'], df['[A]_(M)_25C'])

## Task: Fit to the data at 40 C
fitted_parameters_40C, trash = curve_fit(rate_equation, df['t_(s)'], df['[A]_(M)_40C'])

## Extracts & prints the parameters
k_25C, A0_25C = fitted_parameters_25C
k_40C, A0_40C = fitted_parameters_40C

print(k_25C, A0_25C)
print(k_40C, A0_40C)
```

```
20.081750703221886 5.024371023346721e-05
29.960916302810904 4.990085393325908e-05
```

Having made the fits we plot it together with the two data series.


```
## Your task: Calculate the fits using the chosen rate equation &
## the found parameters found during fitting.
t_smooth = np.linspace(0, 0.05, 100)
A_fit_25C = rate_equation(t_smooth, k_25C, A0_25C)
A_fit_40C = rate_equation(t_smooth, k_40C, A0_40C)

fig, ax = plt.subplots()

## Your task: Plot the two fits
ax.plot(t_smooth, A_fit_25C)
ax.plot(t_smooth, A_fit_40C)

## This plots the data sets
ax.plot(df['t_(s)'], df['[A]_(M)_25C'], 'o', label='[A]_25C', color='C0')
ax.plot(df['t_(s)'], df['[A]_(M)_40C'], 'o', label='[A]_40C', color='C1')

## EXTRA: Sets the axis labels and legend.
ax.set_xlabel('t (s)')
ax.set_ylabel('[A] (M)')
ax.legend()
```




You can go back to the cell where you set `rate_equation` and use another of the three: `zeroth_order`, `first_order`, `second_order`.

(f) Determine reaction order

Based on your analysis above or by using the plot the created by the next cell determine the reaction order and the units of the rate constant for each of the two reactions.

The function makes the same fits that you made, but makes them all at once and plots the results side by side.

```
from fysisk_biokemi.utils.deter_reacti_order_activ import make_plot
rate_laws = {0: zeroth_order, 1: first_order, 2: second_order}
make_plot(df, rate_laws)
```



Information on 'make_plot'

The function 'make_plot' is beyond what is expected for this course, however if you are interested you can see the source code for it on [github](#).

Answer

Both are first order reactions for which the units of the rate constant is $\frac{1}{s}$.

(g) Activation energy

With the assumption that the Arrhenius constant A and the activation energy are temperature independent in the interval measured, use the Arrhenius equation to calculate the activation energy of the isomerization of the compound A.

You can use $R = 8.314 \times 10^{-3} \frac{\text{kJ}}{\text{mol} \cdot \text{K}}$

Caution

You need to derive the correct equation **before** using Python to calculate the result.

Perform the calculation in the cell below.

```
## Your task: Define known & fitted values
k1 = 20.08175
k2 = 29.96091
R = 8.318 * 10**(-3) # kJ / (mol K)
```



```
## Your task: Calculate the activation energy
## Remember: Derive the correct formula first using pen and paper!
activation_energy = R * np.log(k1/k2)/(1/T2 - 1/T1) # kJ/mol
print(activation_energy)
```

```
20.71400117402973
```

! Answer

The Arrhenius equation is

$$k = A \cdot \exp\left(\frac{-E_A}{RT}\right)$$

We have this for the two situations

$$k_1 = A \cdot \exp\left(\frac{-E_A}{RT_1}\right)$$

$$k_2 = A \cdot \exp\left(\frac{-E_A}{RT_2}\right)$$

Giving us two equations with two unknowns A and E_A where E_A is the activation energy that we are interested in.

Taking the log on both sides we get

$$\ln(k) = \ln(A) - \frac{E_A}{RT}$$

Where we can isolate for $\ln(A)$

$$\ln(A) = \ln(k) + \frac{E_A}{RT}$$

We can equate this for both our equations

$$\ln(k_1) + \frac{E_A}{RT_1} = \ln(k_2) + \frac{E_A}{RT_2}$$

Rearranging a bit we get

$$\ln(k_1) - \ln(k_2) = \frac{E_A}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

Isolating for E_A


$$\begin{aligned} E_A &= R \frac{\ln(k_1) - \ln(k_2)}{\left(\frac{1}{T_2} - \frac{1}{T_1} \right)} \\ &= \frac{R \ln\left(\frac{k_1}{k_2}\right)}{\left(\frac{1}{T_2} - \frac{1}{T_1} \right)} \end{aligned}$$

5 The pH effect on an active site histidine residue

Assume that a compound R can react with the unprotonated form of Histidine, His, to form a covalent reaction product, P:



The protonated form of Histidine, HisH⁺, is in equilibrium with His:



The pKa value for this acid-base equilibrium is 6.0. Further assume that the total concentration of Histidine, is

$$[\text{HisH}^+] + [\text{His}] = 10^{-3} \text{ M}$$

(a) Percentage unprotonated

What percentage of Histidine is unprotonated at pH 6.0. What do you notice about the relationship between pH and pKa in this case? (You should not need to do any arithmetic or algebra to answer this exercise.)

```
## Your task: What percentage is unprotonated?  
percentage_unprotonated = 50
```

! Answer

Because pH = pKa the percentage of unprotonated is 50%, so there is no need to do a direct calculation in this case.

(b) Concentration

Determine [His] at pH 6

```
## Your task: Assign known value and calculate [His]  
total_conc = 10**(-3)  
his_conc = total_conc * percentage_unprotonated / 100 # Unit: M  
print(his_conc)  
  
# Or directly  
his_conc = 0.5 * 10**(-3) # Unit: M  
print(his_conc)
```

```
0.0005  
0.0005
```

(c) Reaction order

The reaction equation for the reaction between His and R is

$$v = -\frac{d[\text{His}]}{dt} = k \cdot [\text{His}] \cdot [\text{R}]$$

What's the reaction order?

! Answer

Bimolecular so we expect second order reaction.

(d) High concentrations

If $[\text{R}]$ is much higher than $[\text{His}]$, what can then be concluded regarding the order of the reaction?

! Answer

$[\text{R}]$ will remain roughly constant - we are thus in the regime of a pseudo first order reaction

(e) New rate constant

Show how a new rate constant, k' , can be defined in these conditions. How does k' depend on R ?

! Answer

$$v = k \cdot [\text{His}] \cdot [\text{R}]$$

$$k' = k[\text{R}]$$

$$v = k'[\text{His}]$$

Then k' is linearly proportional with $[\text{R}]$.

(f) SI Units

At pH 6.0 the reaction rate $v = 1 \text{ mM} \cdot \text{s}^{-1}$.

Convert the reaction rate to SI-units given in $\text{M} \cdot \text{s}^{-1}$.

```
# Your task: Calculate v in SI units M / s
reaction_rate = 1 * 10**(-3) # Units: M / s
print(reaction_rate)
```

0.001

(g) Calculate the rate constant

Use the concentration of $[\text{His}]$ at pH 6 calculated in question (b), the reaction rate from (f) and constant $[\text{R}] = 0.2 \text{ M}$ to calculate the rate constant k .

```
## Your task: Set known value and calculate k ##  
## Hint: You can use the his_conc variable ##  
R_conc = 0.2  
k = reaction_rate / (R_conc * his_conc)  
print(k)
```

10.0

6 Qualitative description of Michaelis-Menten kinetics

```
import plotly
import fysisisk_biokemi.widgets as fbw
from fysisisk_biokemi.widgets.utils import enable_custom_widget_colab,
disable_custom_widget_colab
from IPython.display import display, HTML
enable_custom_widget_colab()
```

```
fbw.michaelis_menten_demo()
```

Use the widget created by the cell above to answer the following the questions.

(a) Change K_M

Setting $V_{\max} = 400$ and $[E]_{\text{tot}} = 0.0005$, change K_M from 0.5 to 5 to 10 to 50. Describe the change in curve appearance and explain it using the Michaelis-Menten (MM) equation.

(b) Meaning of K_M

What is the biochemical meaning of K_M ?

(c) Change $V_{\max}$ and $[E]_{\text{tot}}$

Now set $[E]_{\text{tot}} = 0.0005$, and $K_M = 5$. Change $V_{\max}$ step-wise from 40 to 900. Describe the changes you observe and how they relate to the MM equation.

After this, repeat the procedure for $[E]_{\text{tot}}$ increasing it 10-fold.

(d) Substrate concentrations way below K_M

Next, we will explore the regime where $[S] \ll K_M$. Initially, set $K_M = 10$ and zoom in on the regime from 0 to 1 on the X-axis. What does the MM equation look like in this regime? Use the MM equation and a simplifying assumption to show how this occurs.

Next, increase K_M five-fold and try to change $V_{\max}$ som the curve appearance is unchanged (beware of the autoscaling Y-axis).

Imagine an experiment conducted using only $[S]$ values from this range. How would this affect how well you could determine the MM parameters?

(f) Training estimation skills.

Train your estimation skills using the widget below.

```
fbw.michealis_menten_guess()
```

7 Design of an enzyme kinetics experiment

```
import matplotlib.pyplot as plt
from scipy.optimize import curve_fit
import numpy as np
import pandas as pd
from fysisk_biokemi.widgets import DataUploader
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

We have finally arrived at the part of the course where we are ready to analyse the data from your lab exercise in the first quarter. First, however we will analyse a similar data set where we are sure about the data quality (😊) to illustrate the process.

In the Excel document `design-enzyme-kineti-exper.xlsx` you will find a data set in which an enzyme catalyzed formation of product P, with varying start concentration of substrates, [S], was followed over time. The product absorbs light at a specific wavelength with an extinction coefficient of $0.068 \mu\text{M}^{-1} \cdot \text{cm}^{-1}$, and the absorbance was measured in a light path of 1 cm throughout the time course.

You can load the dataset using the cells below;

```
uploader = DataUploader()
uploader.display()
```

Run the next cell **after** uploading the file

```
df = uploader.get_dataframe()
display(df)
```

```
from IPython.display import display
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('design-enzyme-kineti-exper.xlsx') # Load from package for the solution
so it doesn't require to interact.
display(df)
```

	time_(s)	Abs_S1	Abs_S2	Abs_S4	Abs_S8	Abs_S16	Abs_S32	Abs_S64	Abs_S128	Abs_S256
0	0.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
1	1.0	0.002	0.004	0.008	0.015	0.025	0.038	0.051	0.063	0.068
2	2.0	0.004	0.008	0.015	0.028	0.048	0.073	0.099	0.121	0.131
...	...	...	...	...	...	...	...	...	...	...
18	18.0	0.022	0.042	0.080	0.145	0.246	0.376	0.511	0.623	0.677
19	19.0	0.022	0.043	0.082	0.149	0.252	0.385	0.524	0.639	0.693
20	20.0	0.023	0.044	0.084	0.152	0.257	0.394	0.535	0.653	0.709

The headers, like `Abs_S1` refer to the substrate concentration so S1 means a substrate concentration of $1 \mu\text{M}$.

(a) Convert extinction coefficient to SI.

Convert the extinction coefficient to units given in $M^{-1} \cdot cm^{-1}$ and assign it to a variable. Also assign the light path length to a variable.

```
## Your task: Assign known values in SI units.
ext_coeff = 0.068 / 10**(-6) # 1/(M cm)
l = 1 # cm
```

(b) Calculate concentrations

Using Lambert-Beers law, calculate the concentration of product, $[P]$, in M for each time series.

The cell below setups a loop calculate the concentrations for each of these current columns in the dataframe.

```
## Your task calculate the concentration for each 'Abs_SX' column and add it ##
## as a new column.
df['C_S1'] = df['Abs_S1'] / (ext_coeff * l)
df['C_S2'] = df['Abs_S2'] / (ext_coeff * l)
df['C_S4'] = df['Abs_S4'] / (ext_coeff * l)
df['C_S8'] = df['Abs_S8'] / (ext_coeff * l)
df['C_S16'] = df['Abs_S16'] / (ext_coeff * l)
df['C_S32'] = df['Abs_S32'] / (ext_coeff * l)
df['C_S64'] = df['Abs_S64'] / (ext_coeff * l)
df['C_S128'] = df['Abs_S128'] / (ext_coeff * l)
df['C_S256'] = df['Abs_S256'] / (ext_coeff * l)
```

! Alternative solution

A more advanced alternative solution is to use a for-loop like done below, however this is outside the scope of what you're going to learn in this course.

```
for s in [1, 2, 4, 8, 16, 32, 64, 128, 256]:
    df[f'C_{s}'] = df[f'Abs_{s}'] / (ext_coeff * l)
```

And we can check that the columns we expect have been added to the DataFrame.

```
display(df)
```

time_(s)	Abs_S1	Abs_S2	Abs_S4	Abs_S8	Abs_S16	Abs_S32	Abs_S64	Abs_S128	Abs_S256	C_S1	C_S2	C_S4	C_S8	C_S16	C_S32	C_S64	C_S128	C_S256
0	0.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
1	1.0	0.002	0.004	0.008	0.015	0.025	0.038	0.051	0.063	0.068	2.941	5.882	11.764	23.528	47.056	94.112	188.224	376.448
2	2.0	0.004	0.008	0.015	0.028	0.048	0.073	0.099	0.121	0.131	5.882	11.764	23.528	47.056	94.112	188.224	376.448	752.896
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
18	18.0	0.022	0.042	0.080	0.145	0.246	0.376	0.511	0.623	0.677	3.235	6.470	12.940	25.880	51.760	103.520	207.040	414.080

	time_(s)	Abs_S1	Abs_S2	Abs_S3	Abs_S4	Abs_S5	Abs_S6	Abs_S7	Abs_S8	Abs_S9	Abs_S10	Abs_S11	Abs_S12	Abs_S13	Abs_S14	Abs_S15	Abs_S16	Abs_S17	Abs_S18	Abs_S19	Abs_S20	Abs_S21	Abs_S22	Abs_S23	Abs_S24	Abs_S25	Abs_S26	Abs_S27	Abs_S28	Abs_S29	Abs_S30	Abs_S31	Abs_S32	Abs_S33	Abs_S34	Abs_S35	Abs_S36	Abs_S37	Abs_S38	Abs_S39	Abs_S40	Abs_S41	Abs_S42	Abs_S43	Abs_S44	Abs_S45	Abs_S46	Abs_S47	Abs_S48	Abs_S49	Abs_S50	Abs_S51	Abs_S52	Abs_S53	Abs_S54	Abs_S55	Abs_S56	Abs_S57	Abs_S58	Abs_S59	Abs_S60	Abs_S61	Abs_S62	Abs_S63	Abs_S64	Abs_S65	Abs_S66	Abs_S67	Abs_S68	Abs_S69	Abs_S70	Abs_S71	Abs_S72	Abs_S73	Abs_S74	Abs_S75	Abs_S76	Abs_S77	Abs_S78	Abs_S79	Abs_S80	Abs_S81	Abs_S82	Abs_S83	Abs_S84	Abs_S85	Abs_S86	Abs_S87	Abs_S88	Abs_S89	Abs_S90	Abs_S91	Abs_S92	Abs_S93	Abs_S94	Abs_S95	Abs_S96	Abs_S97	Abs_S98	Abs_S99	Abs_S100																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
19	19.0	0.022	0.043	0.082	0.149	0.252	0.385	0.524	0.639	0.693	3.235	3.323	3.392	3.458	3.521	3.581	3.640	3.698	3.755	3.811	3.866	3.920	3.973	4.025	4.076	4.126	4.175	4.223	4.270	4.316	4.361	4.405	4.448	4.490	4.531	4.571	4.610	4.648	4.685	4.721	4.757	4.792	4.826	4.860	4.893	4.925	4.957	4.988	5.019	5.049	5.079	5.108	5.137	5.165	5.193	5.221	5.248	5.275	5.302	5.328	5.354	5.380	5.405	5.430	5.455	5.480	5.504	5.528	5.552	5.576	5.600	5.623	5.646	5.669	5.692	5.715	5.737	5.760	5.782	5.804	5.826	5.848	5.869	5.891	5.912	5.933	5.954	5.975	5.995	6.016	6.036	6.056	6.076	6.096	6.116	6.136	6.155	6.175	6.194	6.214	6.233	6.252	6.271	6.290	6.309	6.328	6.347	6.365	6.384	6.402	6.421	6.439	6.457	6.475	6.493	6.511	6.529	6.547	6.564	6.582	6.600	6.617	6.635	6.652	6.669	6.686	6.703	6.720	6.737	6.754	6.771	6.788	6.805	6.822	6.839	6.855	6.872	6.889	6.905	6.922	6.938	6.955	6.971	6.988	7.004	7.021	7.037	7.053	7.069	7.085	7.101	7.117	7.133	7.149	7.164	7.180	7.196	7.211	7.227	7.242	7.258	7.273	7.288	7.304	7.319	7.334	7.349	7.364	7.379	7.394	7.409	7.424	7.439	7.454	7.469	7.484	7.499	7.514	7.529	7.544	7.559	7.573	7.588	7.603	7.618	7.633	7.647	7.662	7.677	7.691	7.706	7.721	7.735	7.750	7.764	7.779	7.793	7.808	7.822	7.837	7.851	7.866	7.880	7.894	7.909	7.923	7.937	7.951	7.966	7.980	7.994	8.009	8.023	8.037	8.051	8.066	8.080	8.094	8.108	8.122	8.136	8.150	8.164	8.178	8.192	8.206	8.220	8.234	8.248	8.262	8.276	8.290	8.304	8.318	8.332	8.346	8.359	8.373	8.387	8.401	8.415	8.429	8.443	8.456	8.470	8.484	8.498	8.512	8.525	8.539	8.553	8.567	8.580	8.594	8.608	8.621	8.635	8.648	8.662	8.675	8.689	8.702	8.716	8.729	8.742	8.756	8.769	8.782	8.796	8.809	8.822	8.836	8.849	8.862	8.876	8.889	8.902	8.915	8.928	8.941	8.954	8.967	8.980	8.993	9.006	9.019	9.032	9.045	9.058	9.071	9.084	9.097	9.110	9.123	9.136	9.149	9.162	9.175	9.188	9.201	9.214	9.227	9.240	9.253	9.266	9.279	9.292	9.305	9.318	9.331	9.344	9.357	9.370	9.383	9.396	9.409	9.422	9.435	9.448	9.461	9.474	9.487	9.500	9.513	9.526	9.539	9.552	9.565	9.578	9.591	9.604	9.617	9.630	9.643	9.656	9.669	9.682	9.695	9.708	9.721	9.734	9.747	9.760	9.773	9.786	9.799	9.812	9.825	9.838	9.851	9.864	9.877	9.890	9.903	9.916	9.929	9.942	9.955	9.968	9.981	9.994	10.007	10.020	10.033	10.046	10.059	10.072	10.085	10.098	10.111	10.124	10.137	10.150	10.163	10.176	10.189	10.202	10.215	10.228	10.241	10.254	10.267	10.280	10.293	10.306	10.319	10.332	10.345	10.358	10.371	10.384	10.397	10.410	10.423	10.436	10.449	10.462	10.475	10.488	10.501	10.514	10.527	10.540	10.553	10.566	10.579	10.592	10.605	10.618	10.631	10.644	10.657	10.670	10.683	10.696	10.709	10.722	10.735	10.748	10.761	10.774	10.787	10.800	10.813	10.826	10.839	10.852	10.865	10.878	10.891	10.904	10.917	10.930	10.943	10.956	10.969	10.982	10.995	11.008	11.021	11.034	11.047	11.060	11.073	11.086	11.099	11.112	11.125	11.138	11.151	11.164	11.177	11.190	11.203	11.216	11.229	11.242	11.255	11.268	11.281	11.294	11.307	11.320	11.333	11.346	11.359	11.372	11.385	11.398	11.411	11.424	11.437	11.450	11.463	11.476	11.489	11.502	11.515	11.528	11.541	11.554	11.567	11.580	11.593	11.606	11.619	11.632	11.645	11.658	11.671	11.684	11.697	11.710	11.723	11.736	11.749	11.762	11.775	11.788	11.801	11.814	11.827	11.840	11.853	11.866	11.879	11.892	11.905	11.918	11.931	11.944	11.957	11.970	11.983	11.996	12.009	12.022	12.035	12.048	12.061	12.074	12.087	12.100	12.113	12.126	12.139	12.152	12.165	12.178	12.191	12.204	12.217	12.230	12.243	12.256	12.269	12.282	12.295	12.308	12.321	12.334	12.347	12.360	12.373	12.386	12.399	12.412	12.425	12.438	12.451	12.464	12.477	12.490	12.503	12.516	12.529	12.542	12.555	12.568	12.581	12.594	12.607	12.620	12.633	12.646	12.659	12.672	12.685	12.698	12.711	12.724	12.737	12.750	12.763	12.776	12.789	12.802	12.815	12.828	12.841	12.854	12.867	12.880	12.893	12.906	12.919	12.932	12.945	12.958	12.971	12.984	12.997	13.010	13.023	13.036	13.049	13.062	13.075	13.088	13.101	13.114	13.127	13.140	13.153	13.166	13.179	13.192	13.205	13.218	13.231	13.244	13.257	13.270	13.283	13.296	13.309	13.322	13.335	13.348	13.361	13.374	13.387	13.400	13.413	13.426	13.439	13.452	13.465	13.478	13.491	13.504	13.517	13.530	13.543	13.556	13.569	13.582	13.595	13.608	13.621	13.634	13.647	13.660	13.673	13.686	13.699	13.712	13.725	13.738	13.751	13.764	13.777	13.790	13.803	13.816	13.829	13.842	13.855	13.868	13.881	13.894	13.907	13.920	13.933	13.946	13.959	13.972	13.985	13.998	14.011	14.024	14.037	14.050	14.063	14.076	14.089	14.102	14.115	14.128	14.141	14.154	14.167	14.180	14.193	14.206	14.219	14.232	14.245	14.258	14.271	14.284	14.297	14.310	14.323	14.336	14.349	14.362	14.375	14.388	14.401	14.414	14.427	14.440	14.453	14.466	14.479	14.492	14.505	14.518	14.531	14.544	14.557	14.570	14.583	14.596	14.609	14.622	14.635	14.648	14.661	14.674	14.687	14.700	14.713	14.726	14.739	14.752	14.765	14.778	14.791	14.804	14.817	14.830	14.843	14.856	14.869	14.882	14.895	14.908	14.921	14.934	14.947	14.960	14.973	14.986	14.999	15.012	15.025	15.038	15.051	15.064	15.077	15.090	15.103	15.116	15.129	15.142	15.155	15.168	15.181	15.194	15.207	15.220	15.233	15.246	15.259	15.272	15.285	15.298	15.311	15.324	15.337	15.350	15.363	15.376	15.389	15.402	15.415	15.428	15.441	15.454	15.467	15.480	15.493	15.506	15.519	15.532	15.545	15.558	15.571	15.584	15.597	15.610	15.623	15.636	15.649	15.662	15.675	15.688	15.701	15.714	15.727	15.740	15.753	15.766	15.779	15.792	15.805	15.818	15.831	15.844	15.857	15.870	15.883	15.896	15.909	15.922	15.935	15.948	15.961	15.974	15.987	15.999	16.012	16.025	16.038	16.051	16.064	16.077	16.090	16.103	16.116	16.129	16.142	16.155	16.168	16.181	16.194	16.207	16.220	16.233	16.246	16.259	16.272	16.285	16.298	16.311	16.324	16.337	16.350	16.363	16.376	16.389	16.402	16.415	16.428	16.441	16.454	16.467	16.480	16.493	16.506	16.519	16.532	16.545	16.558	16.571	16.584	16.597	16.610	16.623	16.636	16.649	16.662	16.675	16.688	16.701	16.714	16.727	16.740	16.753	16.766	16.779	16.792	16.805	16.818	16.831	16.844	16.857	16.870	16.883	16.896	16.909	16.922	16.935	16.948	16.961	16.974	16.987	16.999	17.012	17.025	17.038	17.051	17.064	17.077	17.090	17.103	17.116	17.129	17.142	17.155	17.168	17.181	17.194	17.207	17.220	17.233	17.246	17.259	17.272	17.285	17.298	17.311	17.324	17.337	17.350	17.363	17.376	17.389	17.402	17.415	17.428	17.441	17.454	17.467	17.480	17.493	17.506	17.519	17.532	17.545	17.558	17.571	17.584	17.597	17.610	17.623	17.636	17.649	17.662	17.675	17.688	17.701	17.714	17.727	17.740	17.753	17.766	17.779	17.792	17.805	17.818	17.831	17.844	17.857	17.870	17.883	17.896	17.909	17.922	17.935	17.948	17.

We want to use Python to determine V_0 for each concentration of S , in order to create a table of V_0 vs $[S]$. To do so we will fit linear functions to the initial parts of the curves, as the slope of these is then V_0 .

Start by defining a linear function;

```
def linear_func(x, a, b):  
    return a * x + b
```

As mentioned we want to get the slope in the initial part of the reaction, where the curves are approximately linear. To do so we need to fit not on all the data-points but only some of them.


We will first do this for the C_S32-column.

```
## Determine the x and y data  
x_data_full = df['time(s)']  
y_data_full = df['C_S32']  
  
## Your task: Set n_points to choose the number of points to fit with  
n_points = 5  
x_data = x_data_full[0:n_points] # This picks the first n_points points.  
y_data = y_data_full[0:n_points] # This picks the first n_points points.  
  
## Your task: Make the fit.  
fitted_parameters, trash = curve_fit(linear_func, x_data, y_data)  
a, b = fitted_parameters
```

The cell below plots your fit alongside the data

```
fig, ax = plt.subplots()  
t_smooth = np.linspace(0, 20)  
y_fit = linear_func(t_smooth, a, b)  
ax.plot(x_data_full, y_data_full, 'o')  
ax.plot(t_smooth, y_fit)  
ax.set_xlabel('Time (s)')  
ax.set_ylabel('Concentration')
```

```
Text(0, 0.5, 'Concentration')
```



You want the line to fit well with the initial approximately linear part of the data, adjust `n_points` such that this is the case.


(e) Using all the datasets

In the cell below `n_points` again determines the number of data-points used for the linear fit, and makes the fits and plots for every dataset.

```
from fysisk_biokemi.utils.design_enzyme_kineti_exper import make_fits_and_plots

## Your task: Select a number of points to fit with that is appropriate in
## all cases
n_points = 5

## This function does what you did in the previous exercise but for all the datasets. ##
slopes, concentrations = make_fits_and_plots(df, n_points)
```



[i](#) Information on 'make_fits_and_plot'

The function 'make_fits_and_plots' is beyond what is expected for this course, however if you are interested you can see the source code for it on [github](#).

Once you're satisfied with the fit you can display the collected slopes with the cell below

```
V0_df = pd.DataFrame({'[S]_(M)': concentrations, 'V0_(M/s)': slopes})
display(V0_df)
```

	[S]_(M)	V0_(M/s)
0	0.000001	2.941176e-08
1	0.000002	5.588235e-08
2	0.000004	1.058824e-07
...	...	...
6	0.000064	6.735294e-07
7	0.000128	8.220588e-07
8	0.000256	8.926471e-07

(f) Understanding of V_0

Why is it important to use V_0 rather than V at a later time point when creating the Michaelis-Menten plot?

! Answer

The initial velocities, V_0 , represent velocities in which the consumption/formation is only dependent on substrate concentration as no product is yet formed.

(g) Plotting again.


Plot V_0 against substrate concentration and estimate k_{cat} and K_M visually (remember units)

```
fig, ax = plt.subplots()

# Your task: Plot the V0_(M/s) versus [S]_(M)
ax.plot(V0_df['[S]_(M)'], V0_df['V0_(M/s)'], 'o')

## EXTRA: Axis labels.
ax.set_xlabel('[S] (M)')
ax.set_ylabel('$V_0$ (M/s)')
```

```
Text(0, 0.5, '$V_0$ (M/s)')
```



Use the figure to estimate the value of K_M and $V_{\max}$ and put those in the cell below.

```
# Your task: Put your estimated values here:
K_M_estimate = 30 * 10**(-6)
V_max_estimate = 10 * 10**(-7)
```

(h) Fitting with Michealis-Menten

Now we can fit with the Michealis-Menten equation, as always in a fitting task we need to implement the model.

```
def michaelis_menten(S, K_M, V_max):
    ## Your task: Implement the Michaelis-Menten model.
    result = (V_max * S) / (K_M + S)
    return result
```

Then use the next two cells to make the fit and plot it.

```
## Uses your estimate as initial guess
initial_guess = [K_M_estimate, V_max_estimate]
fitted_parameters, trash = curve_fit(michaelis_menten, V0_df['[S]_(M)'], V0_df['V0_(M/s)'],
initial_guess)

## Extract and prints the parameters
K_M_fit, V_max_fit = fitted_parameters
print(K_M_fit)
print(V_max_fit)
```

```
3.402856184164823e-05
1.0245295933534427e-06
```


How do these compare to your estimates of the two parameters?

```
fig, ax = plt.subplots()

## Evaluates the fit
S_smooth = np.linspace(0, 0.0003, 100)
V0_fit = michaelis_menten(S_smooth, K_M_fit, V_max_fit)

## Plots the fit and the data.
ax.plot(S_smooth, V0_fit)
ax.plot(V0_df['[S]_(M)'], V0_df['V0_(M/s)'], 'o')

## Axis labels.
ax.set_xlabel('[S] (M)')
ax.set_ylabel('$V_0$ (M/s)');
```



8 Lab Exercise Analysis: Michaelis Menten

```
import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
from scipy.optimize import curve_fit
from fysisk_biokemi.widgets import DataUploader
from IPython.display import display
```

In the “Introduction to the Molecules of Life” course, you performed an experiment where you measured absorption for different substrate concentrations as a function of time. The experiment was done with p-nitrophenylphosphate as the substrate, which turns into p-nitrophenol when the enzyme (alkaline phosphatase) acts on it.

If you do not have data, or you find that your data set is not of sufficient quality, you can use the following `kinetics_data.xlsx`.

Load your dataset using the widget below

```
uploader = DataUploader()
uploader.display()
```

Run the next cell **after** uploading the file

```
df = uploader.get_dataframe()
display(df)
```

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('kinetics_L0') # Load from package for the solution so it doesn't require
to interact.
display(df)
```

	time	0.8	0.4	0.2	0.12	0.05	0.02	0.01	0.003
0	0	0.032	0.101	0.018	0.032	0.015	0.013	0.008	-0.001
1	10	0.049	0.115	0.033	0.047	0.027	0.023	0.016	0.003
2	20	0.065	0.130	0.047	0.060	0.039	0.033	0.023	0.007
...	...	...	...	...	...	...	...	...	...
16	160	0.269	NaN	NaN	NaN	NaN	NaN	NaN	NaN
17	170	0.283	NaN	NaN	NaN	NaN	NaN	NaN	NaN
18	180	0.298	NaN	NaN	NaN	NaN	NaN	NaN	NaN

(a) Absorbance to concentration

The measured absorbances are unitless, they are the log of the ratio between the incoming light intensity and the transmitted light intensity. We would like to work with concentrations instead, so we need to convert the absorbance to concentration. The conversion is done using Lambert-Beer’s law.

The extinction coefficient of the product p-nitrophenol is $\varepsilon = 18000 \text{ M}^{-1}\text{cm}^{-1}$ at 405 nm and the path length of the cuvette is $l = 1 \text{ cm}$. We will construct a new DataFrame with concentrations

We will start by defining a function `lambert_beers` to calculate the concentrations

```
def lambert_beers(absorbance):
    ## Your task: Finish the implementation of LBs-law to calculate concentrations.
    L = 1 # cm
    ext_coeff = 18000 # 1 / (M cm)
    conc = absorbance / (L * ext_coeff) # M
    return conc
```

The next cell makes a new DataFrame by applying the `lambert_beers` function to every column in the original

```
data = {
    'time': df['time'],
    'S_0.8_mM': lambert_beers(df['0.8']),
    'S_0.4_mM': lambert_beers(df['0.4']),
    'S_0.2_mM': lambert_beers(df['0.2']),
    'S_0.12_mM': lambert_beers(df['0.12']),
    'S_0.05_mM': lambert_beers(df['0.05']),
    'S_0.02_mM': lambert_beers(df['0.02']),
    'S_0.01_mM': lambert_beers(df['0.01']),
    'S_0.003_mM': lambert_beers(df['0.003']),
}

df = pd.DataFrame(data)
display(df)
```

	time	S_0.8_mM	S_0.4_mM	S_0.2_mM	S_0.12_mM	S_0.05_mM	S_0.02_mM	S_0.01_mM	S_0.003_mM
0	0	0.000002	0.000006	0.000001	0.000002	8.333333e-07	2.222222e-07	4.444444e-07	5.555556e-08
1	10	0.000003	0.000006	0.000002	0.000003	1.500000e-06	1.277778e-06	6.888889e-07	1.666667e-07
2	20	0.000004	0.000007	0.000003	0.000003	2.166667e-06	1.833333e-06	1.277778e-06	3.888889e-07
...	...	...	...	...	...	...	...	...	...
16	160	0.000015	NaN	NaN	NaN	NaN	NaN	NaN	NaN
17	170	0.000016	NaN	NaN	NaN	NaN	NaN	NaN	NaN
18	180	0.000017	NaN	NaN	NaN	NaN	NaN	NaN	NaN

(b) Plot

Now let's plot the data. Plot each column separately as a function of time.

```
fig, ax = plt.subplots()


# Your task: Plot concentration vs. time for each column of the dataset,
# So 8 lines of code total.
```

```

ax.plot(df['time'], df['S_0.8_mM'], '-o', label='S_0.8_mM')
ax.plot(df['time'], df['S_0.4_mM'], '-o', label='S_0.4_mM')
ax.plot(df['time'], df['S_0.2_mM'], '-o', label='S_0.2_mM')
ax.plot(df['time'], df['S_0.12_mM'], '-o', label='S_0.12_mM')
ax.plot(df['time'], df['S_0.05_mM'], '-o', label='S_0.05_mM')
ax.plot(df['time'], df['S_0.02_mM'], '-o', label='S_0.02_mM')
ax.plot(df['time'], df['S_0.01_mM'], '-o', label='S_0.01_mM')
ax.plot(df['time'], df['S_0.003_mM'], '-o', label='S_0.003_mM')

ax.set_xlabel('Time [s]')
ax.set_ylabel('Concentration [M]')
ax.legend()
plt.show()

```



! Alternative solution: for-loop

An alternative solution is using a for-loop, this is beyond the scope of the course and you will not get exam questions regarding this.

However it is more flexible than writing code for every column - the same code could be used datasets with hundreds or thousands of columns.

```
fig, ax = plt.subplots()

for column_name in df.keys(): # This loops over each column name of the dataframe.
    if column_name != 'time': # Means "if column_name is not equal to 'time'".
        # This only happens when the above condition is met.
        ax.plot(df['time'], df[column_name], '-o', label=column_name)

ax.set_xlabel('Time [s]')
ax.set_ylabel('Concentration [M]')
ax.legend()
plt.show()
```

(c) Finding the slopes

In order to apply the Michaelis Menten equation we need the slopes of these curves as that is the reaction velocity V_0 , luckily we have learned how to make fits! So we make a fit to the linear function to determine the slope a in,

$$y(x) = ax + b$$

As always, start by writing the function

```
def linear_function(x, a, b):
    ## Your task: Implement the linear function
    result = ...
    return result
```

```
def linear_function(x, a, b):
    ## Your task: Implement the linear function
    result = a * x + b
    return result
```

Now we want to find the slope for each column, to help with that we will introduce a function that takes the time column and one of the concentration columns and returns the slope

```
def find_slope(time_data, abs_data):

    ## This selects the times and absorbances that have been measured (not NaN)
    ## Don't worry about this - this is just to help you with this exercise.
    indices = np.where(~abs_data.isna())[0]
    time_data = time_data[indices]
```

```

abs_data = abs_data[indices]

## Find the slope and the intercept using curve_fit and return the slope
fitted_parameters, trash = curve_fit(linear_function, time_data, abs_data)
slope = fitted_parameters[0]
return slope

```

Now we want to use this for each set of measurements at different substrate concentrations.

```

## Your task: Find the slopes for every substrate concentration.
slope_08 = find_slope(df['time'], df['S_0.8_mM'])
slope_04 = find_slope(df['time'], df['S_0.4_mM'])
slope_02 = find_slope(df['time'], df['S_0.2_mM'])
slope_012 = find_slope(df['time'], df['S_0.12_mM'])
slope_005 = find_slope(df['time'], df['S_0.05_mM'])
slope_002 = find_slope(df['time'], df['S_0.02_mM'])
slope_001 = find_slope(df['time'], df['S_0.01_mM'])
slope_0003 = find_slope(df['time'], df['S_0.003_mM'])

## This gathers the slopes and makes an array with the substrate concentrations.
substrate_concentrations = np.array([0.8, 0.4, 0.2, 0.12, 0.05, 0.02, 0.01, 0.003]) *
10**(-3)
slopes = np.array([slope_08, slope_04, slope_02, slope_012, slope_005, slope_002,
slope_001, slope_0003])

```

Now make a plot of the slopes versus the substrate concentrations using the two arrays substrate_concentrations and slopes.


```

fig, ax = plt.subplots()

## Your task plot the slopes vs. substrate concentration.
ax.plot(substrate_concentrations, slopes, 'o')

## Sets x and y axis labels.
ax.set_xlabel('Substrate concentration')
ax.set_ylabel('Slope')
plt.show()

```



```
## Your task: Estimate K_M and V_max from the plot.
K_M_estimate = 0.00001
V_max_estimate = 8 * 10**(-8)
```

(d) Michaelis Menten

Now that we have the slopes we can fit to the Michaelis Menten equation, again we need to implement the equation

```
def michaelis_menten(S, K_M, V_max):
    ## Your task: Implement the Michaelis-Menten model.
    result = (V_max * S) / (K_M + S)
    return result
```

Then fit to the slopes we extracted to the Michaelis-Menten equation

```
## This is your estimates as the initial guess.
initial_guess = [K_M_estimate, V_max_estimate]

## Your task: Make a fit to the Michaelis-Menten equation & extract K_M and V_max.
fitted_parameters, trash = curve_fit(michaelis_menten, substrate_concentrations, slopes,
initial_guess)
K_M_fit, V_max_fit = fitted_parameters

print(K_M_fit)
print(V_max_fit)
```

```
1.1035718037533613e-05
8.121800807252114e-08
```

- What is the physical meaning of K_M and how does it relate to binding affinity?
- What's the meaning of V_{max} ? Why does the curve saturate?


(e) Plot with fit

```
S_smooth = np.linspace(substrate_concentrations.min(), substrate_concentrations.max())
V0_fit = michaelis_menten(S_smooth, K_M_fit, V_max_fit)
```

```
fig, ax = plt.subplots()

## Your task: Plot the data and the fit
ax.plot(substrate_concentrations, slopes, 'o', label='Data')
ax.plot(S_smooth, V0_fit, label='Fit')

## These add x and y axis labels.
ax.set_xlabel('Substrate concentration (M)')
ax.set_ylabel('$V_0$ (M/s)')
plt.show()
```



(f) Turnover number

Given that the enzyme concentration was $3.33 \frac{\text{mg}}{\text{L}}$ and the molecular weight of the enzyme (alkaline phosphatase) is $86000 \frac{\text{g}}{\text{mol}}$, calculate the turnover number k_{cat} .

Start by converting the enzyme concentration to molar units and then calculate the turnover number.

```
## Your task: Calculate the enzyme concentration and use that and your fit to  
## calculate kcat.  
enzyme_conc = (3.33*10**(-3)) / 86000  
kcat = V_max_fit / enzyme_conc  
print(kcat)
```

```
2.097522130401447
```

- What is the biochemical meaning of this number?