

Week 46

Table of contents

1 Estimation of binding affinity	1
2 Dialysis experiment	2
3 Interpretation of binding data.	6
4 Determination of type and strength of cooperativity	10
5 Competition in ligand binding	15
6 Difference between Q and K_{eq}	18
7 Determination of ΔH° for a reaction from the van't Hoff plot.	22
8 ATP hydrolysis	25

```
try:
    import fysisk_biokemi
    print("Already installed")
except ImportError:
    %pip install -q "fysisk_biokemi[pyodide] @ https://au-mbg.github.io/fysisk-biokemi/
    wheels/fysisk_biokemi-0.2.2-py3-none-any.whl"
```

1 Estimation of binding affinity

(a) Train estimation skills

Train your estimation skills using the widget below.

```
from fysisk_biokemi.widgets.utils.colab import enable_custom_widget_colab
from fysisk_biokemi.widgets import estimate_kd

enable_custom_widget_colab()
estimate_kd()
```

(b) Compare to quadratic

The widget below shows the curves for θ using both the simple expression assuming that $L = L_{tot}$ and the quadratic binding expression. Vary K_D and P_{total} to work out a rule of thumb for when the two equations give a similar curve.

```
from fysisk_biokemi.widgets.utils.colab import enable_custom_widget_colab
from fysisk_biokemi.widgets import visualize_simple_vs_quadratic

enable_custom_widget_colab()
visualize_simple_vs_quadratic()
```

```
answer = "They are equal when  $P_{tot} \ll K_D$ "
```

2 Dialysis experiment

```
import numpy as np
import matplotlib.pyplot as plt
import pandas as pd
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

A dialysis experiment was set up where equal amounts of a protein were separately dialyzing against buffers containing different concentrations of a ligand – each measurement was done in triplicate. The average number of ligands bound per protein molecule, $\bar{n}$ were obtained from these experiments. The corresponding concentrations of free ligand and values are given in dataset `dialys-exper.xlsx`.

(a) Load the dataset

```
from IPython.display import display
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('dialysis_experiment') # Load from package for the solution so it doesn't
require to interact.
display(df)
```

	Free_ligand_(uM)	n_bar
0	0.01	0.0059
1	0.01	0.0059
2	0.01	0.0060
...	...	...
39	80.00	1.8769
40	80.00	1.9631
41	80.00	1.8930

(b) Explain calculation of $\bar{n}$

Explain how the values of $\bar{n}$ is calculated when knowing the concentrations of ligand inside and outside the dialysis bag, as well as the total concentration of the protein, $[P_{\text{tot}}]$.

i Calculation of $\bar{n}$

The concentration of free ligand $[L]$ is the same inside and outside the dialysis bag at equilibrium. Outside there is no protein, so only free ligand $L_{outside} = [L]$

The total concentration of ligand inside the dialysis bag is thus the combination of free and bound ligand $L_{inside} = [L] + [PL]$. $[PL]$ can thus be calculated as:

$$[PL] = [L_{inside}] - [L_{outside}]$$

This can be combined with $[P_{tot}]$ to give $\bar{n}$:

$$\bar{n} = \frac{[L_{inside}] - [L_{outside}]}{[P_{tot}]}$$

(c) Molar concentrations

Convert the concentrations of free ligand to SI-units given in M, add it as a row to the DataFrame.

```
df['Free_ligand_(M)'] = df['Free_ligand_(uM)'] * 10**(-6)
display(df)
```


	Free_ligand_(uM)	n_bar	Free_ligand_(M)
0	0.01	0.0059	1.000000e-08
1	0.01	0.0059	1.000000e-08
2	0.01	0.0060	1.000000e-08
...	...	...	...
39	80.00	1.8769	8.000000e-05
40	80.00	1.9631	8.000000e-05
41	80.00	1.8930	8.000000e-05

(d) Plot the data

```
fig, ax = plt.subplots()

## Your task: Plot the raw data
## With the ligand concentration df['Free_ligand_(M)'] on the x-axis
## And with n-bar df['n_bar'] on the y-axis.
ax.plot(df['Free_ligand_(M)'], df['n_bar'], 'o')

## EXTRAS: You do not need to understand this ##
## This adds customization ##
ax.set_xlabel('Free ligand [L] (M)', fontsize=16)
ax.set_ylabel(r'$\bar{n}$', fontsize=16);
```



(e) Prepare for fitting

Now we want to fit the data to extract K_D and n , by using the equation

$$\bar{n}([L_{\text{free}}]) = N \frac{[L_{\text{free}}]}{K_D + [L_{\text{free}}]}$$

To do so we need to implement it as a Python function

```
def nbar(L, N, K_D):
    return N * L / (K_D + L)

## This calculates the function and prints for two different sets of values ##
print(nbar(1, 1, 1)) # Should give 1/2
print(nbar(21, 47, 2.5)) # Should give 42
```

```
0.5
42.0
```

(f) Actually fitting

Finish the code to perform the fitting in the cell below.

```
from scipy.optimize import curve_fit

# Choose the variables from the dataframe
x = df['Free_ligand_(M)']
y = df['n_bar']

## Your task: Make initial guesses for the two parameters ##
```

```

K_D_guess = 10**(-5)
N_guess = 1
initial_guess = [K_D_guess, N_guess]

## Your task: Give the four arguments in the correct order to make calculate ##
## the fitted parameters ##
fitted_parameters, pcov = curve_fit(nbar, x, y, initial_guess)

# Print the parameters
N_fit, K_D_fit = fitted_parameters
print(fitted_parameters)
print(N_fit)
print(K_D_fit)

```

```

[2.00882847e+00 3.39599709e-06]
2.008828473002213
3.3959970874778938e-06

```

Are the parameters you find reasonable? How can you tell if they are reasonable by looking at the plot you made earlier?

(g) Plot with fit

When we have the fitted parameters we can calculate and plot the function. To do so we make an array of values for the independent variable and use our function to calculate the dependent variable

```

## This makes 50 equally spaced points ##
L_smooth = np.linspace(0, x.max()*1.2, 50)

## Your task: Calculate the fitted function using the found parameters and L_smooth
nbar_calc = nbar(L_smooth, N_fit, K_D_fit)

```

Now that we calculated the dependent variable we can plot the fit along with the data.

```


fig, ax = plt.subplots()

## Your task: Plot the fitted function ##
ax.plot(L_smooth, nbar_calc)

## Plots the data ##
ax.plot(df['Free_ligand_(M)'], df['n_bar'], 'o')

## EXTRA: You do not need to understand this ##
## This customizes the plot ##
## It also adds lines indicating the values of n and K_D found ##
ax.axhline(N_fit, color='red', label=r'$N$')
ax.axvline(K_D_fit, color='green', label=r'$K_D$')
ax.set_xlabel('Free ligand [L] (M)', fontsize=16)
ax.set_ylabel(r'$\bar{n}$', fontsize=16)
ax.legend(fontsize=16)

```



3 Interpretation of binding data.

```
import numpy as np
import pandas as pd
from IPython.display import display
import matplotlib.pyplot as plt
from scipy.optimize import curve_fit
pd.set_option('display.max_rows', 6)
```

The `inter-bindin-data.xlsx` contains a protein binding dataset.

(a) Load the dataset

Load the dataset with pandas. Set `data_file_path` to the path of the dataset file named above.

```
from IPython.display import display
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('interpret_week48') # Load from package for the solution so it doesn't
require to interact.
display(df)
```

	[L]_(uM)	nbar
0	0.01	-0.001897
1	0.01	0.001821
2	0.01	0.003624
...	...	...

	[L]_(uM)	nbar
42	160.00	0.939608
43	160.00	0.937402
44	160.00	0.921801

(b) SI Units

Add a new column to the DataFrame with the ligand concentration in SI units.

```
## Your task: Add a column with the concentration in M ##
## Call this column df['[L]_(M)'] ##
## Recall that you can say df['[L]_(M)'] = df["[L]_(uM)"] * XXX ##
## Where XXX is the factor to multiply by. ##
df['[L]_(M)'] = df["[L]_(uM)"] * 10**(-6)
display(df)
```

	[L]_(uM)	nbar	[L]_(M)
0	0.01	-0.001897	1.000000e-08
1	0.01	0.001821	1.000000e-08
2	0.01	0.003624	1.000000e-08
...	...	...	...
42	160.00	0.939608	1.600000e-04
43	160.00	0.937402	1.600000e-04
44	160.00	0.921801	1.600000e-04

(c) Plot the data

Make plots of the binding data directly with a linear and logarithmic x-axis.

Estimate K_D by visual inspection of these plots!

```
# This makes a figure with two axes.
fig, axes = plt.subplots(1, 2)

## We index with [0] to plot in the first axis. ##
## Don't worry about remembering this ##
ax = axes[0]
ax.plot(df['[L]_(M)'], df['nbar'], 'o')

## We index with [1] to plot in the first axis. ##
## Don't worry about remembering this ##
## We will make this a log plot ##
ax = axes[1]

## Your task: Plot the data [L]_(M) vs nbar.
```

```


## Same exact code as above
ax = axes[1]
ax.plot(df['[L](M)'], df['nbar'], 'o')
ax.set_xscale('log')

## EXTRA: You do not need to understand this ##
## Customization: Adds labels for the axis in both plots.
ax = axes[0]
ax.set_xlabel('[L](M)', fontsize=14)
ax.set_ylabel(r'$\bar{n}$', fontsize=14)

ax = axes[1]
ax.set_xlabel('log [L](M)', fontsize=14)
ax.set_ylabel(r'$\bar{n}$', fontsize=14)

```

```
Text(0, 0.5, '$\bar{n}$')
```



i Note

This command `ax.set_xscale('log')` tells matplotlib that we want the x-axis to use a log-scale.

```
K_D_estimate = 5 * 10**(-5)
```

(d) Make a fit

Make a fit to determine K_D , as always we start by implementing the function to fit with


```
def n_bar(L, K_D):
    ## Your task: Implement the function to fit with
    ## Use your biochemical knowledge to decide which function that is.
    return L / (L + K_D)
```

And then we can make the fit

```
## Your task: Choose the variables from the dataframe ##
x = df['[L](M)']
y = df['nbar']

# Initial guess
initial_guess = [K_D_estimate]

## Your task: Finish the curve fitting
fitted_parameters, trash = curve_fit(n_bar, x, y, initial_guess)

# Print the parameters
K_D_fit = fitted_parameters[0]
print(K_D_fit)
```

```
1.2457934647913883e-05
```

(e) Compare

Compare the fitted values with your guess.

```
## These lines make equally spaced points on a normal linear axis ##
## And on a logarithmic axis. ##
L_smooth = np.linspace(0, df['[L](M)'].max(), 100)
L_smooth_log = np.geomspace(df['[L](M)'].min(), df['[L](M)'].max(), 100)

## Your task: Evaluate the fitted function with L_smooth and L_smooth_log ##
nbar_fit = n_bar(L_smooth, K_D_fit)
nbar_fit_log = n_bar(L_smooth_log, K_D_fit)
```

Now we can make a plot.

```
# This makes a figure with two axes.
fig, axes = plt.subplots(1, 2, figsize=(9, 4))

# Index with [0] to plot in the first axis - Linear plot
ax = axes[0]
ax.plot(df['[L](M)'], df['nbar'], 'o', color='C0')
ax.plot(L_smooth, nbar_fit, color='C2')

ax.axvline(K_D_estimate, label='Estimate', color='C1')
ax.axvline(K_D_fit, label='Fit', color='C2')
ax.set_xlabel('[L](M)', fontsize=14)
```


```

ax.set_ylabel(r'\bar{n}', fontsize=14)
ax.legend()

# Index with [1] to plot in the second axis - Log plot.
ax = axes[1]
ax.plot(df['[L](M)'], df['nbar'], 'o', color='C0')
ax.plot(L_smooth_log, nbar_fit_log, color='C2')

ax.set_xlabel('[L](M)', fontsize=14)
ax.set_ylabel(r'\bar{n}', fontsize=14)
ax.axvline(K_D_estimate, label='Estimate', color='C1')
ax.axvline(K_D_fit, label='Fit', color='C2')
ax.legend()
ax.set_xscale('log')

```



(f) Saturation

Based on the value of K_D found from the fit,

- At which concentration do you expect 10% saturation?
- At which concentration do you expect 90% saturation?

```

"""
At K_D/9 and 9*K_D.
"""

```

4 Determination of type and strength of cooperativity

```
import pandas as pd
import matplotlib.pyplot as plt
import numpy as np
from IPython.display import display
from scipy.optimize import curve_fit
```

The binding of NAD⁺ to the protein yeast glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was studied using equilibrium dialysis. The enzyme concentration was 71 μ M. The concentration of $[\text{NAD}_{\text{free}}^+]$ and the corresponding values of $\bar{n}$ were determined with the resulting data found in the dataset `deter-type-streng-coope.xlsx`.

The dataset consists of three independent repetitions of the same experiment.

(a) Load the dataset

Load the dataset with pandas. Set `data_file_path` to the path of the dataset file named above.

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('determination_coop_week48') # Load from package for the solution so it
doesn't require to interact.
display(df)
```

	[NAD+free]_(uM)	nbar1	nbar2	nbar3
0	10	0.250	0.261	0.243
1	13	0.394	0.389	0.379
2	21	0.802	0.801	0.797
...	...	...	...	...
6	68	3.056	3.020	3.091
7	125	3.507	3.543	3.522
8	211	3.704	3.741	3.699

(b) Averaging and units.

Start by adding a new column to the DataFrame with the average value of $\bar{n}$ across the three series



Tip

Remember that you can set a new column based on a computation using one or more other columns, e.g.

```
df['new_col'] = df['col1'] + df['col2']
```

```
## Your task: Calculate the of the values in the three columns ##
## nbar1, nbar2, nbar3 ##
df['nbar_avg'] = (df['nbar1'] + df['nbar2'] + df['nbar3']) / 3
```

Now also add a column with the ligand concentration in SI units with the column-name [NAD+free]_(M).

```
## Task: Calculate the ligand concentration in M and ##
## set it as a new column with the name '[NAD+free]_(M)'
df['[NAD+free]_(M)'] = df['[NAD+free]_(uM)'] * 10**(-6)
display(df)
```

	[NAD+free]_(uM)	nbar1	nbar2	nbar3	nbar_avg	[NAD+free]_(M)
0	10	0.250	0.261	0.243	0.251333	0.000010
1	13	0.394	0.389	0.379	0.387333	0.000013
2	21	0.802	0.801	0.797	0.800000	0.000021
...	...	...	...	...	...	...
6	68	3.056	3.020	3.091	3.055667	0.000068
7	125	3.507	3.543	3.522	3.524000	0.000125
8	211	3.704	3.741	3.699	3.714667	0.000211

Finally, set the concentration of the GADPH in SI units

```
## Task: Convert the GADPH concentration to M ##
## Set it to the variable c_gadph ##
## Add a comment about the unit ##
c_gadph = 71 * 10**(-6) # M
```


(c) Plot

Make a plot of the average $\bar{n}$ as a function of $[\text{NAD}_{\text{free}}^+]$.

```
fig, ax = plt.subplots()

## Your task: Add plot of nbar vs [NAD+_free]_(M)
ax.plot(df['[NAD+free]_(M)'], df['nbar_avg'], 'o')

## EXTRA: You don't need to understand this
## Customization
ax.set_xlabel(r'$[\text{NAD}^+_{\text{free}}]$', fontsize=14)
ax.set_ylabel(r'$\bar{n}$', fontsize=14)
plt.show()
```



(d) Scatchard plot


Make a Scatchard plot based on the average $\bar{n}$.

```
# Your task: Calculate nbar / L
nbar_over_L = df['nbar_avg'] / df['[NAD+free]_(M)']

fig, ax = plt.subplots()

# Your task: Choose the right things to make a Scatchard plot.
ax.plot(df['nbar_avg'], nbar_over_L, 'o')

## EXTRA: You dont need to understand this.
## Adds customization.
ax.set_xlabel(r'$\bar{n}$', fontsize=14)
ax.set_ylabel(r'$\bar{n}/L$', fontsize=14)
ax.set_xlim([0, df['nbar_avg'].max()*1.2])
ax.set_ylim([0, nbar_over_L.max() * 1.2])
```



(e) Binding sites

Estimate by eye how many binding sites GAPDH contains for NAD^+ ?

```
"""
Saturation suggest asymptotic approximation to 4, which seems like the like
intercept of the non-linear scatchard plot.
"""
```

(f) Cooperativity

Based on a visual inspection of the plot above, are there in signs of cooperativity? If so, which kind?

```
"""
Positive cooperativity as witnessed e.g. by the increasing slope in the early
part of the lienar plot
"""
```

(g) Fit

Make a fit using the functional form

$$\bar{n} = N \frac{[L]^h}{K + [L]^h}$$

As usual, start by defining the function in Python

```
def n_bar(L, N, K, h):
    ## Your task: Implement the function
    # Be careful with parentheses!
```

```

result = N * L**h / (K + L**h)
return result

```

Now we can fit

```

## We want to use all the data for the fit, but it was given in three different columns.
## We stitch it together to make a column containing all x-coordinates
## And another column containing all y-coordinates
x = np.concatenate([df['[NAD+free]_(M)'], df['[NAD+free]_(M)'], df['[NAD+free]_(M)'])
y = np.concatenate([df['nbar1'], df['nbar2'], df['nbar3']])

## Your task: Make initial guesses for the three parameters
## The order is that of the function
## So N, K and h.
initial_guess = [1, 1, 1]

## Your task: Make the fit
## Note: The bounds argument has been added to have the function
## only explore positive sets of parameters.
fitted_parameters, trash = curve_fit(n_bar, x, y, initial_guess, bounds=(0, np.inf))

# Print the parameters
N_fit, K_fit, h_fit = fitted_parameters
print('N_fit', N_fit)
print('K_fit', K_fit)
print('h_fit', h_fit)

```

```

N_fit 3.997873553931702
K_fit 5.9798561894818436e-09
h_fit 1.8627077063542798

```

Do the fitted parameters support your intuitive reading of the type of cooperativity?

i Note

In the data sets used, repeated experiments were given as a three separate columns, which is a quite natural way of recording data during an experiment. However, it is not an appropriate format for regression, where we want all the data-points in a single column. In this exercise we did a bit of *data wrangling* to make the data appropriate for what we want to do with it - in this case by creating a new joint column containing all the data by using the `np.concatenate` function.

5 Competition in ligand binding

Below is given the general expression for saturation of a binding site by one of the ligands, L , when two ligands L and C are competing for binding to the same site on a protein. Assume that $[P_T] = 10^{-9}$ M.

$$\theta_L = \frac{[PL]}{[P_T]} = \frac{1}{\frac{K_D}{[L]} \left(1 + \frac{[C]}{K_C}\right) + 1}$$

Consider these four situations

#	$[L_T]$	$[C_T]$	K_D	K_C
1	$1 \cdot 10^{-3}$ M	0	$1 \cdot 10^{-5}$ M	$1 \cdot 10^{-6}$ M
2	$1 \cdot 10^{-3}$ M	$1 \cdot 10^{-2}$ M	$1 \cdot 10^{-5}$ M	$1 \cdot 10^{-6}$ M
3	$1 \cdot 10^{-3}$ M	$1 \cdot 10^{-3}$ M	$1 \cdot 10^{-5}$ M	$1 \cdot 10^{-5}$ M
4	$1 \cdot 10^{-5}$ M	$1 \cdot 10^{-6}$ M	$1 \cdot 10^{-5}$ M	$1 \cdot 10^{-6}$ M

(a) Explain simplification

Explain how the fact that $[P_T]$ is much smaller than $[L_T]$ and $[C_T]$ simplifies the calculations using the above equation.

```
"""
When P(total) is << L(total) and C(total), we can assume that L(total) =[L] and C(total) =
[C].
"""
```

(b) Ligand degree of saturation

Calculate the degree of saturation of the protein with ligand L in the four situations.

Start by writing a Python function that calculates the degree of saturation θ .

```
def bound_fraction(L, C, Kd, Kc):
    """ Your task: Write the equation for binding saturation.
    """ Be careful with parentheses!
    theta = 1 / ((Kd / L)*(1+(C/Kc))+1)
    return theta
```

Then use that function to calculate θ_L for each situation.

```
""" Your task: Calculate the degree of saturation for each set of parameters """
theta_L_1 = bound_fraction(10**(-3), 0, 10**(-5), 10**(-18))
theta_L_2 = bound_fraction(10**(-3), 10**(-2), 10**(-5), 10**(-6))
theta_L_3 = bound_fraction(10**(-3), 10**(-3), 10**(-5), 10**(-5))
theta_L_4 = bound_fraction(10**(-5), 10**(-6), 10**(-5), 10**(-6))

""" Prints the results so we can analyze them.
print('Situation 1', theta_L_1)
print('Situation 2', theta_L_2)
print('Situation 3', theta_L_3)
print('Situation 4', theta_L_4)
```



```

Situation 1 0.9900990099009901
Situation 2 0.0099000099000099
Situation 3 0.49751243781094534
Situation 4 0.3333333333333333

```

(c) Competitor degree of saturation

What is the degree of saturation with respect to the competitor C in #1, #2, #3 and #4?

```

## Your task: Calculate the degree of saturation for each set of parameters ##
## Note: Now you are calculating it for the competitor!
theta_C_1 = 0 # [C] = 0
theta_C_2 = bound_fraction(10**(-2), 10**(-3), 10**(-6), 10**(-5))
theta_C_3 = bound_fraction(10**(-3), 10**(-3), 10**(-5), 10**(-5))
theta_C_4 = bound_fraction(10**(-6), 10**(-5), 10**(-6), 10**(-5))

## Prints the results so we can analyze them.
print('Situation 1', theta_C_1)
print('Situation 2', theta_C_2)
print('Situation 3', theta_C_3)
print('Situation 4', theta_C_4)

```

```

Situation 1 0
Situation 2 0.99000099000099
Situation 3 0.49751243781094534
Situation 4 0.3333333333333333

```

(d) Fraction of $[P_{\text{free}}]$

What is the fraction of $[P_{\text{free}}]$ in #1, #2, #3 and #4?



Tip

Consider how to express the fraction of $[P_{\text{free}}]$ in terms of θ_{L_X} and θ_{C_X} .

```

theta_free_1 = 1 - (theta_L_1 + theta_C_1)
theta_free_2 = 1 - (theta_L_2 + theta_C_2)
theta_free_3 = 1 - (theta_L_3 + theta_C_3)
theta_free_4 = 1 - (theta_L_4 + theta_C_4)

## Prints the results so we can analyze them.
print('Situation 1', theta_free_1)
print('Situation 2', theta_free_2)
print('Situation 3', theta_free_3)
print('Situation 4', theta_free_4)

```

```

Situation 1 0.00990099009900991
Situation 2 9.900009900010165e-05

```

Situation 3 0.004975124378109319
Situation 4 0.3333333333333337

(e) Qualitative analysis

Describe in your own words why these four competitive situations end up with their respective binding patterns. Consider the relative concentrations and affinities of ligands L and C.

```
"""
1. The competitor concentration is zero, so only the ligand L binds. Affinity/concentration
is such that nearly all proteins have an L-ligand bound.

2. The affinity and concentration of C is higher so it binds more and the ratio of affinity
and concentration is high enough that nearly all proteins have a C-ligand bound.


3. The affinity and concentration of L and C are equal and they both bind equally. Again,
the affinities/concentrations are such that nearly all proteins have a ligand bound.

4.  $K_D > K_C$  with the same factor as  $[L_T] > [C_T]$  so they bind equally but
there are still free proteins as the concentrations are low relative to the affinities.
"""
```

6 Difference between Q and K_{eq}

```
import matplotlib.pyplot as plt
import pandas as pd
from IPython.display import display
pd.set_option('display.max_rows', 6)
```

Consider the reaction:



The time course of the reaction was followed monitoring formation of the product B . The starting concentration of A was 50 μM , and as you will learn later in the course, the reversible reaction going from A to B follows first order kinetics.

(a) Load & convert

The data is contained in the `diff_q_keq_data.xlsx` file.

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('diff_q_keq') # Load from package for the solution so it doesn't require
to interact.
display(df)
```

	Time_(min)	[B]_(uM)
0	0	0.000000

	Time_(min)	[B]_(uM)
1	1	12.756091
2	2	21.896230
...	...	...
33	33	44.999248
34	34	44.999461
35	35	44.999614

Convert the measured concentrations of B , $[B]$, to concentrations in the SI-unit M (mol/L). Repeat for the starting concentration of A , $[A_0]$.

```
## Your task: Convert the concentration of B to M and set in new column ##
df['[B]_(M)'] = df['[B]_(uM)'] * 10**(-6) # Unit: M

## Your task: Set the initial concentration of A, in the variable A0.
## Remember to convert to M.
A0 = 50 * 10**(-6) # Unit: M
```

(b) Law of mass balance

Use the laws of mass balance to calculate $[A]$ as a function of time in the unit M, add it as a column to the DataFrame

```
df['[A]_(M)'] = A0 - df['[B]_(M)']
```


Make a plot of the time vs. the concentration of A and of time versus the concentration of B .

```
fig, ax = plt.subplots()
## Your task: Plot time vs. concentrations.
ax.plot(df['Time_(min)'], df['[A]_(M)'], label='[A]') # Time vs. concentration of A
ax.plot(df['Time_(min)'], df['[B]_(M)'], label='[B]') # Time vs. concentration of B

## This shows the labels.
ax.legend()

## EXTRA: You do not need to understand this:
ax.set_xlabel('Time [min]')
ax.set_ylabel('Concentrations [M]')
```

```
Text(0, 0.5, 'Concentrations [M]')
```



(c) Mass action ratio

Calculate and plot the mass action ratio Q as a function of time.

```
## Your task: Calculate the mass action ratio
Q = df['[B]_(M)'] / df['[A]_(M)']
```


Now plot the mass action ratio versus time.

```
fig, ax = plt.subplots()

## Your task: Make a plot of the axis from the dataframe df['Time_(min)'] vs Q
ax.plot(df['Time_(min)'], Q)

## EXTRA: You do not need to understand this
## Adds customization.
ax.set_xlabel('Time (min)')
ax.set_ylabel('Mass action ratio $Q$')
```

```
Text(0, 0.5, 'Mass action ratio $Q$')
```



(d) Q vs K_{eq}

What is the difference between Q and K_{eq} ?

```
"""
```

Q is the mass action ratio at any time throughout the reaction. K_{eq} is the mass action ratio at equilibrium only, and tells about the energetic favorisation of one direction over the other.

```
"""
```

(e) Favoured compound

Which of the two compounds A and B is favored at equilibrium? (Favored: highest concentration)

```
"""
```

When the reaction reaches close to equilibrium, more B is present as compared to A , telling that B is favored.

$K_{eq} > 1 \rightarrow B$ is favored

```
"""
```

(f) Equilibrium constant

What is the equilibrium constant of this reaction?

```
"""
```

Can also just read the concentrations values at eq from the graph and calculate K_{eq} from those.

But it can of course also just be calculated.

```
"""
```

```
K_eq = df['[B](M)'][35] / df['[A](M)'][35]
print(K_eq)
```

8.999228315035122

7 Determination of ΔH° for a reaction from the van't Hoff plot.

```
import pandas as pd
import matplotlib.pyplot as plt
from scipy.optimize import curve_fit
import numpy as np
from IPython.display import display
```

The equilibrium constant for a reversible reaction going from A to B was measured as a function of temperature.

The data obtained is given in the Excel document `deter_delta_h_data.xlsx`. Load the data with pandas.

```
from fysisk_biokemi.datasets import load_dataset
df = load_dataset('deter_delta_h') # Load from package for the solution so it doesn't
require to interact.
display(df)
```

	T_(K)	Keq
0	283	0.87
1	288	0.90
2	293	0.93
3	298	0.97
4	303	1.00
5	308	1.03

(a) Transform & plot the data

Tip

In NumPy the `np.log` function calculates the natural logarithm that we often write as $\ln$ and `np.log10` calculates the base 10 logarithm.

Add new columns to the DataFrame containing $\ln(K_{eq})$ and $1/T$ and make a plot with $1/T$ on the x-axis and $\ln(K_{eq})$ on the y-axis.


```
## Your task: Add a column containing 1/T and one containing ln(Keq) ##
df['1/T'] = 1/df['T_(K)']
df['ln(Keq)'] = np.log(df['Keq'])
```

Now you can make the plot in the cell below

```
fig, ax = plt.subplots()

## Your task: Make a plot of 1/T vs ln(Keq)
ax.plot(df['1/T'], df['ln(Keq)'], 'o')

## EXTRA: Customization you don't need to edit or understand this
## It adds axis labels to the plot you made.
ax.set_xlabel(r'1/T [K]')
ax.set_ylabel(r'$\ln(K_{eq})$');
```



(b) Fit trendline

Make a linear trendline by fitting a linear function to the data.

Start by defining a linear function

```
def vant_hoff(T, a, b):
    ## Your task: Implement a linear function
    result = a * T + b
    return result
```

Now we can use regression to find the parameters

```
## Your task: Make a fit with curve_fit to find the parameters ##
fitted_parameters, trash = curve_fit(vant_hoff, df['1/T'], df['ln(Keq)'])
a_fit, b_fit = fitted_parameters

## Print ##
print(a_fit)
print(b_fit)
```

-599.0182388271635
1.975814074075527

Then make a plot with the trendline added


```
fig, ax = plt.subplots()

## Your task: Choose points to evaluate the fit
## Look at the x-axis above and estimate a value for the minimum and maximum.
## Then calculate the fit at those points
max_x = 0.0032
min_x = 0.0036
T_trend = np.array([min_x, max_x])
y_fit = vant_hoff(T_trend, a_fit, b_fit)

## Your task: Plot the fit
ax.plot(T_trend, y_fit)

## Plots the data
ax.plot(df['1/T'], df['ln(Keq)'], 'o')

## EXTRA: You do not need to edit or understand this.
## It adds axis labels to the plot you made.
ax.set_xlabel('1/T [K]')
ax.set_ylabel(r'$\ln(K_{eq})$');
```



(c) Reaction type

Calculate the reaction enthalpy and entropy, at $T = 298$ K, assuming that $R = 8.314472 \times 10^{-3} \frac{\text{kJ}}{\text{mol}\cdot\text{K}}$. Is the reaction endo- or exothermic?


```
## Your task: Set known values and add units as comments e.g. # kJ/(mol K) ##
R = 8.314472 * 10**(-3) # Unit: # kJ/(mol K)
T = 298 # Unit: K

## Your task: Calculate ΔH and ΔS ##
## Consider the units!
dH = -R * a_fit # kJ/mol
dS = R * b_fit # kJ/(mol K)

print(dH)
print(dS)
```

```
4.980520374217764
0.016427850796106896
```

Tip

The units of a slope is the units of the y-axis divided by the units of the x-axis.

In the van't Hoff plot the y-axis is unitless and the x-axis is in 1/K.

(d) Free energy change

Calculate the free energy change, ΔG° , at $T = 298$ K, assuming that $R = 8.314472 \times 10^{-3} \frac{\text{kJ}}{\text{mol}\cdot\text{K}}$

```
## Your task: Calculate ΔG
dG = dH - T * dS

print(dG)
```


```
0.08502083697790841
```

8 ATP hydrolysis

```
import numpy as np
```

Below is shown the structure of ATP.

ΔG° for the hydrolysis of ATP to ADP and P_i is -30.5 kJ/mol.



(a) ΔG

In cells at 298 K the steady state concentrations of ATP, ADP og Pi resulted in a value for the reaction quotient $Q = \frac{[\text{ADP}][\text{Pi}]}{[\text{ATP}]} = 0.002$. Calculate $\Delta G'$ under these conditions.

```
## Your task: Define known values
## Add units as comments! E.g. # kJ / mol
Q = 0.002
T = 298 # K
dG_std = -30.5 # kJ / mol
R = 8.314 * 10**(-3) # kJ / (mol K)

## Your task: Calculate dG
dG = dG_std + R*T*np.log(Q)
print(dG)
```

-45.89713901562406

(b) **Equilibrium or not?**

Assuming that the temperature within these cells is 298 K, are these cells in equilibrium with respect to the hydrolysis of ATP?

```
K_eq = np.exp(-dG_std/(R*T))
print(K_eq)
```

222001.499219259

```
"""
```

```
Q is very different from this value, and therefore these cells are very far from
equilibrium with respect to hydrolysis of ATP. This value of Q further favors the
hydrolysis of ATP
```

```
"""
```

(c) ATP synthesized

Calculate the number of moles of ATP synthesized by a resting person in a period of 24 hours, assuming that the energy consumption by this person is 8368 kJ per 24 hours and that 40% of this energy is converted directly into ATP without loss of energy. Use the $\Delta G'$ value calculated in question a, and assume that 298 K.

```
# Your task: Set the known value of the energy spent for ATP.
energy_atp = 0.4 * 8368 # kJ

# Your task: Calculate the number of moles of ATP synthesized
synth_atp = energy_atp / np.absolute(dG)
print(synth_atp)
```

```
72.92829295657326
```



Tip

You can use `np.absolute` to calculate the absolute value of a number.

(d) Turnover

Next, calculate the average number of times each ATP molecule is turned over (synthesis followed by hydrolysis) in 24 hours when the total amount of ATP in the body equals 0.2 mol.

```
## Your task: Set the known total_atp value
total_atp = 0.2 # Unit: mol

## Your task: Calculate the turnover number.
turn_over = synth_atp / total_atp # Unit mol/mol -> Unitless
print(turn_over)
```

```
364.64146478286625
```