

---

# Index

- $2 \times 2$  table, 135, 137, 492–496, 546, 557, 666, 697  
Bayesian analysis, 642  
small-sample analysis, 494  
 $C_p$ , 404, 408, 413  
 $F$  distribution, 65–68, 76, 140, 486  
 $F$  statistic, 367, 378  
 $N(\mu, \sigma^2)$  distribution, *see* normal distribution  
 $O$  and  $o$  notation, 35  
 $p^*$  formula, 651, 665, 674  
 $t$  distribution, *see* Student  $t$  distribution  
 $\alpha$ -particle data, 696  
 $\chi^2$  distribution, *see* chi-squared distribution  
P-value, *see* significance level
- ABO blood group system, 137, 475  
accelerated life model, 541–553  
acceptance-rejection algorithm, *see* rejection algorithm  
adaptive rejection sampling, 82, 93, 626  
added variable, 414  
adjusted dependent variable, 474  
admissibility, 633  
age-specific failure rate, *see* hazard function  
AIC, 152, 235, 236, 308, 404, 407, 408, 413  
corrected, 403, 404, 524  
air-conditioning failure data, 696  
Akaike information criterion, *see* AIC 152  
Alofi rainfall data, 697  
analysis of covariance, 446, 448  
analysis of deviance, *see* deviance, analysis of  
analysis of variance, 378–386  
Latin square, 434  
one-way layout, 426  
random effects, 451  
split-unit experiment, 455  
two-way layout, 431, 437, 465  
ancillary statistic, 646–656, 693, 698  
Anderson–Darling statistic, 328  
ARCH process, 272  
ARIMA process, 271  
ARMA process, 270, 297, 697  
asymptotic relative efficiency, 51  
autocorrelation function, 267  
autoregressive moving average process, *see* ARMA process  
autoregressive process, 109, 267, 274, 618  
average, 15, 16, 24, 30, 41, 52, 54, 66, 67, 74, 75  
trimmed, 16, 86  
backfitting, 536, 623  
backshift operator, 270  
backward recurrence time, 298  
balanced incomplete block design, 432  
bandwidth, 307, 308, 520  
Barndorff–Nielsen’s formula, *see*  $p^*$  formula  
Bartlett adjustment, 340  
Basu’s theorem, 590, 649  
Bayes factor, 582–587, 593, 595, 596, 640  
approximate, 598  
Bayes information criterion, *see* BIC  
Bayes risk, 632  
Bayes rule, 631  
Bayes’ theorem, 11, 565–568  
Bayesian model averaging, 638  
beaver body temperature data, 266, 268, 698  
beetle data, 697  
belief network, 251  
Bernoulli distribution, 31, 89, 566, 568  
Bayesian analysis, 594, 637, 639  
information, 115  
Jeffreys prior, 575, 576  
sufficient statistic, 104  
Bernoulli trials, 570, 577  
best linear unbiased predictor (BLUP), 458, 463, 464, 467  
beta distribution, 89, 172, 219, 566  
beta function, 168  
beta-binomial distribution, 518, 636, 698  
bi-Weibull distribution, 189, 190  
bias, 300  
BIC, 152, 404, 599, 641  
binary data, 487–492, 517, 554, 684, 698  
complete separation, 489  
conditional inference, 694  
conditional likelihood, 665, 683  
deviance, 497, 559  
dichotomization, 488  
model checking, 490  
binomial distribution, 8, 61  
Bayesian analysis, 636  
confidence interval, 56, 58, 62, 345  
conjugate prior, 579  
cumulants, 49  
estimation, 314  
exponential family, 169, 180  
information, 110  
orthogonal parameter, 691  
Poisson approximation, 49  
relation to Bernoulli distribution, 30  
sufficient statistic, 315  
test, 91  
variance function, 481  
biological control data, 294  
birth data, 17, 19, 20, 23, 25, 26, 41, 54, 59, 60, 76, 83, 135, 177, 696  
birth order data, 158  
birth process, 288  
biweight, 375, 376  
Blalock–Taussig shunt data, 192, 198  
blocking, 419  
blood data, 450, 610  
blood group data, 175, 697  
boiling point data, 697

- bootstrap, 376
- Box–Cox transformation, 389, 391, 455, 551
- Box–Tidwell transformation, 412
- boxplot, 20
- branching process, 324
- breakdown point, 17, 27
- breast cancer data, 226, 230, 241
- Brownian bridge, 696
- Brownian motion, 92
- brush and spin plot, 696
- Burt twins data, 697
- cake data, 453
- calcium data, 469, 478, 678
- calibration, 415
- capture-recapture model, 106, 149
- cardiac surgery data, 579, 621
- case diagnostics, *see* model checking
- cat heart data, 446
- Cauchy distribution, 33, 48, 693, 698
  - information, 120
  - likelihood, 96, 100, 101, 127
  - simulation, 81, 91
- Cauchy–Schwarz inequality, 36
- causal inference, 423, 464
- cement data, 354, 379, 381, 385, 399, 408, 593, 697
- censoring, 4, 190, 217, 641
  - discrete data, 193
  - information, 112
  - left, 191
  - random, 190
  - right, 5, 112, 191
  - Type I, 190, 220
  - Type II, 190, 220
- central limit theorem, 30, 696
- Challenger data, 6, 97, 100, 122, 130, 139, 603, 697
- changepoint, 142, 583, 698
- characteristic function, 44, 48
- cherry tree data, 697
- chi-squared distribution, 63–64, 67, 76, 139, 219, 558
  - cumulants, 45
  - noncentral, 51
  - simulation, 78
- chi-squared statistic, 133
- chick bone data, 432, 697
- chimpanzee learning data, 485
- Cholesky decomposition, 89, 623
- classical inference, *see* repeated sampling
- clique, 245, 254, 255
- cloth data, 698
- cloth fault data, 514
- coal-mining disaster data, 698
- coefficient of variation, 51
- collinearity, 398, 697
- competing risks, 198–201, 218, 221
- completeness, 311, 315
  - bounded, 311, 340
- components of variance, 449–464
- computer bug data, 299, 643
- condition number, 398
- conditional inference, 143, 177
- conditional predictive ordinate, 589
- conditionality principle, 569, 639
- confidence interval, 54
  - equi-tailed, 56, 343
  - interpretation, 58
  - maximum likelihood estimate, 120
  - normal linear model, 371
  - one-sided, 56
  - Student  $t$ , 67, 90, 92
  - two-sample, 74
- confidence limit, 343
  - conservative, 345
- configuration, 186, 187, 650, 655
- confounding, 420, 442, 448, 466
- conjugate density, 573
- consistency, 29
  - strong, 123
- constructed variable, 391, 413, 487
- contingency table, 135, 500–507
- continuation ratio model, 510
- continuity correction, 671
- contrast, 443, 445, 465
- control variate, 85
- convergence, 28–37
  - diagnostics, 607, 638
  - in distribution, 30, 31
  - in probability, 28, 36
- Cook statistic, 362, 394, 396
  - approximate, 477
- correlation, 32, 36, 69, 90, 347
  - partial, 261, 264
- correlogram, 267, 297
  - partial, 267
- count data, 498–511
- counterfactual, 424
- counting process, 552
- covariance, 32, 36, 68
  - matrix, 68
  - partial, 261
- covariate, *see* explanatory variable
- coverage error, 345
- Cramér–Rao lower bound, 302, 319, 325, 377
  - multivariate, 304
- Cramér–von Mises statistic, 328
- credible set, 579, 594, 640, 641
  - highest posterior density (HPD), 579
- critical region, 333
  - invariant, 342
  - similar, 339, 665
  - unbiased, 337
  - uniformly most powerful, 336
- cross-validation, 308, 314, 395, 399, 408, 524, 533, 537
  - generalized, 399, 524, 525, 533, 537
- cumulant-generating function, 44–48, 167, 487, 671
- cut, 182, 501
- cycling data, 356, 362, 372, 388, 395, 444, 466
- daily rainfall data, 293
- Danish fire data, 277, 285, 328, 688
- data augmentation, 638, 698
- de Finetti’s theorem, 619
- decision rule, 631
  - minimax, 633, 637
- decision theory, 631–636, 638
- defective distribution, 189
- delta method, 33–35, 59, 122
  - several variables, 34
- dependent data, 323–324
- design matrix, 354
- detailed balance, 231, 238, 613
- deviance, 471, 483, 556, 559
  - analysis of, 484, 486
  - normal, 472
  - overdispersed, 515
  - penalized likelihood, 537
  - scaled, 471, 483
- diagnostic test, 567
- differencing, 271, 274
- Dirac comb, 310, 315
- Dirac delta function, 12

- directed deviance statistic, *see* signed likelihood ratio statistic
- directional data, 172
- Dirichlet distribution, 181
- discrimination, 631, 637
- dispersion parameter, 480, 487
- distribution constant, 184, 647, 649
- DNA data, 225, 230, 234, 236, 292
- double exponential distribution, *see* Laplace distribution
- dummy variable, 356
- Edgeworth series, 671, 672
- efficiency, 111
  - asymptotic relative, 303
- eigendecomposition, 230, 237, 238
- EM algorithm, 210–218, 223, 296, 297, 463, 563, 638, 697
- empirical Bayes, 627–638
- empirical distribution function (EDF), 19, 30, 277, 278
- empirical logistic transform, 36, 490, 509, 559
- endpoint, 146
- envelope simulation, *see* rejection algorithm
- equivalence relation, 107
- equivariant estimator, 185
- ergodic average, 230, 608
- ergodic model, 323
- error
  - Type I, 333
  - Type II, 333
- estimate, 23
- estimating equation, 316, 512
  - generalized, 507
- estimating function, 315–325, 555
  - optimal, 318
- estimation, 300–315
  - efficient, 303
  - non-regular, 304
  - unbiased, 300
- estimator, 23
- evolutionary distance data, 295
- exchangeability, 619, 626
- expectation space, 169
- expected information, 109–115, 124, 138, 144, 166, 179, 575
  - comparison with observed, 120
  - transformation, 156
- expert system, 293
- explanatory variable, 4, 161
- exponential distribution, 79, 119, 350, 680
  - Bayesian analysis, 639, 641
  - censored, 112, 220
  - conditional inference, 693
  - confidence interval, 314
  - estimation, 697
  - exponential family, 168
  - failure time, 6
  - grouped, 159
  - hazard, 188, 192
  - lack-of-memory property, 39
  - likelihood, 95, 125, 127
  - mixture, 149
  - nested in Weibull, 96, 130
  - order statistics, 39
  - orthogonal parameter, 691, 695
  - probability plot, 26
  - shifted, 145, 149, 350
  - simulation, 78, 89, 91
  - sufficient statistic, 105, 108
  - test, 326, 344
  - truncated, 698
- exponential family, 166–183, 215, 218, 336, 340, 350, 493, 636, 680
  - $(p, q)$ , 174
  - complementary mean parameter, 689
  - completeness, 312
  - conditional density, 180
  - conditional inference, 674
  - conjugate prior, 573, 577, 578, 639
  - curved, 174, 182, 677
  - inference, 176
  - likelihood, 179
  - linear, 490
  - marginal density, 180
  - minimal representation, 172
  - natural, 167, 172
  - order  $p$ , 171, 176, 573
  - order 1, 167, 168
  - regular, 167
  - steep, 170, 220
- exponential scores, 26, 40
- exponential tilting, 167, 168
- eye data, 505
- factor, 356
  - crossed, 452
- factorial experiment, 356, 391, 436, 439, 441, 442, 444, 448, 697
  - replicated, 436
- factorization criterion, 104, 410, 566
- field concrete mixer data, 434, 445, 448
- financial data, 33
- fir seedling data, 640
- first-passage time, 229, 243
- Fisher information, *see* expected information
- Fisher scoring, 118
- fitted value, 361, 362
- force of mortality, *see* hazard function
- forensic evidence, 584
- forward recurrence time, 298
- frailty, 201–202, 218, 221, 555, 563
  - shared, 562
- frequentist inference, *see* repeated sampling
- FTSE data, 266, 271, 273, 697
- full conditional density, 245, 605
- funnel plot, 209
- galaxy data, 213
- gamma distribution, 23, 64, 487
  - cumulants, 48
  - estimation, 57, 696
  - exponential family, 181, 182, 219
  - generalized, 132
  - hazard, 190
  - information, 115
  - inverse, 182
  - probability plot, 26
  - simulation, 89
  - small-sample inference, 674
  - test, 339
- gamma function, 23, 617
  - properties, 27
- GARCH process, 273
- Gauss–Markov theorem, 374
- Gaussian distribution, *see* normal distribution
- generalized additive model, 538, 541, 555, 623, 698
- generalized extreme-value distribution, 50, 279, 291
- generalized linear model, 480–487, 518, 541, 554–556, 558
- generalized Pareto distribution, 284, 286, 291, 292, 299, 317, 688, 697
- genetic linkage data, 223
- genetic pedigree, 249
- geometric distribution, 229
  - Bayesian analysis, 639, 640
  - exponential family, 219
  - information, 115

- likelihood, 101
  - relation to negative binomial, 50
  - simulation, 89
- Gibbs sampler, 605–612, 618, 621, 638, 642, 643, 698
- goodness of fit, 131–138, 177, 327
  - posterior predictive, 592
- Graeco-Latin square, 466
- graph
  - ancestral subset, 255
  - directed acyclic, 249–253, 255
  - moral, 250, 251, 255, 262, 265, 296
- graphical design, 21, 28
- graphical model, 260–266, 292
- Greenwood's formula, 197
- group, 183
- group action, 183
- group transformation model, 183–188, 218, 329
  - composite, 187
- grouped data, 368, 414
- Gumbel distribution, 203, 279, 297, 413, 475
- half-normal distribution, 79, 696
- half-normal plot, 444
- Hammersley–Clifford theorem, 246, 253, 255, 292, 296, 605
- hat matrix, 362, 369, 385, 413
- hazard function, 188, 203, 275, 286
  - bathtub, 190
  - cause-specific, 198, 221
  - cumulative, 189
  - estimation, 220
- head size data, 697
- Heaviside function, 12
- Hermite polynomial, 672
- hierarchical model, 464, 638
  - Bayesian, 619–627
  - Poisson, 600, 698
- histogram, 19, 305, 349
- Hotelling's  $T^2$  statistic, 260
- Huber estimator, 321, 325, 350, 375, 376
- human lifetime data, 194
- HUS data, 142, 177, 583, 697, 698
- hypergeometric distribution, 495, 557
- hyperparameter, 573
- hypothesis
  - alternative, 326
  - composite, 326, 339–343
  - null, 325
  - simple, 326
- hypothesis test, 325–348, 582
  - comparison, 333
  - invariant, 342
  - nonparametric, 331
  - one-sided, 329, 337, 350
  - randomized, 336, 347
  - relation to confidence interval, 343–346
  - similar, 339
  - two-sided, 329, 337
- Hölder's inequality, 182
- ignorable non-response, 204
- image analysis, 245, 292
- imaginary observations, 596
- importance sampling, 87, 618, 641
  - Bayesian application, 602–605
  - ratio estimator, 603
  - raw estimator, 87
  - weight, 87
- incidence matrix, 533
- inference function, *see* estimating function
- infinitesimal generator, 238
- influence, 394, 477, 539
- influence function, 321
- information
  - expected, 222
- information distance, *see* Kullback–Leibler discrepancy
- information sandwich, 147, 151, 377
- intensity function
  - complete, 286
  - conditional, 288
- interaction, 424, 436, 439, 466
  - first-order, 440
- interest-preserving reparametrization, 645
- interquartile range (IQR), 17, 20, 37, 43, 61
- interval estimation, 313–314
- invariant, 184
  - maximal, 184, 186, 329, 343
- inverse gamma distribution, 580, 588, 640
- inverse Gaussian distribution, 182, 487, 680
- inverse probability, 637
- inversion algorithm, 78–79, 89
- IQR, *see* interquartile range
- Ising model, 248
- iterated expectation, 65
- jacamar data, 470, 483, 502
- Japanese earthquake data, 288, 518, 525
- Jeffreys prior, 639
- Jeffreys–Lindley paradox, 586
- Jensen's inequality, 123
- Kaplan–Meier estimator, *see* product-limit estimator
- kernel, 306, 520
  - effective, 521
  - tricube, 520
- kernel density estimation, 697
- kernel density estimator, 305–309, 314, 608
- Kolmogorov–Smirnov statistic, 328
- Kronecker delta, 12
- Kullback–Leibler discrepancy, 123, 147, 150
- kurtosis, 46
- Lake Constance data, 697
- Laplace approximation, 596–602, 617, 636, 638, 641
  - posterior density, 599
  - posterior distribution, 599
- Laplace distribution, 22, 24, 28, 85, 125, 157, 698
  - regression model, 377
- Laplace transform, 312
- large deviation region, 652
- Latin square, 434, 438, 446, 466, 697
- law of small numbers, *see* Poisson approximation to binomial distribution
- least squares estimation, 163, 359–369, 697
  - generalized, 369
  - geometrical interpretation, 362, 369, 378
  - iterative generalized (IGLS), 463
  - iterative weighted, 472–476, 479, 554–556
  - penalized, 560
  - restricted iterative generalized (RIGLS), 659
  - robustness, 376
  - weighted, 368, 369, 409, 514

- least trimmed squares estimation, 376
- leukaemia data, 541, 545, 697, 698
- leverage, 362, 393, 394, 476, 539
- Lexis diagram, 191, 218, 561
- likelihood, 94
  - basic properties, 99–100
  - complete-data, 210, 215
  - conditional, 557, 646, 665, 677, 683, 694
  - dependent data, 98
  - exponential family, 179
  - interpretation, 100–101
  - local, 527, 540
  - log, 99
  - marginal, 646, 656, 665
  - modified, 458
  - modified profile, 680–691, 694
  - non-regular properties, 140–148, 697
  - observed-data, 210
  - partial, 544, 545, 554, 561, 656, 665
  - penalized, 156, 531, 535, 555
  - profile, 117, 140, 544, 680, 694
  - pure, 101
  - quadratic summary, 109, 125
  - relative, 99, 102, 109, 119
  - reparametrization, 99, 116
  - restricted, 458, 657, 690
  - summary, 101
- likelihood equation, 116
- likelihood principle, 568–571, 589, 638, 639
- likelihood ratio statistic, 126–139, 330, 340, 366
  - generalized, 128
  - large-sample distribution, 126, 138
  - signed, *see* signed likelihood ratio statistic
- linear congruential generator, 77, 696
- linear exponential family
  - conditional inference, 683
  - modified profile likelihood, 682
  - orthogonal parameter, 689
- linear mixed model, 456, 463, 657
- linear model, 353–417, 479, 554, 661
  - Bayesian analysis, 641
  - normal, 359, 370–374
  - terms, 380
- linear predictor, 480
- linear process, 269
- link function, 480
  - binary data, 488, 497
  - canonical, 482, 487
  - complementary log-log, 488, 497
  - identity, 482
  - inverse, 482, 486
  - log, 482, 486
  - log-log, 488, 497
  - logit, 8, 484, 488, 490, 497
  - probit, 488, 558
- lizards data, 697
- local characteristic, *see* full conditional density
- local polynomial estimation, 519–530, 539, 555
  - bias and variance, 522, 529
  - degrees of freedom, 521
  - smoothing parameter, 523
- location, 15, 27, 61
- location model, 183, 649, 655
- location-scale model, 61, 157, 185, 187
  - Bayesian analysis, 639
  - conditional inference, 694
  - goodness of fit, 588
  - Jeffreys prior, 576
  - orthogonal parameter, 695
- log odds, 169
- log rank test, 545
- log-concave density, 81
- log-linear model, 498–500, 554, 556, 559, 697
- log-logistic distribution, 190
- log-normal distribution, 37, 91, 190, 303, 304
  - probability plot, 26
- logistic distribution, 156, 202, 316
- logistic regression, 97, 100, 122, 130, 490–492, 498, 505, 509, 515, 636, 665, 676, 683, 684, 697, 698
  - sufficient statistic, 108
- longitudinal data, 456, 457, 555, 660
- look-up method, 78
- loss function, 631
- lowess, 521
- lung cancer data, 8, 503, 693
- M-estimator, 375, 376
- MAD, *see* median absolute deviation
- magical mystery formula, 651
- magnesium data, 208
- main effect, 440
- maize data, 1, 67, 68, 74, 129, 130, 140, 309, 329, 332, 365, 372, 381, 386, 410, 580
- Manaus river height data, 697
- Mantel–Haenszel test, 558
- marginal model, 505, 554, 559
- Markov chain, 225–245, 292, 606, 697
  - classification of states, 229
  - continuous-time, 237
  - first-order, 234, 247, 293, 660
  - geometrically ergodic, 231
  - inhomogeneous, 242
  - reversible, 231, 238
  - second-order, 236, 254, 293
  - simulation, 244
  - stationary, 228
  - two-state, 231, 240
  - variable-order, 235
  - zeroth-order, 234
- Markov chain Monte Carlo, 605–617, 638, 642, 666
  - output analysis, 607
- Markov process, 98, 267, 268, 273
  - continuous-time, 294, 295
- Markov property, 98, 228, 244
  - global, 251, 262, 296
  - local, 251, 253, 254, 296
  - pairwise, 296
- Markov random field, 244–255, 292, 293, 622, 626
- martingale, 323, 552
- masking, 388
- matched pairs, 372
- maths marks data, 256, 259, 261, 263
- max stability, 291
- maximum likelihood estimator, 102, 115–126, 210, 324, 346
  - computation, 115
  - conditional, 677
  - consistent, 122
  - large-sample distribution, 118, 124
  - usual regularity conditions, 118
- mean, 22
- mean excess life function, 203
- mean parameter, 169
- mean residual life plot, 285
- mean squared error, 300–305
  - integrated, 308
- measuring machines, 569
- median, 331
- median absolute deviation (MAD), 17, 28, 43
- meta-analysis, 206, 223
- method of moments, *see* moment estimator

- Metropolis–Hastings algorithm, 612–617, 626, 638, 642, 667, 698
  - random walk, 613, 618, 643
- Michaelis–Menton model, 695
- midrange, 50
- millet plant data, 697
- minimal representation, 172, 182
- minimax decision rule, 633
- minimum variance unbiased estimation, 309–313
- missing at random (MAR), 204, 205, 222
- missing completely at random (MCAR), 204, 205
- missing data, 203–218
- missing information principle, 211
- mixture distribution, 213, 219, 639
- mode, 27
- model
  - averaging, 407, 592
  - nested, 127, 131, 133, 139, 586
  - non-linear, 503
  - parametric, 22
  - saturated, 471
  - selection, 150–155
  - uncertainty, 83, 406
  - wrong, 147, 377
- model building
  - linear model, 397–408
- model checking, 476–479
  - Bayesian, 587–592
  - linear model, 386–397
- moderate deviation region, 652
- moment estimator, 316, 636
- moment-generating function, 37, 44, 481, 487
- moments, 44–48
- Monte Carlo integration, 87
- motorette data, 615, 698
- mouse data, 200, 546
- moving average process, 269, 274, 297
- multilevel model, 461, 464
- multimodal distribution, 642
- multinomial distribution, 37, 139, 475, 654
  - cumulants, 47, 51
  - estimation, 697
  - exponential family, 175, 220
  - fit, 133
- multiplicative model, 359
- multivariate  $t$  distribution, 297
- multivariate normal distribution, 138, 247, 255–267
- Nadaraya–Watson estimator, 522, 530, 540
- natural cubic spline, 530, 533, 535, 560
- natural observation, 168
- natural parameter, 168
- negative binomial distribution, 630
  - exponential family, 219
  - genesis, 50
  - likelihood, 517
  - orthogonal parameter, 691
- neighbourhood system, 244–246, 248
- nested variation, 450
- network information criterion, 152
- neurological data, 697
- Newton–Raphson algorithm, 116, 211, 213, 215, 217, 473
- Neyman–Pearson lemma, 335, 632
- nodal involvement data, 490, 676, 684
- non-additivity test, 390, 391, 486
- non-ignorable non-response (NIN), 204, 205
- nonlinear model, 678, 695
- nonlinearity, 389
- normal distribution, 62–63, 159, 347, 481, 646
  - Bayesian analysis, 580, 640
  - bivariate, 70, 71, 90, 207, 608
  - confidence interval, 121
  - conjugate prior, 574
  - cumulants, 45, 70
  - empirical Bayes analysis, 627
  - exponential family, 178
  - extremes, 280
  - goodness of fit, 588
  - information, 111
  - Jeffreys prior, 577
  - likelihood, 116, 129, 180
  - linear combination, 45, 72–73, 90, 92, 164, 165
  - mixture, 85, 145, 149, 644, 697
  - modified profile likelihood, 690
  - multivariate, 68–77, 89, 90, 220, 259–266, 697
  - orthogonal parameter, 689
  - risk, 633
  - rounding, 114, 616
  - sample median, 41
  - sampling, 613, 642
  - simulation, 78, 80, 91, 696
  - standard, 63
  - sufficient statistic, 109, 125
  - test, 139, 140, 159, 333, 337
  - trivariate, 72, 73
  - unbiased estimation, 301, 312
- normal equations, 360
- normal hierarchical model, 620
- normal linear model, 474, 479, 593, 681
  - Bayesian analysis, 589, 595
- normal nonlinear model, 474, 479
- normal scores plot, 26, 387
- notation, 12
- nuclear plant data, 401, 404, 664
- null distribution, 326
- observational study, 10, 648
- observed at random (OAR), 217
- observed information, 102, 109–115, 138, 144, 166, 179
  - transformation, 156
- Ockham's razor, 150, 378
- offset, 498
- one-way layout, 426, 438, 449, 459, 465, 467
- opinion polling, 56, 58
- orbit, 183
- order statistic, 37–44, 106, 186, 190, 220, 296, 350
  - extreme, 41
  - summary, 16
- order statistics, 276, 279
- ordinal response, 507–510, 555
- orthogonal polynomials, 383, 445, 697
- outlier, 17, 149, 320, 388, 697
- overdispersion, 177, 511–518, 527, 698
- paired comparison, 3, 140, 419, 421, 425
- panel data, 225
- parameter, 3, 22
  - identifiable, 144
  - interest, 127, 645
  - nuisance, 127, 645
  - orthogonal, 487, 685–690
  - redundant, 144, 149
  - space, 94, 140

- parametrization, 23
  - corner-point, 440
- Pareto distribution, 41, 348, 594
- partial likelihood, 561
- partial spline model, 537
- PBC data, 549, 698
- pea data, 160
- Pearson's statistic, 135, 140, 160, 177, 234, 237, 483, 485, 497, 517, 696
- people data, 696
- permutation group, 184
- permutation test, 341, 352
- pig diet data, 431, 438
- pigeon data, 172
- pivot, 53, 61, 67, 313, 343
  - approximate, 56, 74
  - basis of test, 60
  - exact, 139
  - properties, 56
  - Student  $t$ , 66
  - two-sample, 74
- plotting position, 26
- pneumoconiosis data, 508
- point process, 274–293
  - clustering, 277
  - length-biased sampling, 298
  - marked, 288
  - orderly, 275, 286
  - self-exciting, 289
  - spatial, 293
  - thinning, 291
- poisons data, 391, 436, 440, 464
- Poisson approximation to binomial distribution, 49, 282
- Poisson birth process, 98, 108, 146
- Poisson dispersion test, 177
- Poisson distribution, 9, 23, 49, 142, 511
  - Bayesian analysis, 626, 637, 640, 644
  - complete, 311
  - conditional inference, 694
  - confidence interval, 61, 160, 697
  - conjugate prior, 573, 577
  - cumulants, 46
  - estimation, 696
  - exponential family, 170, 177, 340, 481
  - goodness of fit, 135
  - likelihood, 94
  - marginal inference, 665, 693
  - mixture, 697
  - orthogonal parameter, 690
  - saddlepoint approximation, 670
  - sufficient statistic, 109
  - truncated, 37, 696
  - unbiased estimation, 310, 313, 315
  - variance stabilization, 59
- Poisson process, 40, 274–287, 293, 486, 498, 562, 693, 697
  - Bayesian analysis, 596, 643
  - empirical Bayes analysis, 629
  - estimation, 696
  - homogeneous, 277, 285
  - information, 112
  - inhomogeneous, 283, 299, 557, 643, 697, 698
  - intensity, 275
  - simulation, 298
- pollution data, 697
- polynomial regression, 354
- positive stable distribution, 563
- positivity condition, 246, 253, 254
- positron emission tomography, 216
- posterior density, 566
  - marginal, 578
  - normal approximation, 578
- posterior predictive density, 568, 577, 602, 617
  - normal linear model, 591
  - Poisson distribution, 577
- potential, 246
- power, 333
  - local, 338
- prediction, 60–61, 150, 568, 592
- prediction decomposition, 98
- prediction interval, 60, 165
  - normal linear model, 371, 372
- Premier League data, 498
- principal components, 397
- principle of insufficient reason, 577, 637
- principle of parsimony, *see* Ockham's razor
- prior density, 566, 572–577, 638
  - conjugate, 567, 573, 640
  - elicitation, 638
  - ignorance, 574
  - improper, 574, 580, 640
  - Jeffreys, 575
  - non-informative, 574
- probability integral transform, 39
- probability plot, 26, 28, 131, 203
  - exponential, 159, 277, 278, 286, 696
  - Gumbel, 297
  - half-normal, 696
  - normal, 49, 63, 92, 165, 179, 696
  - Weibull, 50
- probability weighted moment estimators, 317
- product moment correlation coefficient, *see* correlation
- product-limit estimator, 196–198
- profile likelihood, 127–131, 479
- proportion data, 487–498
- proportional hazards model, 543–555, 562, 563, 656, 698
- proportional odds model, 508
- prospective study, 493
- pseudo-random numbers, 77–78
- publication bias, 206–210
- pump failure data, 160, 600, 644, 698
- quantile, 22
- quantile-quantile (Q-Q) plot, 26, 28
- quartile, 16
- quasi-likelihood, 512–517, 555, 698
- quasi-random numbers, *see* pseudo-random numbers
- random effects model, 449, 456, 458, 610
- random sample, 21–24
  - normal, 66–68
- randomization, 417–426
  - distribution, 422
- randomized block design, 429
- rank statistic, 561
- Rao–Blackwell theorem, 309
- rare events, *see* statistics of extremes
- rat growth data, 459
- ratio, 34
- ratio of uniforms algorithm, 81, 91
- regression-scale model, 661–665, 681, 691
- rejection algorithm, 79–82, 89
  - adaptive, 81
- relevant subset, 647
- renewal process, 287
- repeated measures, 456
- repeated sampling, 52, 58, 119
- replication, 464
- residual
  - binary response, 492
  - Cox–Snell, 541, 548
  - deletion, 362, 395, 590, 591
  - deviance, 477, 517, 548, 551
  - martingale, 548, 551, 552

- Pearson, 517
- properties, 386
- raw, 165, 179, 362, 370, 387, 554, 588
- serial correlation, 387
- standardized, 362, 387, 396, 414, 479, 590, 649
- standardized deviance, 477, 479, 485, 556
- standardized Pearson, 477, 479, 556
- time series, 269, 274
- residual sum of squares, 163, 361, 371
- resistant statistic, 17
- response, 161
- restricted maximum likelihood
  - estimation (REML), 458, 461, 464, 467, 659, 665
- retrospective study, 493
- return level, 280, 688
- return period, 280
- reversibility, 607
- ridge regression, 398, 697
- risk function, 632
- risk set, 193, 543
- robustness, 319–322
- roughness penalty, 216, 530–535
- rounding, 113, 115, 145
- rug, 19
- run, 229, 243
- Rényi representation, 40
- saddlepoint approximation, 560, 638, 668–673, 680, 698
  - double, 670, 675
- saddlepoint equation, 669
- salinity data, 697
- sample, 15
  - average, 51
  - maximum, 16, 50, 279, 291
  - mean, *see* average
  - median, 16, 20, 37, 41, 51, 61, 157, 324
  - minimum, 16, 41, 42, 50, 279
  - moment, 15, 24, 75
  - quantile, 16
  - range, 50
  - shape, 18, 28
  - skewness, 28
  - space, 94
  - space derivative, 652, 682
  - variance, 15, 25, 28, 31, 50, 66–68, 74, 75
- sampling variation, 24–25
- sampling-importance resampling (SIR), 618
- sandwich covariance matrix, 507, 513
- scale, 15, 27, 61
  - choice, 58
- scale model, 347, 639
- scatterplot, 4, 20
  - matrix, 256
- score statistic, 116, 138, 144, 149, 315, 338, 346
- score test, 132, 338
- seed, 78
- seed germination data, 698
- selection bias, 210, 218
- self-consistency, 223
- semiparametric regression, 518–540, 555
- Shakespeare's vocabulary data, 629
- shoe data, 421
- short time series, 659
- shrinkage, 459, 621, 625, 628, 634
- sign test, 331, 332, 334
- signed likelihood ratio statistic, 128, 346
  - modified, 653, 663, 676
- significance level, 325, 582
  - conflict with likelihood principle, 570
  - interpretation, 326
  - mid- $p$ , 495, 671
- significance trace, 525
- simple random sample, *see* random sample
- Simpson's paradox, 256–258
- simulation, 77–90
- size, 333
- skew-normal distribution, 91, 160
- skewness, 18, 46
- slash distribution, 85
- Slutsky's lemma, 31–33
- smoking and the Grim Reaper, 258, 494
- smoother, 518
  - linear, 532, 539
- smoothing matrix, 521, 537
- spacing, 43, 277
- spectral decomposition, 73, 397
- speed limit data, 697
- speed of light data, 696
- spline, 555
- split-unit experiment, 452
- spring barley data, 533, 538, 622, 626
- spring failure data, 4, 95, 96, 100, 119, 120, 127, 130, 132, 154
- standard error, 52
- stationarity, 267, 607, 641
  - second-order, 267
  - strict, 267
- statistic, 15
- statistical formulae
  - mindless repetition, 88
- statistical genetics, 292
- statistics of extremes, 278–286, 293
- Stein effect, 635
- stem-and-leaf display, 28
- Stirling's formula, 617
- stochastic matrix, 229
- straight-line regression, 115, 159, 161–166, 186, 219, 220, 317, 322, 353, 354, 361, 394, 410, 412, 413, 697
- Student  $t$  distribution, 64–65, 74, 76, 85, 140, 187, 651
  - linear model, 374
  - simulation, 78
- Student  $t$  statistic, 67, 129, 139, 140, 164, 368, 696
  - coverage, 696
  - regression, 379
  - robustness, 84
- Student  $t$  test, 330, 332, 341, 342
- studentized statistic, 32
- sufficiency principle, 569, 639
- sufficient partition, 107, 109
- sufficient statistic, 103–108, 176
  - minimal, 107, 166, 410, 566
- sum of squares, 163, 360, 380
  - orthogonal, 382
  - penalized, 532
- surrogate variable, 564
- survival data, 188–203, 218, 376, 540–554
- survivor function, 203
- Swan of Avon, 629
- symbolic rank deficiency, 149
- teaching methods data, 427
- teak plant data, 697
- test, 60
  - Monte Carlo exact, 668, 680
  - unimodality, 697
- test statistic, 325
- threshold stability, 291

- time series, 266–274, 293
- time-dependent covariate, 547, 562
- Titanic data, 697
- tolerance distribution, 488, 508
- tolerance interval, *see* prediction interval 60
- toxoplasmosis data, 515, 527, 636
  - empirical Bayes analysis, 628
- transformation, 58, 122, 697
  - exponential, 34
  - interest-preserving, 129
  - symmetrizing, 558
  - variance-stabilizing, 59
- transition matrix, 229
- transition probability, 228
- trinomial distribution, 508
- two-sample model, 3, 73–75, 140, 341, 365, 372, 419, 425
  - Bayesian analysis, 595
- two-way layout, 429, 464, 485, 538
- ulcer data, 495, 668, 697
- unemployment rate, 61
- uniform distribution, 669
  - Bayesian analysis, 594
  - exponential family, 180
  - exponential tilting, 167, 170
  - likelihood, 103, 109, 149
  - not complete, 312
  - order statistic, 43
  - order statistics, 38, 50, 51
  - simulation, 77
  - unbiased estimation, 304, 315
- unit-treatment additivity, 421, 424
- urine data, 698
- variability band, 525
- variable selection, 400
  - $C_p$ , 403, 412
  - backward elimination, 400, 408, 412
  - forward selection, 400, 408, 412
  - likelihood criteria, 402
  - stepwise, 400, 408, 412, 697
- variance function, 59, 170, 182, 481, 512
  - linear, 171, 511
  - quadratic, 511, 517
- variance reduction, 85–89
- variance-stabilizing transformation, 170
- variance-time curve, 288, 298
- Venice sea level data, 161, 164, 165, 186, 205, 465, 475, 477
- volatility, 272
- von Mises distribution, 172, 174
- weak law of large numbers, 28, 152
- Weibull distribution, 50, 553
  - Bayesian analysis, 615
  - estimation, 697
  - hazard, 189
  - information, 157
  - likelihood, 96, 100, 117, 125, 127, 130, 154
  - moment estimation, 319
- weight of evidence, 583
- white noise process, 267
- Wiener process, 696
- Wilcoxon signed-rank test, 331, 332, 351
- Wilcoxon two-sample test, 351
- Wishart distribution, 260
- Yahoo share price data, 92
- Yarmouth sea level data, 281