

# **Establishment of Self-sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia**

Aubrey Moore PhD

University of Guam College of Natural and Applied Sciences

April 23, 2020

# Contents

|   |           |
|---|-----------|
| <b>1. Project Description</b>   | <b>3</b>  |
| 1.1. Background . . . . .   | 3         |
| 1.2. Statement of Need . . . . .  | 6         |
| 1.3. Goals and Objectives . . . . .   | 7         |
| 1.3.1. Objective 1: Survey to Determine Background OrNV Incidence . .                                     | 7         |
| 1.3.2. Objective 2: Establish Sustainable CRB-G Biocontrol by Autodis-<br>semination of OrNV . . . . .    | 7         |
| 1.3.3. Objective 3: Establish Island-wide Monitoring Systems for CRB and<br>Coconut Palm Health . . . . . | 8         |
| 1.4. Timeline . . . . .   | 9         |
| 1.5. Potential Benefits . . . . .   | 10        |
| <b>2. Budget</b>  | <b>11</b> |
| <b>3. References</b>  | <b>12</b> |
| <b>Appendix A. Supporting Technical Data</b>  | <b>14</b> |
| A.1. Self-sustaining Biological Control of CRB in Fiji Using OrNV . . . . .                               | 14        |
| A.2. Automated Monitoring of CRB Damage Using Roadside Video Surveys . .                                  | 15        |
| <b>Appendix B. DOI-OIA Grant D17AP00107 Progress Report 4</b>   | <b>16</b> |
| <b>Appendix C. Extracts from the 22nd Micronesian Islands Forum Joint Commu-<br/>        nique</b>        | <b>28</b> |
| <b>Appendix D. Letters of Support</b>   | <b>32</b> |
| D.1. Guam Invasive Species Council . . . . .  | 32        |
| D.2. Regional Invasive Species Council for the Western Pacific . . . . .                                  | 34        |
| <b>Appendix E. Cover Letter</b>   | <b>36</b> |

# 1. Project Description

This grant proposal is a request for funds to support a one-year project with a major goal of establishing effective, self-sustaining biological control of the coconut rhinoceros beetle (CRB) on Guam. *Oryctes rhinoceros*.

## 1.1. Background

**The coconut rhinoceros beetle, *Oryctes rhinoceros*, is a major pest of coconut palm, oil palm and other palm species.** Palms are damaged when adult beetles bore into the crowns of palms to feed on sap. Tree mortality occurs when beetles destroy the growing tip (meristem). Immature beetles (grubs) do no damage. They feed on dead, decaying vegetation in breeding sites. Preferred breeding sites are dead, standing coconut stems, and piles of decaying vegetation such those left behind by typhoons or after re-planting of oil palm plantations. If a CRB population is not suppressed, it is possible for a positive feed-back cycle to initiate whereby adult beetles kill massive numbers of palms, thereby generating more food for even more grubs which turn into adults which kill even more palms. An outbreak following this scenario occurred in the Palau Islands during the late 1940s resulting in about 50% coconut palms being killed by CRB throughout the archipelago and 100% mortality on some of the smaller islands (Gressitt 1953). A similar outbreak, initiated by Typhoon Dolphin (2015), is currently impacting Guam.

**Following 40 years of no geographical range expansion, CRB is again “on the move” in the Pacific.** CRB was recently detected for the first time at several Pacific Island locations including Saipan (2006), Guam (2007), Port Moresby, Papua New Guinea (2010), Oahu, Hawaii (2013), Honiara, Solomon Islands (2015), Rota, CNMI (2017), and Aguiguan, CNMI (2019).

Eradication of CRB has been attempted many times but is extremely difficult, having been achieved only once, on Niuatoputapu (formerly known as Keppel Island), a tiny island belonging to Tonga, with an area of only 16 km<sup>2</sup> (3% the area of Guam) (Catley 1969).

Failing eradication, the usual response to CRB infestations during the second half of the 20th century was introduction of *Oryctes* nudiviruses (OrNV), the biological control agent of choice for this pest Jackson 2009 . OrNV attacks only CRB, typically reducing damage by up to 90% with population suppression lasting indefinitely (Bedford 2013). OrNV is auto-disseminated, meaning the pathogen is carried between feeding and breeding sites by CRB adults. Like many biocontrol agents, OrNV is density-dependent, working best at high population densities. After release, OrNV sustains itself within the CRB population, limiting damage to very low levels (See Appendix A.1: Self-sustaining biological control of CRB in Fiji using OrNV).

**Current invasions of Pacific Islands by CRB involve a new invasive biotype that has escaped from biological control by OrNV.** Discovery of OrNV nudivirus in the 1960s enabled the successful management of CRB populations in Pacific Island Countries (Huger 2005). Augmentative release of OrNV continues to be an important mechanism for CRB management in both coconut and oil palm growing regions. For about 40 years after adoption of this biocontrol strategy, no new outbreaks of CRB were reported from uninfested palm growing islands in the Pacific ensuring continuity of palm based village economies.

However, the situation has recently changed. For the first time in 40 years, CRB invasion into completely new areas has been reported. Additionally, Pacific areas with established CRB populations (e.g. Palau) have reported increased severity and frequency of CRB damage. Common to all these areas is the high incidence of severe palm damage by beetles not seen since the introduction of OrNV.

Initial attempts to introduce OrNV into the Guam CRB population were unexpectedly unsuccessful, raising the possibility that the population that invaded Guam is tolerant or resistant to the commonly applied OrNV isolates. Subsequent DNA analysis showed that the Guam population is genetically different from other populations in the region. On the basis of distinct genetics and tolerance to currently available OrNV isolates, the Guam population has been designated a new biotype, CRB-Guam (CRB-G) (Sean David Goldie Marshall et al. 2015; Sean D. G. Marshall et al. 2017).

DNA analysis from an ongoing survey has detected the CRB-G biotype in Guam, Hawaii, Palau, Port Moresby (PNG) and Honiara (Solomon Islands). Thus, current invasions in the Pacific involve the CRB-Guam biotype and it is expected that these populations are tolerant isolates of OrNV previously used as biocontrol agents. However, Recent work has identified OrNV isolates which are new biocontrol candidates for CRB-G. (See the **Recent progress** section below [1.1].)

**Uncontrolled CRB-G outbreaks on islands may kill most palms within a few years and risk of accidental spread to other islands is high.** A worse case scenario for a CRB infestation may be triggered by a massive outbreak of adult CRB emerging from abundant breeding sites made by large amounts of decaying vegetation left in the wake of a typhoon, from large scale land clearing or large environmental destruction during a war. The current uncontrolled outbreak on Guam was initiated by Typhoon Dolphin which visited Guam in May, 2015. Massive amounts of decaying vegetation left in the wake of this storm provided abundant CRB breeding sites. Very high feeding activity by adults emerging from these breeding sites killed mature coconut palms, leaving standing dead coconut trunks which became ideal breeding sites for subsequent generations of beetles.

During a severe CRB outbreak, there will be an increased risk of further spread to uninfested islands throughout the Pacific. Palms are important on Pacific Islands for various reasons: as a cash crop for nuts, oil and lumber, as an ornamental tree appreciated by residents and tourists. On some of the smaller, more traditional islands, coconut palm is referred to as *the tree of life*. On these islands, this species is an essential natural resource providing income, housing, food, oil, soap, clothing, mats, baskets, and other containers. The smaller, poorer Pacific islands will suffer the most if spread of CRB-Guam cannot be controlled. If CRB-G infests islands and atolls where the coconut palm as the *tree of life*, islanders may have to migrate to larger population centers.

**Recommended response to CRB-G invasions.** Entomologists working on the CRB-G problem agree that the most feasible way to prevent massive palm mortality during outbreaks is establishment of biological control using an isolate of OrNV which is highly pathogenic to CRB-G (Jackson 2015; Vaqalo et al. 2015; Secretariat of the Pacific Community 2017).

The consensus among Pacific-based entomologists is that the most feasible way to stop massive palm mortality during CRB-G outbreaks is to find a find and release a have met several times to plan a response to CRB-G invasions. In a special meeting on CRB-G at the XXVth International Congress of Entomology , the following strategic plan was suggested:

A coordinated regional project should be organized and adequately staffed and funded to accomplish 3 objectives:

1. Survey CRB populations throughout the Asian/Pacific region to delimit the geographical distribution of CRB-G and identify its centre of origin.
2. Survey CRB-G populations from the centre of origin to find isolate(s) of OrNV (or other pathogens) that are highly pathogenic for the CRB-G biotype.
3. Implement *in vivo* or *in vitro* propagation of selected OrNV isolates for auto-dissemination on islands infested with CRB-G.

The CRB-G problem is not limited to American-affiliated islands. Attempts to find financial support for a well-coordinated Pacific-wide response to this problem have failed. However, there is an *ad hoc* international community of entomologists, the *CRB-G Action Group*, which meets annually (Table 1.1).

Table 1: Meetings of the CRB-G Action Group

|      |   |
|------|---|
| 2015 | Pacific Entomology Conference, Honolulu, HI, USA              |
| 2016 | International Congress of Entomology, Orlando, USA            |
| 2017 | Japanese Society for Insect Pathology, Tokyo, Japan           |
| 2018 | Society for Invertebrate Pathology, Gold Coast, Australia     |
| 2019 | XIX International Plant Protection Congress, Hyderabad, India |
| 2020 | (tentative): Pacific Plant Protection Organization, Guam      |

**Recent progress.** Recent work at the University of Guam, supported by grants from DOI-OIA and USDA-APHIS, has produced encouraging results (See Appendix B: *DOI-OIA Grant D17AP00107 Progress Report 4* for details):

- Laboratory tests indicate that OrNV from two sources can be considered as potential biocontrol agents CRB-G: OrNV isolate V23B maintained in insect tissue culture by AgResearch New Zealand and OrNV isolate UOGTW from bodies of CRB collected in Taiwan by the University of Guam CRB-G Biocontrol Project. Further laboratory testing of these virus samples is underway.
- PCR tests of recently collected CRB-G adults on Guam indicate presence of OrNV in this population. This virus could be from OrNV autodissemination earlier in the Guam CRB project or from fortuitous introduction.

## 1.2. Statement of Need

This proposal is, in part, a response to an urgent need expressed by Micronesian Island leaders in the 22nd Micronesian Islands Forum Communique (Appendix C):

In recognition of the urgent need for a Pacific-wide project to find an effective biological control agent for the CRB, the Leaders instructed RISC to seek financial support for such a project, to be conducted with partners at the University of Guam, the Secretariat of the Pacific Community, New Zealand, the USDA and others, as appropriate.

In addition to loss of coconut as ornamental plants and an emergency food supply, the uncontrolled CRB-G outbreak on Guam is a major environmental disaster rivaling that caused by the brown treesnake (BTS). BTS killed the birds in Guam's forests. CRB-G is now killing the trees. A 2002 US Forest Survey reported that the three most populous trees in Guam's forests were *Cycas micronesica*, *Cocos nucifera* and *Heterospathe elata* accounting for 16%, 12% and 11% of total trees with a stem diameter of five inches or greater (Moore 2018). These three species, 39% of the trees in Guam's forests, are being attacked by CRB-G<sup>1</sup>. If the Guam CRB-G outbreak is not brought under control, the island's forest health will continue to decline, accidental export of CRB-G to other islands in the American Pacific (in addition to Oahu and Rota) will be inevitable and cascading impacts from loss of forests will cause damage to other systems (erosion leading to reef fouling for example).

---

<sup>1</sup>Attack of *Cycas micronesica* by CRB-G adults is a recent discovery.

Despite the severity of the Guam CRB-G problem, the US federal government has provided relatively little support for response efforts. USDA-APHIS granted Plant Protection Act funding for CRB-G work in Hawaii and Guam for several years (Moore 2020). However, a grant proposal requesting \$331,4904 for Guam from FY2020 PPA funding was unexpectedly rejected. A proposal requesting support (\$3.5M) for a cooperative CRB biocontrol project to be performed by North Carolina State University, Colorado State University and the University of Guam has been submitted to DOD’s Strategic Environmental Research and Development Program (SERDP). If granted, this SERDP project will begin in the middle of 2021. Thus, **work towards mitigating the CRB-G problem on Guam during FY2020 is currently unfunded.**

The immediate challenge is to procure bridge funding to retain the Guam Biological Control Project’s insect pathologist, Dr. James Grasela, for an additional year so that we can begin propagation and field release of the two OrNV biological control candidates he has identified. Specialized skills of an insect pathologist are essential to implementing successful biological control of CRB-G on Guam and elsewhere.

### **1.3. Goals and Objectives**

Recent laboratory bioassays indicate that two recently tested OrNV isolates, V23B and UOGT are potential biocontrol candidates. There is also recent evidence indicating that there is already OrNV actively spreading within the Guam CRB-G population.

#### **1.3.1. Objective 1: Survey to Determine Background OrNV Incidence**

CRB adults collected from breeding sites and pheromone traps throughout Guam will be tested for presence of OrNV using PCR. Laboratory bioassays will be performed on OrNV isolated from these beetles to evaluate potential for biological control.

#### **1.3.2. Objective 2: Establish Sustainable CRB-G Biocontrol by Autodissemination of OrNV**

OrNV biocontrol candidates will be propagated *in vivo* using established methods (Huger 2005) and released into the Guam CRB-G population by autodissemination. Autodissemination involves infecting healthy CRB adults with OrNV. These infected beetles are then released at points dispersed throughout the island where they vector disease to conspecifics. A permit for field release of OrNV on Guam has already been obtained from USDA-APHIS. Field releases on CEMML on DOD land and by UOG on the rest of Guam. All released beetles will be marked by etching unique numbers on their elytra using a computer-controlled laser engraving system already in use for this application at UOG.

Beetles for *in vivo* propagation and autodissemination will be field-collected from breeding sites and pheromone traps because this is far more efficient than rearing beetles in the lab at the current time. Impact of virus releases will be monitored using pheromone traps and a novel roadside video analysis system (see Subsection 1.3.3). A subset of beetles captured in traps will be used to estimate the virus infection rate. Concurrent with virus releases, we will continue to screen OrNV isolates to find candidate biocontrol agents.

### 1.3.3. Objective 3: Establish Island-wide Monitoring Systems for CRB and Coconut Palm Health

The CRB-G outbreak on Guam is currently unmonitored on an island-wide basis. An island-wide pheromone trapping system, using about 1500 traps, was operated by the University of Guam from 2008 to 2014. This monitoring system was transferred to the Guam Department of Agriculture which abandoned the effort at the end of February, 2016. Currently, many coconut palms are being killed by CRB-G. But, in the absence of a monitoring system, we do not have an estimate of tree mortality or whether or not the damage is increasing or decreasing. Clearly, establishment of a monitoring system is necessary to evaluate success of the proposed biocontrol project, or any other mitigation efforts. We intend to re-establish island-wide trapping and to establish a sustainable roadside video survey which uses artificial intelligence to detect CRB damage in dash-cam videos.

**Pheromone Traps** We plan to installed 150 CRB pheromone monitoring traps. These will be baited with oryctalure and serviced semimonthly. These traps catch approximately equal numbers of males and females which remain alive in the traps for several weeks. Collected beetles will be used for autodissemination of virus and a subsample will be used for virus detection. Traps will be deployed at least 3 months prior to initiation of autodissemination.


A web database already exists for Guam CRB trap data and it is available for use by this project (URL: [mysql.guaminsects.net](http://mysql.guaminsects.net); database: **oryctes**; user: **readonlyguest**; password: **readonlypassword**; main tables: **trap** (2,265 records) and **trap\_visit** (89,114 records)).

**Roadside Surveys** Damage symptoms such as v-shaped cuts to fronds, bore holes, and dead standing coconut palm stems are readily observed during roadside surveys. Survey data will be collected on a smart-phone dash-cam app which georeferences each image. Initially, images of coconut palm damage by CRB-G will be detected, classified and tagged by a technician. When a large number of images have been tagged, these will be used to train an object detector. This work will result in a fully automated CRB damage detection and monitoring system which generates detection alerts and damage maps. This automated system will be useful as an early detection device for CRB. Roadside surveys on Guam will be performed bimonthly and the system will also be tested on Tinian, an island just north of Guam on which CRB has never been detected.



The envisioned system has already been successfully prototyped (See Supporting Technical Data [2](#)). A custom object detector for CRB damage has been trained using the TensorFlow implementation of the Faster R-CNN Deep Learning model (Moore, unpublished).

### 1.4. Timeline



## 1.5. Potential Benefits

- This project will directly benefit Guam. Without implementation of effective biological control, it is likely that 50% or more coconut palms will be killed by CRB-G. In addition to loss of coconut as ornamental plants and an emergency food supply, the uncontrolled CRB-G outbreak on Guam is a major environmental disaster with 39% the trees in Guam's forests are at risk of attack by CRB-G. If the outbreak is not brought under control, the island's forest health will continue to decline, accidental export of CRB-G to other islands in the American Pacific (in addition to Oahu and Rota) will be inevitable and cascading impacts from loss of forests will cause damage to other systems (erosion leading to reef fouling for example).
- This project will indirectly benefit all other islands in Micronesia and other areas of the Pacific. With very high populations of CRB-G in Guam risk of accidental introduction to other islands is extremely high. CRB-G has already been intercepted twice on Saipan, it is established on Rota and it has been detected on Aguiguan. If CRB-G infests smaller islands and atolls in the FSM and RMI where the coconut palm as the *tree of life*, islanders may have to migrate to larger population centers. If funded, the proposed project will be run under the *Open Science* concept. All data and analyses will be publicly shared on the internet in near-real time and samples of biocontrol candidates will be also be shared other Pacific islands battling CRB-G.
- It is expected that establishment of OrNV as an effective biological control agent will provide permanent self-sustaining population suppression of CRB-G. This is in contrast to temporary results from invasive species control programs which rely on population suppression using pesticide application, trapping or physical removal.
- Technology developed for automated roadside video surveys may be used for early detection of CRB on islands on which this pest has not yet been detected. Roadside videos may be of use for monitoring spread of invasive weeds such as Mexican creeper, *Antigonon leptopus*.

## 2. Budget

| Item               | Cost             |
|--------------------|------------------|
| Personnel          | \$146,200        |
| Benefits           | \$43,200         |
| Travel             | \$4,000          |
| Supplies           | \$15,290         |
| <b>SUBTOTAL</b>    | <b>\$208,690</b> |
| Administrative fee | \$31,304         |
| <b>TOTAL</b>       | <b>\$239,994</b> |

**Personnel** includes salary for an insect pathologist (Dr. James Grasela, 1 FTE, \$64,000), a laboratory technician (Mr. Chris Cayanan, 1 FTE, \$40,000) and a field technician (vacant, 1 FTE, \$40,000). Benefits for these 3 positions are calculated at 30% \* salary. Also includes PI's salary compensation calculated at 0.02 FTE \* \$110,000.

**Travel** includes airfare and other relocation expenses for Dr. Grasela who resides in Missouri.

**Supplies** includes laboratory and insect rearing supplies (\$5,340) and fuel and maintenance for the project's field vehicle (\$4,000), purchase of 100 pheromone traps (\$4,000), and purchase of 520 pheromone lures (\$1,950).

**Administrative fee** equal to 15% of the total grant award is charged by the Research Corporation of the University of Guam for services provided.

### 3. References

- Bedford, Geoffrey O. (2013). “Long-Term Reduction in Damage by Rhinoceros Beetle *Oryctes Rhinoceros* (L.) (Coleoptera: Scarabaeidae: Dynastinae) to Coconut Palms at *Oryctes* Nudivirus Release Sites on Viti Levu, Fiji”. In: *African J. Agricultural Research* 8.49, pp. 6422–6425. URL: [https://www.researchgate.net/publication/320921021\\_Long-term\\_reduction\\_in\\_damage\\_by\\_rhinoceros\\_beetle\\_Oryctes\\_rhinoceros\\_L\\_Coleoptera\\_Scarabaeidae\\_Dynastinae\\_to\\_coconut\\_palms\\_at\\_Oryctes\\_Nudivirus\\_release\\_sites\\_on\\_Viti\\_Levu\\_Fiji](https://www.researchgate.net/publication/320921021_Long-term_reduction_in_damage_by_rhinoceros_beetle_Oryctes_rhinoceros_L_Coleoptera_Scarabaeidae_Dynastinae_to_coconut_palms_at_Oryctes_Nudivirus_release_sites_on_Viti_Levu_Fiji).
- Catley, A (1969). “The Coconut Rhinoceros Beetle *Oryctes Rhinoceros* (L)”. In: *International Journal of Pest Management: Part A* 15.1, pp. 18–30. ISSN: 0434554690941. DOI: [10.1080/04345546909415075](https://doi.org/10.1080/04345546909415075).
- Gressitt, Linsley J (1953). *The Coconut Rhinoceros Beetle (Oryctes Rhinoceros) with Particular Reference to the Palau Islands*. Honolulu, pp. 1–83. URL: <https://books.google.com/books?id=0tcZAQAIAAJ>.
- Huger, Alois M. (2005). “The *Oryctes* Virus: Its Detection, Identification, and Implementation in Biological Control of the Coconut Palm Rhinoceros Beetle, *Oryctes Rhinoceros* (Coleoptera: Scarabaeidae)”. In: *Journal of Invertebrate Pathology* 89.1, pp. 78–84. DOI: [10.1016/j.jip.2005.02.010](https://doi.org/10.1016/j.jip.2005.02.010). URL: <http://www.guaminsects.net/CRB/docs/Huger2005.pdf>.
- Jackson, Trevor A (2009). “The Use of *Oryctes* Virus for Control of Rhinoceros Beetle in the Pacific Islands”. In: *Use of Microbes for Control and Eradication of Invasive Arthropods A.E. Hajak et al. [Eds.]* Ed. by Anne E. Hajek, T R Glare, and Maureen O’Callaghan. Vol. 6. Progress in Biological Control, pp. 133–140. URL: <http://guaminsects.net/CRB/docs/Jackson2009.pdf>.
- Jackson, Trevor A (2015). “Need for Emergency Response for a New Variant of Rhinoceros Beetle (Guam Biotype)”. In: *International Association for the Plant Protection Sciences Newsletter* XI. URL: <https://www.plantprotection.org/portals/0/documents/newsletters/2015/iapps%2011-2015.pdf>.
- Marshall, Sean D. G., Aubrey Moore, Maclean Vaqalo, Alasdair Noble, and Trevor A. Jackson (2017). “A New Haplotype of the Coconut Rhinoceros Beetle, *Oryctes Rhinoceros*, Has Escaped Biological Control by *Oryctes Rhinoceros* Nudivirus and Is Invading Pacific Islands”. In: *Journal of Invertebrate Pathology* 149, pp. 127–134. ISSN: 0022-2011. DOI: [10.1016/j.jip.2017.07.006](https://doi.org/10.1016/j.jip.2017.07.006). URL: <http://www.sciencedirect.com/science/article/pii/S0022201117300289> (visited on 08/26/2017).

- Marshall, Sean David Goldie, Maclean Vaqalo, Aubrey Moore, Roland Quitugua, and Trevor A Jackson (2015). “A New Invasive Biotype of the Coconut Rhinoceros Beetle (*Oryctes Rhinoceros*) Has Escaped from Biocontrol by *Oryctes Rhinoceros Nudi-virus*”. In: *International Congress on Invertebrate Pathology and Microbial Control and the 48th Annual Meeting of the Society for Invertebrate Pathology*. URL: <http://www.sipmeeting.org/van1/SIP2015-Full%20Program.pdf>.
- Moore, Aubrey (2018). “Failed Attempts to Establish IPM for Asian Cycad Scale and Coconut Rhinoceros Beetle on Guam”. URL: <https://zenodo.org/record/2545065/files/Moore-Vancouver-2018.pdf> (visited on 01/21/2019).
- Moore, Aubrey (2020). *USDA APHIS Farm Bill and Plant Protection Act Funding for Coconut Rhinoceros Beetle in Guam and Hawaii*. URL: <https://github.com/aubreymoore/wiki/wiki/USDA-APHIS-Farm-Bill-and-Plant-Protection-Act-Funding-for-Coconut-Rhinoceros-Beetle-in-Guam-and-Hawaii> (visited on 04/15/2020).
- Secretariat of the Pacific Community (2017). *Pest Alert 52: A New Biotype of Coconut Rhinoceros Beetle Discovered in the Pacific*. URL: [https://www.spc.int/sites/default/files/documents/blog/Pest-Alert-52-Coconut-Rhino-Beetle\\_finaledits.pdf](https://www.spc.int/sites/default/files/documents/blog/Pest-Alert-52-Coconut-Rhino-Beetle_finaledits.pdf) (visited on 04/18/2020).
- Vaqalo, Maclean, Sean Marshall, Trevor Jackson, and Aubrey Moore (2015). “Pest Alert 51: An Emerging Biotype of Coconut Rhinoceros Beetle Discovered in the Pacific”. In: pp. 2–2. URL: <http://westernipm.org/index.cfm/center-projects/signature-programs/invasive-species/coconut-rhinoceros-beetle/pest-alert-coconut-rhino-beetle-final-pdf/>.

## A. Supporting Technical Data

### A.1. Self-sustaining Biological Control of CRB in Fiji Using OrNV




Figure 1: Reduction in coconut palm damage following release of *Oryctes rhinoceros* nudivirus in Fiji. (Jackson 2009)

Reduction in palm damage recorded over 36 months after release of *Oryctes virus* on five sites in the Fiji Islands from 1970 to 1972. No virus was released in the Lautoka area where damage remained high 18 months after the start of the program but natural incidence of disease was recorded in the area after 36 months coinciding with a decline in visible damage. Population suppression of CRB by OrNV in Fiji was still in effect 35 years after virus introduction (Bedford 2013).

## A.2. Automated Monitoring of CRB Damage Using Roadside Video Surveys




Figure 2: Training an Object Detector to Locate Coconut Palms Damaged or Killed by Coconut Rhinoceros Beetle. <https://youtu.be/zzSorqcm9U>.

Result of a first attempt to train an object detector (Faster R-CNN) to locate coconut trees killed or damaged by coconut rhinoceros beetle in a video. Dead palms are in red boxes, damaged palms are in green boxes. Not perfect, but it does serve as a proof of concept.

## B. DOI-OIA Grant D17AP00107 Progress Report 4

Please see next page.



DOI-OIA Grant D17AP00107

Progress Report 4

Performance Period: March 1, 2019 - February 29, 2020

# Coconut Rhinoceros Beetle Biological Control

Aubrey Moore, University of Guam

March 23, 2020

<https://github.com/aubreymoore/doi-CRB-biocontrol-project/blob/master/report4/report4.pdf>

## Abstract

**CRB Biological Control.** The primary objective of this project is to find an isolate of *Oryctes rhinoceros* nudivirus (OrNV) which can be used as an effective biological control agent for the CRB-G biotype of coconut rhinoceros beetle (CRB).

Laboratory tests indicate that OrNV from two sources can be considered as potential biocontrol agents CRB-G: OrNV isolate V23B maintained in insect tissue culture by AgResearch New Zealand and OrNV in bodies of CRB collected in Taiwan for the current project. Further laboratory testing of these virus samples is underway.

PCR tests of recently collected CRB-G adults on Guam indicate presence of OrNV in this population. This virus could be from OrNV autodissemination early in the current project or from fortuitous introduction.

**CRB Damage Survey.** A secondary objective of this project is to develop a CRB damage monitoring system.

A digital image analysis system has been developed to detect and quantify V-shaped cuts to fronds and coconut palm mortality caused by CRB. The heart of this system is an object detector, trained by deep learning technology, which locates CRB damage symptoms on frames from georeferenced roadside video surveys. This object detector can be used to automate detection, quantification and to map changes in CRB damage over time and space and can also be used for early detection of CRB invasion.

A working prototype of the system has been built.

**Regional Collaboration.** Uncontrolled outbreaks of CRB-G is a major problem for Pacific islands. Outbreaks of this highly invasive biotype are damaging and killing palms in Guam, Rota, Hawaii, Palau, Papua New Guinea, and the Solomon Islands. Without effective control of these outbreaks, the problem will spread to other Pacific islands, resulting in a human tragedy when it reaches atolls where islanders still rely on coconut palm as the *tree of life*.

Project resources, time and effort were used to facilitate communication among an *ad hoc* collaboration of entomologists working on the CRB-G problem throughout the Pacific. Project staff participated in a symposium and planning meeting of the CRB-G Action Group at the XIX International Plant Protection Congress in November 2019.

# Contents

|          |  |           |
|----------|--|-----------|
| <b>1</b> | <b>Background</b>  | <b>4</b>  |
| <b>2</b> | <b>Staffing</b>  | <b>4</b>  |
| <b>3</b> | <b>Biological Control</b>  | <b>4</b>  |
| 3.1      | Laboratory Bioassays of OrNV Isolates . . . . .                                    | 4         |
| 3.2      | Virus Transmission Experiment . . . . .  | 5         |
| 3.3      | PCR Tests for OrNV Detection . . . . .   | 6         |
| 3.4      | CRB Rearing . . . . .  | 7         |
| 3.4.1    | Urgent Need for a Pathogen-free CRB Laboratory Colony . . . . .                    | 7         |
| 3.4.2    | Laboratory Information Management System . . . . .                                 | 7         |
| 3.4.3    | Acquisition of a Virus-Susceptible CRB Biotype for Comparative Bioassays . . . . . | 7         |
| <b>4</b> | <b>CRB Damage Survey</b>   | <b>8</b>  |
| <b>5</b> | <b>Regional Collaboration</b>  | <b>8</b>  |
| 5.1      | Participation in Scientific Meetings . . . . .                                     | 9         |
| 5.2      | Development of Online Resources . . . . .  | 9         |
|          | <b>References</b>  | <b>10</b> |

# 1 Background

The major goal of this project was to find an effective biological control agent for coconut rhinoceros beetle biotype G (CRB-G).

Prior to arrival of CRB-G on Guam during 2007, coconut rhinoceros beetle infestations of Pacific islands were readily dealt with by classical biological control using *Oryctes* nudiviruses (OrNV), a pathogen specific to rhinoceros beetles. Following a lack of response to release of OrNV on Guam, research showed that the Guam CRB population is a genetically distinct virus-resistant biotype which has become known as CRB-G (Marshall et al. 2017). This biotype is highly invasive and is causing massive damage to coconut and oil palms after recent invasion of Papua New Guinea and the Solomon Islands. CRB-G has also invaded Oahu and Rota. Eradication attempts have been launched on these two islands.

Additional goals for this project are to establish a CRB damage survey to evaluate efficacy of biocontrol and other tactics, and to maintain and facilitate collaboration with other Pacific island entomologists working to find solutions for CRB-G management.

## 2 Staffing

Staff for this project currently comprises of the PI, a post-doc and a technician.

- Funding from the Department of Interior, Office of Island Affairs was used to hire an insect pathologist for a 2 year term. Dr. James Grasela was recruited and started work at UOG on June 24, 2018.
- Funding from this USDA-APHIS project was used to hire Mr. Chris Cayanan as a technician. Mr. Cayanan was hired during December, 2019, as a replacement for Mr. Ian Iriarte who resigned to accept another job.

## 3 Biological Control

### 3.1 Laboratory Bioassays of OrNV Isolates

Four isolates of OrNV were evaluated as candidate biological control agents for CRB-G in a series of laboratory bioassays. Virus sample preparations came from Dr. Sean Marshall's lab at AgResearch New Zealand where they are maintained in insect cell culture.

**DUG42** Collected from Dumaguete, Negros Island, Philippines in 2017

**MALB** Collected from Malaysia, details not available.

**PNG** Collected from Rabaul, Papua New Guinea in 1988

**V23B** Collected from southern Luzon, Philippines in 1980

During laboratory bioassays, we dosed CRB-G adults with samples of the OrNV isolates, and observed mortality and changes in mass for one month. Each beetle was kept in isolation and individual records were stored in a laboratory information system developed for this application (Section 3.4.2).

Bioassay results, displayed in Table 1, indicate that one of the isolates, V23B, is pathogenic for CRB-G when doses are applied by placing droplets of virus suspension on mouthparts of adult beetles.

Table 1: *Oryctes rhinoceros* nudivirus (OrNV) bioassay results summary.

| OrNV isolate | bioassay   | method <sup>1</sup> | beetles | replicates | virus mortality ( $p$ ) <sup>2</sup> | inactivated virus mortality ( $p$ ) <sup>3</sup> |
|--------------|--|---------------------|---------|------------|--------------------------------------|--|
| DUG42        | DUG42 (Moore and J. J. Grasela 2019a)              | injection           | 30      | 2          | 40% (0.65)                           | 40% (0.65)                                       |
| MALB         | MALB (Moore and J. J. Grasela 2019b)               | injection           | 30      | 2          | 50% (0.37)                           | 0% (1.00)  |
|              | MALBperOS (Moore and J. J. Grasela 2019c)          | per os              | 13      | 1          | -60% (1.00)                          | 20% (1.00)                                       |
| PNG          | PNG (Moore and J. J. Grasela 2019d)                | injection           | 81      | 4          | 90% (0.00)                           | 5% (1.00)  |
|              | PNGperOS (Moore and J. J. Grasela 2019e)           | per os              | 21      | 1          | 0% (1.00)                            | 0% (1.00)  |
| V23B         | V23B (Moore and J. J. Grasela 2019f)               | injection           | 66      | 4          | 88% (0.00)                           | 0% (1.00)  |
|              | V23BperOS (Moore and J. J. Grasela 2019h)          | per os              | 32      | 2          | 80% (0.07)                           | 20% (0.69)                                       |
|              | V23-large_bioassay (Moore and J. J. Grasela 2019g) | per os              | 53      | 1          | 42% (0.00)                           | -  |
|              | V23_perOSIN (Moore and J. J. Grasela 2019i)        | per os              | 16      | 1          | 60% (0.06)                           | -  |

<sup>1</sup> Adult beetles were dosed either by direct injection of virus suspension into the haemocoel or by applying a droplet containing virus to mouthparts.

<sup>2</sup> Percent mortality in beetles treated with virus, adjusted for untreated control mortality; number in parentheses is the  $p$ -value resulting from a Fisher's exact test of significant difference between mortality of treated and untreated beetles.

<sup>3</sup> Percent mortality in beetles treated with heat inactivated virus, adjusted for untreated control mortality; number in parentheses is the  $p$ -value resulting from a Fisher's exact test of significant difference between mortality of treated and untreated beetles.

In a separate experiment, macerated guts from OrNV infected beetles from field collections of CRB in Taiwan were fed to CRB-G field-collected on Guam. PCR results indicated that the Taiwanese OrNV propagated in the Guam beetles (See Subsection 3.3).

We now have two isolates of OrNV which can be considered as candidate biocontrol agents for further testing: V23B and Taiwan.

## 3.2 Virus Transmission Experiment

This experiment was performed to determine if OrNV isolate V23B can be transmitted from a dosed CRB adult to an undosed CRB adult. If a virus is not contagious in the lab, it will probably have no potential as a biocontrol agent.

Unfortunately, the experiment failed due to very high mortality in the experimental control group and results are inconclusive. For details, see J. Grasela and Moore 2020. This experiment needs to be repeated using pathogen-free, laboratory-reared adults (See Section 3.4).

### 3.3 PCR Tests for OrNV Detection

Previously, our laboratory relied on outside collaboration for molecular testing to determine presence of OrNV in CRB tissues. We recently acquired access to equipment and supplies which allow us to use PCR (polymerase chain reaction) techniques for OrNV detection. We tested five different primer pairs for detection of OrNV in reference samples from AgResearch New Zealand and were successful with all five. Identity of DNA fragments was confirmed as OrNV using a commercial DNA sequencing service.

Details of our PCR technique and initial series of tests can be found in J. J. Grasela and Moore [2020a](#), J. J. Grasela and Moore [2020b](#) and J. J. Grasela and Moore [2020c](#).

Results from PCR tests indicate:

- OrNV is present in the Guam adult CRB population: 18% (10 of 47) gut samples dissected from field collected CRB tested positive for OrNV.
- OrNV is present in the adult Taiwanese CRB population: 7% (5 of 67) gut samples dissected from field collected CRB tested positive for OrNV.
- OrNV from Taiwanese adult CRB propagates in Guam adult CRB: 12% (5 of 41) Guam CRB adults dosed with Taiwanese gut preparations tested positive.

## 3.4 CRB Rearing

Experimental beetles were field-collected on Guam as prepupae, pupae or adults from breeding sites or as adults from pheromone traps. Each beetle was given a serial number and was reared individually in a Mason jar stored in one of three environmental chambers set for 30 degrees Celsius, 80% relative humidity and 12 hour photoperiod. Each adult beetle was fed weekly with a slice of banana. Detailed information on the CRB rearing facility can be found in a document prepared in support of a USDA-APHIS permit to allow importation of live CRB for laboratory bioassays (Moore 2019a).

### 3.4.1 Urgent Need for a Pathogen-free CRB Laboratory Colony

To date, CRB used in bioassays and other laboratory experiments were field collected on Guam, rather than from reared in a laboratory colony. This was done because a very high population density of rhino beetles on Guam made field collection far more efficient in terms of time and resources than laboratory rearing. However, field collected beetles are often infected with pathogens, especially the fungus, *Anisopliae majus*, which was introduced on Guam as a classical biological control agent. Presence of these pathogens in experimental animals resulted in unpredictable and often high mortality rates in experimental control groups, leading to inconclusive results such as experienced with the recent virus transmission experiment (3.2).

Recent discovery of OrNV within the wild Guam rhino beetle population (3.3) now makes it mandatory to establish a sterile CRB-G laboratory colony to supply experimental animals.

### 3.4.2 Laboratory Information Management System

We developed an online database which we use as a laboratory information management system for maintaining individual records for beetles in our rearing program (Moore 2019b). This system was developed using the [web2py python web framework](#) and it is available on the web at <http://aubreymoore.pythonanywhere.com/rearing3>.

### 3.4.3 Acquisition of a Virus-Susceptible CRB Biotype for Comparative Bioassays

Since discovery of the CRB-G biotype on Guam (Marshall et al. 2017), we have been operating under the hypothesis that this biotype is significantly more tolerant (resistant) to pathogenic effects of OrNV isolates previously used as effective biocontrol agents for CRB invading Pacific Islands. It has also been hypothesized that CRB-G has different behavioral characteristics, such as a significantly reduced attraction to oryctalure. However, comparative laboratory bioassays have not been performed to test these hypotheses.

We applied for and have been granted a USDA-APHIS import permit for live CRB which will allow us to establish a laboratory colony of CRB from presumed non-virus-resistant populations (See (Moore 2019a) and (USDA-APHIS 2019)). We have designed custom shipping containers to facilitate secure transport of live CRB (Moore and Quitugua 2017).

We plan to import CRB from American Samoa with assistance from our collaborator, Dr. Mark Schmaedick, American Samoa Community College. Plans for project staff to visit American Samoa in December 2019 failed because of travel restrictions prompted by a measles outbreak and in March 2020 because of travel restrictions prompted by the corona virus pandemic.

## 4 CRB Damage Survey

The objective of this component of the project was to develop an automated system to evaluate CRB damage by image analysis of roadside video surveys. We completed a *proof of concept* trial in which deep learning algorithms were used to train an object detector which locates and counts dead and CRB-damaged coconut palms in video streams. Visual results are presented in a YouTube post (Moore 2019d) and technical details are available in an Open Science Framework Project (Moore 2019c).

## 5 Regional Collaboration

An informal collaboration, the *CRB-G Action Group*, has been formed among Pacific-based entomologists working on the CRB-G problem. Participants from Guam, Hawaii, Palau, Papua New Guinea, Solomon Islands, Fiji, Malaysia, Japan and New Zealand have met several times and future meetings are planned (Table 2). This is an *ad hoc* group which has been organized by Dr. Trevor Jackson and Sean Marshall of AgResearch New Zealand. AgResearch is recognized as a global center for expertise on biological control of CRB. AgResearch scientists have worked on CRB in the south Pacific for several decades and they maintain a library of OrNV isolates in cell culture. The New Zealand government has recently committed several million dollars to aid response to CRB-G in the south Pacific islands. Although individual institutions working to find a solution to the CRB-G problem on American-affiliated islands in the northern Pacific have secured funding from multiple, short-term grants, attempts to secure funding to support a sustainable well-coordinated regional project have been unsuccessful.



Table 2: Meetings of the CRB-G Action Group

---

|  |
|--|
| 2015 Pacific Entomology Conference, Honolulu, HI, USA              |
| 2016 International Congress of Entomology, Orlando, USA            |
| 2017 Japanese Society for Insect Pathology, Tokyo, Japan           |
| 2018 Society for Invertebrate Pathology, Gold Coast, Australia     |
| 2019 XIX International Plant Protection Congress, Hyderabad, India |
| 2020 (tentative): Pacific Plant Protection Organization, Guam      |

---

## 5.1 Participation in Scientific Meetings

Moore and Grasele participated at the XIX International Plant Protection Congress in a symposium entitled *The challenge of coconut rhinoceros beetle, *Oryctes rhinoceros*, to palm production and prospects for control in a changing world*. Moore made an oral presentation at this meeting (Moore, J. J. Grasele, and Marshall 2019). They also participated in a CRB-G Action Group meeting with colleagues from throughout the Pacific and Asia.

## 5.2 Development of Online Resources

Project resources were used to build and maintain the following:

- CRB Bibliography <https://github.com/aubreymoore/CRB-Bibliography>
- Interactive CRB Invasion History Map <https://aubreymoore.github.io/crbdist/mymap.html>
- CRB Wiki Site <https://guaminsects.net/CRBG>
- CRB-G Facebook Site <https://www.facebook.com/groups/crbg07>

## References

- Grasela, James J. and Aubrey Moore (2020a). *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 1*. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-1.pdf>.
- Grasela, James J. and Aubrey Moore (2020b). *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 2*. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-2.pdf>.
- Grasela, James J. and Aubrey Moore (2020c). *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 3*. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-3.pdf>.
- Grasela, James and Aubrey Moore (2020). *Guam CRB Biocontrol Project Technical Report: OrNV Transmission Experiment*. URL: <https://github.com/aubreymoore/OrNV-Transmission/blob/master/ornv-transmission.pdf>.
- Marshall, Sean D. G., Aubrey Moore, Maclean Vaqalo, Alasdair Noble, and Trevor A. Jackson (2017). “A New Haplotype of the Coconut Rhinoceros Beetle, *Oryctes Rhinoceros*, Has Escaped Biological Control by *Oryctes Rhinoceros* Nudivirus and Is Invading Pacific Islands”. In: *Journal of Invertebrate Pathology* 149, pp. 127–134. ISSN: 0022-2011. DOI: 10.1016/j.jip.2017.07.006. URL: <http://www.sciencedirect.com/science/article/pii/S0022201117300289> (visited on 08/26/2017).
- Moore, Aubrey (2019a). *Additional Documentation in Support of a Permit Application for Importation of Live Coconut Rhinoceros Beetles to Guam*. URL: <https://github.com/aubreymoore/CRB-Import-Permit/blob/master/CRB-import-permit-request-additional.pdf>.
- Moore, Aubrey (2019b). *Coconut Rhinoceros Beetle Rearing Database*. URL: <http://aubreymoore.pythonanywhere.com/rearing>.
- Moore, Aubrey (2019c). *Open Science Framework: Digital Image Analysis for Detection of CRB Damage in Videos*. URL: [https://osf.io/czr8t/?view\\_only=cdec7cc47b514b31b33115113193a](https://osf.io/czr8t/?view_only=cdec7cc47b514b31b33115113193a).
- Moore, Aubrey, director (2019d). *Training an Object Detector to Locate Coconut Palms Damaged or Killed by Coconut Rhinoceros Beetle*. URL: <https://www.youtube.com/watch?v=zzSorqcmt9U> (visited on 10/09/2019).
- Moore, Aubrey and James J. Grasela (2019a). *Bioassay Report: DUG42*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-DUG42.pdf>.
- Moore, Aubrey and James J. Grasela (2019b). *Bioassay Report: MALB*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-MALB.pdf>.

- Moore, Aubrey and James J. Grasela (2019c). *Bioassay Report: MALBperOS*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-MALBperOS.pdf>.
- Moore, Aubrey and James J. Grasela (2019d). *Bioassay Report: PNG*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-PNG.pdf>.
- Moore, Aubrey and James J. Grasela (2019e). *Bioassay Report: PNGperOS*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-PNGperOS.pdf>.
- Moore, Aubrey and James J. Grasela (2019f). *Bioassay Report: V23B*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23B.pdf>.
- Moore, Aubrey and James J. Grasela (2019g). *Bioassay Report: V23B Large Bioassay*. University of Guam. URL: [https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23B%5C\\_large%5C\\_bioassay.pdf](https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23B%5C_large%5C_bioassay.pdf).
- Moore, Aubrey and James J. Grasela (2019h). *Bioassay Report: V23BperOS*. University of Guam. URL: <https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23BperOS.pdf>.
- Moore, Aubrey and James J. Grasela (2019i). *Bioassay Report: V23BperOSIN*. University of Guam. URL: [https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23B%5C\\_perOSIN.pdf](https://github.com/aubreymoore/rearing3/raw/master/bioassay-V23B%5C_perOSIN.pdf).
- Moore, Aubrey, James J. Grasela, and Sean D. G. Marshall (2019). *Presentation: Status of a Major Outbreak of Coconut Rhinoceros Beetle, Oryctes Rhinoceros Biotype G, on Guam and Attempts at Establishing Biological Control*. XIX International Plant Protection Congress, Hyderabad, India. Nov. 11, 2019. URL: [https://github.com/aubreymoore/IAPPS-2019-Presentation/raw/master/Moore\\_IAPPS-2019.odp](https://github.com/aubreymoore/IAPPS-2019-Presentation/raw/master/Moore_IAPPS-2019.odp).
- Moore, Aubrey and Roland Quitugua (2017). *Container for Secure Shipment of Live Coconut Rhinoceros Beetle Adults*. Guam Coconut Rhinoceros Beetle Project Technical Report CRB-2017-05-27. University of Guam Cooperative Extension Service, p. 3. URL: [https://github.com/aubreymoore/CRB-Import-Permit/raw/master/CRB\\_Shipping\\_Container.pdf](https://github.com/aubreymoore/CRB-Import-Permit/raw/master/CRB_Shipping_Container.pdf).
- USDA-APHIS (2019). *CRB Import Permit*. URL: <https://github.com/aubreymoore/CRB-Import-Permit/blob/master/2019-CRB-Import-Permit.pdf>.

## C. Extracts from the 22nd Micronesia Islands Forum Joint Communique



**EDDIE BAZA CALVO**  
GOVERNOR OF GUAM  
Chairman/Host



**PETER CHRISTIAN**  
PRESIDENT OF THE  
FEDERATED STATES OF MICRONESIA  
Member



**TOMMY REMENGESAU, JR.**  
PRESIDENT OF THE REPUBLIC OF PALAU  
Member




**HILDA HEINE**  
PRESIDENT OF THE REPUBLIC  
OF THE MARSHALL ISLANDS  
Member



**RALPH D.L.G. TORRES**  
GOVERNOR OF  
THE COMMONWEALTH OF THE  
NORTHERN MARIANA ISLANDS  
Member



**JOHNSON ELIMO**  
GOVERNOR OF CHUUK  
Member



**LYNDON H. JACKSON**  
GOVERNOR OF KOSRAE  
Member



**MARCELO PETERSON**  
GOVERNOR OF POHNPEI  
Member



**JAMES YANGETMAI**  
LT. GOVERNOR OF YAP  
Member

# 22nd Micronesian Islands Forum

MAY 01, 2017 ~ MAY 03, 2017 | GUAM



The Leaders called for the support of Micronesia Challenge Plus (MC+), which would fully integrate global and regional challenges, including Climate Change, Disaster Risk Management, and Invasive Species toward enhanced and holistic conservation measures and to move the 2020 commitments beyond 2020 and into the future. This will keep the MC in line with the revised and expanded UN Sustainable Development Goals.

## 2. Regional Invasive Species Council (RISC)

The Regional Invasive Species Council (RISC) reported on progress since the 21<sup>st</sup> MIF. RISC identified and discussed two major issues needing supporting action from the Leaders, and an additional two developments in the United States which needed to be brought to the attention of the Leaders. RISC also met with the MC SC to discuss ways that RISC can support the MC.

The first and most urgent issue is the need for regional coordination of invasive species activities in Micronesia, including implementation of the Regional Biosecurity Plan (RBP). RISC recommended the creation of a Regional Invasive Species Coordination Office to be staffed by a Regional Coordinator and to be housed in the Micronesia Center for a Sustainable Future (MCSF). The Regional Coordinator would be responsible to coordinate implementation of the RBP across the region, as well as to coordinate other activities and initiatives.

The second very urgent issue is the growing threat of the coconut rhinoceros beetle (CRB) to the region. CRB is devastating coconut trees in Guam, Palau, and other Pacific Islands. There is a high risk of their spread to other islands in Micronesia, and there is currently no effective control. A regional project is therefore urgently needed to develop an effective biological control for CRB.

RISC met with the MC SC to discuss ways that RISC can support the MC in their efforts to minimize the impacts of invasive species in terrestrial and marine conservation sites throughout the region as an integral component of effective conservation. Together, they identified the critical need to work together to establish baselines of invasive species in conservation sites. RISC will continue to assist the MC as they work to effectively conserve natural resources by protecting conservation sites from invasive species.

# 22nd Micronesian Islands Forum


MAY 01, 2017 ~ MAY 03, 2017 | GUAM

  
**EDDIE BAZA CALVO**  
GOVERNOR OF GUAM  
Chairman/Host

  
**PETER CHRISTIAN**  
PRESIDENT OF THE  
FEDERATED STATES OF MICRONESIA  
Member

  
**TOMMY REMENGESAU, JR.**  
PRESIDENT OF THE REPUBLIC OF PALAU  
Member

  
**HILDA HEINE**  
PRESIDENT OF THE REPUBLIC  
OF THE MARSHALL ISLANDS  
Member

  
**RALPH D.L.G. TORRES**  
GOVERNOR OF  
THE COMMONWEALTH OF THE  
NORTHERN MARIANA ISLANDS  
Member

  
**JOHNSON ELIMO**  
GOVERNOR OF CHUUK  
Member

  
**LYNDON H. JACKSON**  
GOVERNOR OF KOSRAE  
Member

  
**MARCELO PETERSON**  
GOVERNOR OF POHNPEI  
Member

  
**JAMES YANGETMAI**  
LT. GOVERNOR OF YAP  
Member

## ➤ Regional Coordination for Implementation of the Regional Biosecurity Plan (RBP)

The Leaders supported the establishment of a Regional Coordinator position in the to be established Regional Invasive Species Coordination Office housed in the MCSF. The Leaders committed themselves, through the MCSF, to provide initial funding support for the first two years to help ensure the effective implementation of the RBP.

## ➤ US Presidential Executive Order on Promoting Agriculture and Rural Prosperity in America

The Leaders signed a joint letter to the Secretary of the US Department of Agriculture (USDA) welcoming the Executive Order signed by President Donald Trump on April 25, 2017, listing accomplishments of the Micronesian Region in the battle against invasive species and outlining needs for further progress. The Leaders look forward to working closely with the USDA and the Interagency Task Force to promote rural agricultural opportunity, food security and rural prosperity in the islands of Micronesia.

## ➤ Regional biocontrol project for Coconut Rhinoceros Beetle (CRB)

In recognition of the urgent need for a Pacific-wide project to find an effective biological control agent for the CRB, the Leaders instructed RISC to seek financial support for such a project, to be conducted with partners at the University of Guam, the Secretariat of the Pacific Community, New Zealand, the USDA and others, as appropriate.

## ➤ RISC/MC SC cooperation – capacity building for baseline surveys

The Leaders recognized the value of the collaboration between RISC and the MC SC and the progress they have made toward integrating invasive species prevention and management into effective conservation. The Leaders support the plan for RISC to work with the MC Measures Working Group to develop a capacity-building project to enable all jurisdictions to conduct baseline surveys for invasive plants, including development of a regional MC invasive species database for use in planning and decision-making. The Leaders expect that this will



  
**EDDIE BAZA CALVO**  
GOVERNOR OF GUAM  
Chairman/Host

  
**PETER CHRISTIAN**  
PRESIDENT OF THE  
FEDERATED STATES OF MICRONESIA  
Member

  
**TOMMY REMENGESAU, JR.**  
PRESIDENT OF THE REPUBLIC OF PALAU  
Member

  
**HILDA HEINE**  
PRESIDENT OF THE REPUBLIC  
OF THE MARSHALL ISLANDS  
Member

  
**RALPH D.L.G. TORRES**  
GOVERNOR OF  
THE COMMONWEALTH OF THE  
NORTHERN MARIANA ISLANDS  
Member

  
**JOHNSON ELIMO**  
GOVERNOR OF CHUUK  
Member

  
**LYNDON H. JACKSON**  
GOVERNOR OF KOSRAE  
Member


  
**MARCELO PETERSON**  
GOVERNOR OF POHNPEI  
Member

  
**JAMES YANGETMAI**  
LT. GOVERNOR OF YAP  
Member


# 22nd Micronesian Islands Forum


MAY 01, 2017 ~ MAY 03, 2017 | GUAM


MAY 02 2017


  
**EDDIE BAZA CALVO**  
Governor  
U. S. Territory of Guam


  
**TOMMY E. REMENGESAU, JR.**  
President  
Republic of Palau


  
**PETER M. CHRISTIAN**  
President  
Federated States of Micronesia


  
**HILDA C. HEINE, Ed. D**  
President  
By Minister John M. Silk  
Republic of the Marshall Islands

  
**JOHNSON ELIMO**  
Governor  
Chuuk State, FSM

  
**RALPH D.L.G. TORRES**  
Governor  
Commonwealth of the  
Northern Mariana Islands

  
**MARCELO K. PETERSON**  
Governor  
Pohnpei State, FSM

  
**LYNDON H. JACKSON**  
Governor  
Kosrae State, FSM

  
**JAMES YANGETMAI**  
Lt. Governor  
Yap State, FSM

## **D. Letters of Support**

### **D.1. Guam Invasive Species Council**

Please see next page.





**Lourdes A. Leon Guerrero**  
Governor  
**Joshua F. Tenorio**  
Lt. Governor

# Department of Agriculture Dipåttamenton Agrikottura

163 Dairy Road, Mangilao, Guam 96913



**Chelsa Muña-Brecht**  
Director  
**Adrian Cruz**  
Deputy Director

April 20, 2020

Douglas W. Domenech  
Assistant Secretary Insular and International Affairs  
U.S. Department of the Interior  
1849 C Street, N.W.  
Mail-Stop 3119  
Washington, D.C. 20240

Håfa adai Assistant Secretary Domenech,

As the Chairperson of the Guam Invasive Species Council, I am pleased to submit a letter of support for Dr. Moore's grant proposal entitled *Establishment of Self-sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia*.

We have identified the coconut rhinoceros beetle (CRB) as a priority invasive species that threatens Guam's forest and agricultural ecosystems. CRB has killed large numbers of coconuts and other indigenous palms on Guam. It will continue to do so and continue to spread to other islands in Micronesia and beyond if we cannot get the Guam outbreak under control.

Dr. Moore at the University of Guam is offering a promising, sustainable solution to our problem. Unfortunately, USDA-APHIS is not supporting CRB work on Guam during FY2020. Dr. Moore has invested several years into viral biocontrol development and stoppage at this critical point may cause an irreparable setback. I urge your office to fund his proposal.

Si Yu'os Ma'åse,

**Chelsa Muña-Brecht**  
Director

## **D.2. Regional Invasive Species Council for the Western Pacific**

Please see next page.

Mr. Leonard Sigrah, Chair  
Regional Invasive Species Council  
P.O. Box 51 Lelu, Kosrae, FM 96944



Douglas W. Domenech  
Assistant Secretary Insular and International Affairs  
U.S. Department of the Interior  
1849 C Street, N.W.  
Mail-Stop 3119  
Washington, D.C. 20240

Dear Assistant Secretary Domenech,

As Chair of the Regional Invasive Species Council for the Western Pacific, a standing committee of the Micronesia Islands Forum's nine jurisdictions, I am pleased to submit a letter of support for Dr. Moore's grant proposal entitled *Establishment of Self-sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia*.

In virtually every travel destination across Micronesia, *Cocos nucifera* (coconut tree) is synonymous with the island's image and community identity. The tree is ecologically resilient, culturally valued, and a daily part of the economy and vitality of island life. Unfortunately, this plant has recently undergone dramatic mortality in the region.

The effect this new wave of Coconut Rhinoceros Beetle infestations has created is felt on every island jurisdiction today, from the Marshall Islands to Palau. To illustrate the extreme, Fuana (a small, uninhabited atoll island in Palau's southwest districts) is now devoid of *C. nucifera*, while anecdotal and documented evidence throughout the region reveal extensive damage and mortality to both farmed and natural tree populations.

We have reviewed Dr. Moore's proposal, applaud his commitment, and fully support the University and our Partner's request for assistance. At this juncture, we believe that his research will progress towards an ecologically-sound method of long-term coastal forest recovery and resilience in the region, and urge your office to fund his proposal.

Respectfully yours,

  
Leonard Sigrah, Invasive Species Coordinator

## E. Cover Letter

Please see next page.



Aubrey Moore, PhD  
College of Agriculture and Life Sciences  
University of Guam  
Mangilao, Guam 96923

April 23, 2020

Douglas W. Domenech  
Assistant Secretary, Insular and International Affairs  
U.S. Department of the Interior  
1849 C Street, N.W.  
Mail-Stop 3119  
Washington, D.C. 20240

Dear Assistant Secretary Domenech,

I am pleased to submit a grant proposal entitled *Establishment of Self-sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia* for consideration under the Coral Reef and Natural Resources Initiative (CFDA 15.875).

The Guam Invasive Species Council has identified the coconut rhinoceros beetle as a priority invasive species which threatens Guam's forestry and agriculture. CRB has already killed large numbers of coconuts and other palms on Guam. It will continue to do so, and spread to other islands in Micronesia and beyond if we cannot get the Guam outbreak under control.

Dr. James Grasela, an insect pathologist hired under previous DOI-OIA funding has identified two isolates of *Oryctes nudivirus* (OrNV) which can be used as biological control agents for CRB-G, the rhino beetle biotype which is causing damage in Guam, Hawaii, CNMI, Palau, Papua New Guinea, and the Solomon Islands.

I am asking DOI-OIA support for a one year project to implement biological control for CRB-G on Guam by propagating and releasing the biological control agents identified by Dr. Grasela. A USDA-APHIS release permit has already been approved.

The consensus among Pacific-based entomologists is that biological control is the best, and possibly the only way, to stop palms from being killed on an island-wide basis and to reduce the risk of accidental transport to other islands in Micronesia and beyond. Based on previous CRB biocontrol projects, we expect population suppression of CRB-G by OrNV to be self-sustaining.

Yours sincerely,

Aubrey Moore, PhD