

Establishment of Self-sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia

Aubrey Moore PhD

University of Guam College of Natural and Applied Sciences

Federal ID D20AP00060

Reporting period 2020-06-01 through 2020-12-31

Contents

1 Objective 1: Survey to Determine Background OrNV Incidence	3
2 Objective 2: Establish Sustainable CRB-G Biocontrol by Autodissemination of OrNV	4
3 Objective 3: Establish Island-wide Monitoring Systems for CRB and Coconut Palm Health	5
3.1 Objective 3a: Pheromone Traps	5
3.2 Objective 3b: Roadside Video Surveys	6
3.2.1 Guam Roadside Video Survey 1	6
3.2.2 Guam Roadside Video Survey 2	11
3.2.3 Rota Roadside Video Survey 1	12
4 References	13

Notes for the Reader

- Each objective, as stated in the funded grant proposal [1], is presented in a frame at the start of each section.

Todo list

1 Objective 1: Survey to Determine Background OrNV Incidence

CRB adults collected from breeding sites and pheromone traps throughout Guam will be tested for presence of OrNV using PCR. Laboratory bioassays will be performed on OrNV isolated from these beetles to evaluate potential for biological control.

Our lab currently uses CRB-G adults collected from pheromone traps as test animals in bioassays evaluating OrNV isolates as biocontrol agents under the assumption that the Guam beetle population contains only the CRB-G biotype and is free from OrNV infection. In 2019 we gained the capacity to perform PCR in our lab and began testing these assumptions. PCR results indicated that untreated, field-collected beetles, used as experimental controls in our bioassays, were all CRB-G, but 18% of these tested positive for OrNV [2, 3, 4].

Based on these results, the PI decided to suspend bioassays until we had conclusive evidence of OrNV infection in the Guam CRB-G population. This decision was made for two reasons:

1. If field-collected CRB insects where already infected with OrNV, these could not be used as test insects in laboratory bioassays designed to evaluate virulence of OrNV isolates.
2. If an OrNV strain was already spreading within the Guam CRB population, it would be prudent to evaluate this strain as an effective biological control agent.

An experimental plan [5] was developed and executed. One hundred beetles were collected from each of two pheromone trapping sites (Leo Palace Resort in Southern Guam and the UOG Ag. Expt. Stn. in northern Guam). Gut samples were obtained from these beetles and tested using PCR in our lab and also in Sean Marshall's lab at AgResearch New Zealand.

In PCR results from both labs all beetles tested positive for CRB-G biotype and negative for OrNV infection [6].

We conclude that previous OrNV positive tests were the result of lab contamination (not false positives).

Current evidence suggests that all CRB on Guam belong to the CRB-G biotype and there is no OrNV within this population. In other words, the background OrNV incidence is zero.

2 Objective 2: Establish Sustainable CRB-G Biocontrol by Autodissemination of OrNV

OrNV biocontrol candidates will be propagated *in vivo* using established methods [7] and released into the Guam CRB-G population by autodissemination. Autodissemination involves infecting healthy CRB adults with OrNV. These infected beetles are then released at points dispersed throughout the island where they vector disease to conspecifics. A permit for field release of OrNV on Guam has already been obtained from USDA-APHIS. All released beetles will be marked by etching unique numbers on their elytra using a computer-controlled laser engraving system already in use for this application at UOG.

Beetles for *in vivo* propagation of OrNV and autodissemination will be field-collected from breeding sites and pheromone traps because this is far more efficient than rearing beetles in the lab at the current time. Impact of virus releases will be monitored using pheromone traps and a novel roadside video analysis system (see Objective 3). A subset of beetles captured in traps will be used to estimate the virus infection rate. Concurrent with virus releases, we will continue to screen OrNV isolates to find candidate biocontrol agents.

We have not made progress on this objective for two reasons:

1. High mortality of untreated insects in experimental control groups and discovery of OrNV in some of these insects, presumably from laboratory contamination, have thrown results of previous bioassays into question. None of the several OrNV isolates we have tested in laboratory bioassays jumps out as a good candidate for effective biological control of CRB-G.
2. We require permission from the US Environmental Protection Agency (USEPA) prior to release of OrNV as a biological control agent. Data demonstrating efficacy of OrNV as a biological control agent is required by USEPA.

3 Objective 3: Establish Island-wide Monitoring Systems for CRB and Coconut Palm Health

The CRB-G outbreak on Guam is currently unmonitored on an island-wide basis. An island-wide pheromone trapping system, using about 1500 traps, was operated by the University of Guam from 2008 to 2014. This monitoring system was transferred to the Guam Department of Agriculture which abandoned the effort at the end of February, 2016. Currently, many coconut palms are being killed by CRB-G. But, in the absence of a monitoring system, we do not have an estimate of tree mortality or whether or not the damage is increasing or decreasing. Clearly, establishment of a monitoring system is necessary to evaluate success of the proposed biocontrol project, or any other mitigation efforts. We intend to re-establish island-wide trapping and to establish a sustainable roadside video survey which uses artificial intelligence to detect CRB damage in dash-cam videos.

3.1 Objective 3a: Pheromone Traps

We plan to installed 150 CRB pheromone monitoring traps. These will be baited with oryctalure and serviced semimonthly. These traps catch approximately equal numbers of males and females which remain alive in the traps for several weeks. Collected beetles will be used for autodissemination of virus and a subsample will be used for virus detection. Traps will be deployed at least 3 months prior to initiation of autodissemination.

A web database already exists for Guam CRB trap data and it is available for use by this project (URL: mysql.guaminsects.net; database: **oryctes**; user: **readonlyguest**; password: **readonlypassword**; main tables: **trap** (2,265 records) and **trap_visit** (89,114 records)).

We have not deployed new pheromone traps. But we continue to maintain trap lines in central Guam, at the Leo Palace Resort, and in northern Guam, at the University of Guam Agricultural Station in Yigo.

3.2 Objective 3b: Roadside Video Surveys

Damage symptoms such as v-shaped cuts to fronds, bore holes, and dead standing coconut palm stems are readily observed during roadside surveys. Survey data will be collected on a smart-phone dash-cam app which georeferences each image. Initially, images of coconut palm damage by CRB-G will be detected, classified and tagged by a technician. When a large number of images have been tagged, these will be used to train an object detector. This work will result in a fully automated CRB damage detection and monitoring system which generates detection alerts and damage maps. This automated system will be useful as an early detection device for CRB. Roadside surveys on Guam will be performed bimonthly and the system will also be tested on Tinian, an island just north of Guam on which CRB has never been detected.

The envisioned system has already been successfully prototyped. A custom object detector for CRB damage has been trained using the TensorFlow implementation of the Faster R-CNN Deep Learning model (Moore, unpublished).

Bimonthly automated roadside video surveys for CRB damage are now operational on Guam and the system has been tested on Rota. Videos recorded with a smart phone attached to a vehicle are analyzed using custom-designed artificial intelligence software which recognizes coconut palms and measures CRB damage. The software was developed in collaboration with OnePanel Inc. A nontechnical description of the survey method is given in the next section.

A presentation on this new CRB survey methodology was made at the December 9 2020 meeting of the CRB-G Action Group. Videos of all presentations at this meeting are available online [9].

3.2.1 Guam Roadside Video Survey 1

The following four images were extracted from a draft of the University of Guam's Western Pacific Tropical Research Center impact report for 2020. They provide a nontechnical overview of the new automated roadside video survey for CRB damage and results from the first Guam survey in October, 2020.




Figure 1: Feature article in the University of Guam's Western Pacific Tropical Research Center impact report for 2020.

Everyone living on Guam has seen damage to coconut palms caused by coconut rhinoceros beetles (CRB). CRB has been on Guam since 2007, however, until recently, the number of palms being damaged and killed on Guam was unknown. Standardized surveys of CRB damage are needed to monitor changes over time and space, especially in response to control activities and for early detection of CRB in new geographic areas.

UOG entomologist Aubrey Moore has developed a highly automated method for routine island-wide monitoring of CRB damage using a cell phone and artificial intelligence (AI).

Methods for monitoring CRB damage have been developed. But these rely on direct observation or image analysis by human experts and are too time-consuming and expensive for routine monitoring over large areas.

Dr. Trevor Jackson, an entomologist working for AgResearch New Zealand, has developed a survey method based on a five-level scale for classifying CRB damage to individual coconut palms. Jackson's method is being used extensively on CRB-infested islands in the South Pacific. Moore decided to develop an island-wide roadside CRB damage survey for Guam based on an automated version of Jackson's method.



A smart phone is attached to a vehicle using a magnetic mount. As the car travels, the phone records videos that are analyzed by open-source software.


In the automated survey, a smart phone mounted on a car or truck records continuous videos while the vehicle is driven along all major roads on Guam. The smart phone uses a couple of free apps: OpenCamera records videos and GPSLogger records GPS coordinates.

Recent technical breakthroughs in AI have made it much easier to train computers to recognize objects in digital images. Moore collaborated with OnePanel Inc., an AI tech



Above: medium to severe CRB damage detected in the Royal Palms area of Dededo.

Below: each dot on the map represents a video frame in which one or more coconut palms was detected. The image at the top is a frame extract indicated by the camera icon.



Continue reading →

8

Figure 2: [Continued] Feature article in the University of Guam's Western Pacific Tropical Research Center impact report for 2020.




Figure 3: [Continued] Feature article in the University of Guam's Western Pacific Tropical Research Center impact report for 2020. This interactive web map is publicly available at: <https://aubreymoore.github.io/new-crb-damage-map>

WPTRC 2019 IMPACT REPORT

company, to develop and train a couple of object detectors using a technique called "deep learning".

The first object detector finds all images of coconut palms in the survey videos and classifies each one using Jackson's damage scale. The second object detector locates and counts v-shaped cuts in the fronds of each coconut palm. Data extracted from the videos by the object detectors are saved in a SpatialLite database.

To visualize survey results, Moore uses Quantum GIS, to make a publicly available, interactive web map. There are links on the web map to download the survey database and QGIS map project for more detailed analysis.

The first operational island-wide survey on Guam, completed during October 2020, indicated that about 19% of Guam's coconut palms show CRB damage symptoms. The Guam surveys will be conducted bimonthly. An island-wide roadside video survey is also being done on Rota for early detection of CRB damage.

There is interest in use of roadside video surveys for CRB damage elsewhere in the Pacific and Moore plans to evaluate drone imagery for use on islands without extensive roads. The Guam roadside video survey was designed to be adaptable by using only free



open-source software (FOSS) components. Custom-written software for the project as well videos, databases, and GIS projects from surveys will also be made available for download from public repositories hosted on GitHub.

It is interesting to note that this is not the first time that Moore has dabbled with AI. Thirty years ago he trained an artificial neural network to identify free-flying mosquitoes.

Funded by the US Department of the Interior-Office of Insular Affairs, US Forest Service, USDA-APHIS

Thanks to UOG entomology technician Christian Cayanan for doing the surveys.

Further reading:

Moore, A. 2018. The Guam Coconut Rhinoceros Beetle Problem: Past, Present and Future. Zenodo. doi.org/10.5281/zenodo.1185371}.

Moore, A. 1991. Artificial neural network trained to identify mosquitoes in flight. Journal of Insect Behavior 4, 391–396.

Aubrey Moore
(671) 735-2068
aubreymoore@triton.uog.edu

10

Figure 4: [Continued] Feature article in the University of Guam's Western Pacific Tropical Research Center impact report for 2020.

3.2.2 Guam Roadside Video Survey 2

The proportion of coconut palms damaged by CRB increased significantly from 19.2% in October 2020 to 21.5% in December 2020 ($p < 0.001$; Fisher's exact test).




Figure 5: Screenshot of an interactive web map of results from a roadside video survey of CRB damage on Guam in December 2020 <https://aubreymoore.github.io/Guam-CRB-damage-map-2020-12/webmap/v1/>.

3.2.3 Rota Roadside Video Survey 1

Rota was invaded by CRB in 2017 and eradication efforts by Rota Department of Land and Natural Resources have successfully kept the population at a very low level. In October 2020, a smart phone and associated equipment was sent to the so that they could do an initial roadside video survey in support of their CRB control efforts. In addition to the equipment, a survey setup guide and [10] and a setup video [11] were prepared and sent.

The survey was performed by Mark Mangolana, DLNR and the phone containing videos from the survey was returned to the University of Guam. Videos were analyzed using the workflow developed for the Guam surveys. The resulting web map contained many false positives for CRB damage, but there is one hit which shows a classic v-shaped cut probably caused by CRB. For convenience, data for this hit (images, date, location) were documented as an iNaturalist observation (Figure 6). If this v-shaped cut was caused by CRB, there will be a bore hole. Rota-DLNR are following up to see if this is the case.




Figure 6: Screenshot of an iNaturalist observation documenting probable coconut rhinoceros beetle damage detected during a roadside video survey performed by Rota DLNR. <https://www.inaturalist.org/observations/69534809>.

4 References

- [1] Aubrey Moore. *Grant Proposal: DOI-OIA Coral Reef and Natural Resources Initiative FY2020: Establishment of Self-Sustaining Biological Control of Coconut Rhinoceros Beetle Biotype G in Micronesia*. Apr. 23, 2020. URL: https://github.com/aubreymoore/2020-DOI-CRB-Biocontrol/blob/master/doi_proposal.pdf (visited on 05/22/2020).
- [2] James J. Grasela and Aubrey Moore. *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 1.* 2020. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-1.pdf>.
- [3] James J. Grasela and Aubrey Moore. *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 2.* 2020. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-2.pdf>.
- [4] James J. Grasela and Aubrey Moore. *Technical Report: Polymerase Chain Reaction (PCR) Analysis of the Coconut Rhinoceros Beetle (CRB), Oryctes Rhinoceros. Part 3.* 2020. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/blob/master/pcrdata/PCR-Summary-Part-3.pdf>.
- [5] Aubrey Moore, James J. Grasela, and Sean David Goldie Marshall. *Experimental Plan: Determining Presence of OrNV in the Guam CRB-G Population*. 2020. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/raw/master/docs/Guam%20rNV%20Experimental%20Plan.pdf>.
- [6] James J. Grasela and Aubrey Moore. *Investigation to Determine the Presence of OrNV in the Guam CRB Population*. 2020. URL: <https://github.com/aubreymoore/FY19-PPA-Report-1/raw/master/docs/Determining%20the%20Presence%20of%20rNV%20in%20the%20Guam%20CRB.pdf>.
- [7] Alois M. Huger. “The Oryctes Virus: Its Detection, Identification, and Implementation in Biological Control of the Coconut Palm Rhinoceros Beetle, Oryctes Rhinoceros (Coleoptera: Scarabaeidae)”. In: *Journal of Invertebrate Pathology* 89.1 (May 2005), pp. 78–84. DOI: [10.1016/j.jip.2005.02.010](https://doi.org/10.1016/j.jip.2005.02.010). URL: <http://www.guaminsects.net/CRB/docs/Huger2005.pdf>.
- [8] Aubrey Moore. “Ideas for Re-Establishment of CRB Lab Colonies at UoG”. Dec. 30, 2020. URL: <https://docs.google.com/document/d/1wo0cDjz06LNsvDX9rlHY3bz1R09R704PzLRhrv-8pw/edit?usp=sharing>.
- [9] Aubrey Moore. *Video Recording of the CRBG Action Group Meeting: December 9, 2020 [Conducted via Zoom]*. Dec. 9, 2020. URL: <https://aubreymoore.github.io/CRBG-action-group-webinar-20201209/>.

- [10] Aubrey Moore. *Set Up for Automated Roadside Video Surveys for Detecting and Monitoring Coconut Rhinoceros Beetle Damage on Rota*. Oct. 26, 2020. URL: <https://api.zotero.org/groups/511387/items/3X83KK58/file/view>.
- [11] Aubrey Moore. *YouTube Video: Mounting a Smart Phone on a Vehicle for Roadside Video Surveys of Coconut Rhinoceros Beetle Damage*. 2020. URL: <https://www.youtube.com/watch?v=Pa-UslJV6BI>.