

Coconut Rhinoceros Beetle, *Oryctes rhinoceros*

Coleoptera: Scarabaeidae

A Major Threat to Hawaii's Coconut and Palm Trees




Aubrey Moore UG

Arnold H. Hara
University of Hawaii at Manoa
College of Tropical Agriculture & Human Resources
875 Komohana St. Hilo, Hawaii
E-mail: arnold@hawaii.edu

Coconut rhinoceros beetle (CRB)

Native range: Southeastern Asia (M. Schmaedick 2005 Am. Samoa)

- *CRB was introduced throughout the Pacific primarily due to increased sea traffic during World War II.
- *Most recently, CRB was discovered in Guam in September 2007.
- *Primary damage is caused by adults boring from the petioles of fronds into the crown, cutting through developing leaves, and feeding on the exuded sap.
- *The beetle breeds in dead, standing coconut palms killed by pest/disease/ lightning, and in decaying organic materials, such as compost and sawdust heaps. (Bedford, 1980).
- *Eggs hatch in 8-12 days; larvae feed on decaying coconut/palm debris for 82-207 days.
- *Prepupal and pupal stage is 25-35 days; adult remains in the pupal cell for 17-22 days.
- *Adults live for 4-9 months; each female lays 50-100 eggs.



Coconut rhinoceros beetle (CRB) - Damage Symptoms

**Primary damage is caused by adults boring from the petioles of fronds into the crown, cutting through developing, unopened fronds, and feeding on the exuded sap.*

V-shaped cut on open fronds.

Similar to mechanical pruning damage to unopened fronds



Active adult boring hole in petiole causing "wet look"



Active adult boring hole thru petiole



Coconut fibers resulting from adults' boring



**DAMAGE TO EMERGING FROND ON
FOXTAIL PALM**



**EXPOSED COCONUT FIBER
CAUSED BY BORING**



Area attacked by adult beetle



Site of old adult boring hole



Adult boring hole thru frond petiole



Site of old boring holes in trunk



Symptoms of CRB attack: V-cuts with associated lateral boring thru fronds



Semi-circular damage caused by boring through folded new terminal growth



Major breeding site among coconut trimming debris - Asan, Guam



Major Breeding Site with all CRB stages - Asan, Guam



Asan Beach, Guam



Dead coconut
tree with CRB



Adults and
grubs found
in rotting
coconut trunk
terminal





Most Efficient CRB Trap in Guam

- 50 gal steel barrel filled with rotting coconut debris, live CRB, grubs and aggregation pheromone.
- Ultraviolet LED light over trap.
- Chicken wire allows CRB to enter but not fly out (with spread wings).



UV LED light

(Moore & Quitugua, pers. com. 09/13)



UV LED light



Pheromone
w/CRB



hooded vane trap



UV LED light plus
pheromone
increase trap
efficiency
(Moore & Quitugua,
pers. com. 09/13)

Trap with chicken wire
instead of vane



Traps with CRB pheromone



Rotting coconut stumps
placed beneath trap with
CRB pheromone increases
trap efficiency

Guam 09/13

