

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/330832017>

ADVANCES IN ENTOMOLOGICAL LASER RADAR

Conference Paper · October 2018

CITATIONS

0

READS

21

2 authors:



Mikkel Brydegaard

Lund University

90 PUBLICATIONS 659 CITATIONS

[SEE PROFILE](#)



Samuel Jansson

Lund University

27 PUBLICATIONS 72 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Inelastic hyperspectral lidar [View project](#)



Entomological lidar [View project](#)

ADVANCES IN ENTOMOLOGICAL LASER RADAR

Mikkel Brydegaard^{1,2,3,4*}, Samuel Jansson^{1,2}

¹Lund Laser Centre, Dept. Physics, Lund University, Sölvegatan 14, Lund, Sweden

²Center for Animal Movement Research, Dept. Biology, Lund University, Sölvegatan 37, Lund, Sweden

³Norsk Elektro Optikk AS, Prost Stabells vei 22, Skedsmokorset, Norway

⁴FaunaPhotonics ApS, Ole Maaløes Vej 3, København N, Denmark

* mikkel.brydegaard@fysik.lth.se

Keywords: Laser Radar, Biophotonics, Remote Sensing, Pest Monitoring, Entomological Ecology

Abstract

During last decade we have developed and applied optical remote sensing instrumentation for *in situ* remote surveillance and quantification of the aerofauna. The sparse structure of aerofauna makes optical focusing challenging, but we solved this issue through applying the century old Scheimpflug condition. With this approach we have managed to reduce size, cost and complexity of atmospheric lidars and accomplished an effective tool for ecological entomology capable of counting thousands of insects per hour. Because of the high sensitivity and resolution in time and space, we can retrieve target modulation signatures in the kHz range for target classification purposes. As opposed to the cm waves in entomological radar, we rely on near infrared light around one micron. This allow superior beam quality, negligible ground clutter and applications close over ground or within vegetation structure. Near infrared light can assess both molecular and microstructural properties of the target through differential absorption and depolarization. Here we give the background of entomological lidar, summarize our recent progress and put it in context with contemporary work. We outline applications, ongoing activities and state of the art. We discuss future prospects and challenges.

1. Background

Following WWII development of radar, radar ornithology^{1,2} and later radar entomology^{3,4} emerged in the last century. This century entomological lidar emerged⁵⁻⁸ with time-of-flight lidars at kHz rates developed in Montana. Our group in Lund started out using a slower atmospheric lidar at hand⁹⁻¹² in particular explorer feasibility of fluorescence lidar. We later also developed kHz methods^{13,14}. As opposed to insect sampling with sweep nets and traps (see e.g.¹⁵) remote sensing has the advantages of being non-intrusive, non-perturbing and with known biases and detection ranges and limits. Further, remotely sensed insects are to some extent classified automatically^{16,17}, whereas analysis of caught insects is tedious but provide more details^{18,19}. Most radar entomology studies have targeted high migrating insects by vertical profiling. The size ratio between a large aircraft and insects is roughly the same at the wavelength ratio between microwaves in radar and infrared light and as opposed to microwaves, laser and optics offer great beam qualities allowing investigations of entomological ecology close over ground or within forest without ground clutter. Lidar and photonics methods also offer assessment of molecular and microstructural information through absorption, fluorescence

and depolarization processes. Applications of entomological lidar includes monitoring abundance and dispersal of agricultural and forestry pest, disease vectors for humans and livestock and not least assessing biodiversity of pollinators. We will here give an overview of the recent progress in the field.

2. Instrumentation

2.1 Passive lidar

As in passive radar²⁰, insects can be sensed optically using existing radiation sources such as the sun for diurnal species²¹, the moon²² or thermal infrared emission²³ for nocturnal species. Sunlight offer a range of wavelengths from UV to IR and some 1kWm^{-2} of power. We started out developing the remote dark-field method²¹ which gives lidar like signals across the spectrum without need of laser development. This method is based on directing a telescope at a remote black cavity and collect the light scattered from airborne insects. We demonstrated powder tagging and dark-field modulation spectroscopy²⁴ and heading assessment by quadrant detection¹³ (corresponding to beam wobbling in radar). The method can also be employed at several spectral bands at kHz rates for determining body and wing melanisation^{25,26}. For some time we attempted passive ranging²⁷ but only recently provided a solution using quadrant detection²⁸.

2.2 Time-of-Flight entomological lidar

Equivalently to radar, conventional lidar is based on transmission of ns long laser pulses and recording the backscattered intensity with fast detectors and RF electronics as a function of round-trip delay. Applied systems ranges from slow conventional multiband systems^{10,29,30} to fast single band systems^{8,31}.

2.3 Scheimpflug lidar

Similarly to bistatic radar, atmospheric Scheimpflug lidars utilise transmitted and receiver separated by a small baseline of roughly a meter^{13,32-35}. The Scheimpflug principle allow infinite focal depth with large apertures¹⁴, therefore weak signals from insect scattering can be retrieved at kHz rates. The average emitted power can greatly exceed pulsed lidars despite the much smaller and inexpensive lasers. The systems can efficiently collect high number of observations^{32,34} (See Fig.1). Scheimpflug lidars can be expanded to retrieve several spectral bands^{31,36,37} or polarisation modes^{33,35}. For example for assessing melanin (See Fig.2). Scheimpflug lidar

can also retrieve fluorescence³⁸ with applications for insect powder tagging^{10,24,39}, however violet light may affect the insects⁴⁰.




Fig.1. Fraction of entomological data from a Scheimpflug lidar. The lidar was profiling bees over a clover field, the hive was situated 180 m from the lidar. Bees oscillates around 200 Hz indicated by red color code (data from FaunaPhotonics).




Fig.2. Dual band remote in insect observation using a short-wave infrared Scheimpflug lidar, the melanisation of body and wings can be derived (adapted from³¹).




Fig.3. a) Polarimetric image of a dry ♂ *Anopheles Coluzzii* mosquito, 808 nm, 162° backscatter. b) Aspect dependent backscatter cross sections with- and without wings (oscillatory part). c) Corresponding extinction cross section, note the different scale (adapted from⁴¹).

2.4 Reference measurements

Effort was put into quantifying and understanding optical backscatter- and extinction-cross sections for insects for comparison and feasibility of remote target classification. In

analogy with early radar referencing⁴² the Montana group recorded polarimetric backscatter of pinned bees⁷. Our group recorded oscillatory cross sections of free flying insects^{43,44} including spectral and polarisation modes feasible for lidar. We also applied hyper spectral imaging⁴⁵, goniometry⁴⁶ and polarisation imaging⁴¹ of pinned insects (See Fig.3). Referencing can also be done by releasing known insects into the lidar beam in field⁴⁷ or sampling insect along the beam with a trap⁴⁸.

3. Methods, purpose and outcome.

3.1 Organism sizing

If the lidar beam-FoV overlap function can be determined, e.g., by a molecular Rayleigh echo and assumptions of homogeneous atmosphere³², then backscatter intensities can be converted into Lidar Cross Sections in mm². Whereas RCS related to liquid water mass, Lidar Cross Section was defined as the projected area had the target been a 100% Lambertian reflector such as a Teflon beat. In practice insect reflectance in near infrared^{34,41}, 808 nm, is in the order of 20%, the reflectance in short wave infrared⁴⁵, 1320 nm, is in the order of 80% whereas specular reflectance⁴⁶ could reach 5000%. Despite this, organism sizing allow estimation of the aerofauna biomass spectrum³² and insect and predating vertebrate can display a bimodal biomass spectrum³⁴. Biomass spectra are popular in marine science,

3.2 Modulation spectroscopy for target classification

The wing beat cycle causes backscatter and extinction cross section to oscillate with a fundamental tone and a harmonic spectrum. This spectrum has proven valuable for target classification^{16,49,50} or even biodiversity assessment⁵¹. The methods are directly applicable for entomological lidar applications but the work is mainly based on so called E-traps somewhat simpler than lidar and often operating in extinction mode. The drawback is that the counts per day are much lower than lidar^{32,34,50}. Much of this work involve advanced machine learning but even the fundamental tone can differentiate species, e.g.⁵². Some challenges are that the tone is temperature dependent⁵³, that harmonic spectrum is aspect dependent^{28,43,44} and that identification of the fundamental tone is difficult⁵⁴. Modulation spectroscopy can be expanded by additional spectral and polarisation bands (See Fig.4).




Fig.4. Frequency components of co- and de-polarised backscatter from a *Drosophila Melanogaster* at 808 nm. The high harmonics are from glossy wings and therefore co-polarised (for instrument details see⁴⁴).

3.3 Powder tagging

Insect can be tagged optically using fluorescent powders

from the printing and advertising community, and several colours can be used simultaneously. This is referred to as mark-without-recapture studies and yield the greatest certainty about target identity. Social insects such as bees can be auto-tagged by installing a tray in the hive exit. Powdered individuals can be detected remotely either by sunlight²⁴ or violet lasers^{10,29,38}. Applications of tagging include landscape dispersal studies, population size estimates and verifications of classification, e.g., by modulation spectroscopy. Corresponding tagging in radar entomology is accomplished by harmonic radar by attaching electrical diodes to the insects⁵⁵.

3.4 Heading assessment

Heading assessment is important for understanding dispersal and migration properties, not least from epidemiological aspects of economic and medical entomology. It can be accomplished in radar by polarisation anisotropy⁵⁶, beam wobbling or tracking⁵⁷. In optics heading can be assessed by quadrant detectors^{13,26,58}, heading could also be inferred by transit times³³ or even odd harmonics⁴³ however specular contributions from wings deviates from this model²⁸.

3.5 Tracking individuals

Reported entomological lidar studies typically report single timings and positions of many individuals, whereas tracking approaches report many timings and positions of a single individual. Tracking radar ornithology is widespread⁵⁷ whereas report from radar entomology are limited⁵⁵. In optics efficient tracking can be accomplished by static passive systems^{23,59}, although active lock-in servo systems have been reported⁶⁰.

4. Doppler assessment of movements

Movements of animals and their bodypart may yield small shift in backscatter photon energy, this have been demonstrated with weather radars⁶¹ and micro Doppler radar technology⁶². The reports on applications of widespread Doppler lidars are limited⁶³.

4. Conclusion, challenges and outlook

We have revised the development of entomological lidar instrumentation and compared it to radar. We have demonstrated aspects of modulation signatures, dual band and polarimetric insect scattering. We have discussed various types of studies and outcome. A current challenge is that there are no table values for scatter cross sections of the millions of insect species. Entomological lidar scientists should be encouraged to calibrate and standardize cross sections for comparison and not report arbitrary units. Future work could benefit from improved reference measurement, and linking lidar recording both to modulation spectroscopy⁵⁰ and museum 3D insect libraries⁶⁴.

5. Acknowledgements

We thank Mariam Andersson, Peggy Atkinson, Klas Rydhmer, Alfred Strand, Birte Boelt for contributed material. Entomological lidar has been supported by amound others: Swedish Research Council, Norsk Elektro Optikk, FaunaPhotonics, Lund University, Stellenbosch University,

South China Normal University, Physiographic Society in Lund, Linnaeus grant to Lund Laser Centre and Center for Animal Movement Research.

6. References

- [1] S. A. Gauthreaux and C. G. Belser, "RADAR ornithology and biological conservation," *The Auk*, vol. **120**, pp. 266-277, 2003/04/01 2003.
- [2] E. Eastwood, *Radar ornithology*: Methuen, 1967.
- [3] J. Riley, "Remote sensing in entomology," *Annual review of entomology*, vol. **34**, pp. 247-271, 1989.
- [4] V. A. Drake and D. R. Reynolds, *Radar entomology : observing insect flight and migration* Wallingford, Oxfordshire ; Boston, MA: CABI, 2012.
- [5] D. S. Hoffman, A. R. Nehrir, *et al.*, "Range-resolved optical detection of honeybees by use of wing-beat modulation of scattered light for locating land mines," *Appl. Opt.*, vol. **46**, pp. 3009-3012, 2007.
- [6] K. S. Repasky, J. A. Shaw, *et al.*, "Optical detection of honeybees by use of wing-beat modulation of scattered laser light for locating explosives and land mines," *Appl. Opt.*, vol. **45**, pp. 1839-1843, 2006.
- [7] J. A. Shaw, N. L. Seldomridge, *et al.*, "Polarization lidar measurements of honey bees in flight for locating land mines," *Opt. Expr.*, vol. **13**, pp. 5853-5863, 2005.
- [8] M. J. Tauc, K. M. Fristrup, and J. A. Shaw, "Development of a wing-beat-modulation scanning lidar system for insect studies," in *Lidar Remote Sensing for Environmental Monitoring*, San Diego, USA, 2017.
- [9] M. Brydegaard, P. Lundin, *et al.*, "Feasibility study: fluorescence lidar for remote bird classification," *Appl. Opt.*, vol. **49**, pp. 1-15, 2010.
- [10] Z. Guan, M. Brydegaard, *et al.*, "Insect monitoring with fluorescence lidar techniques: Field experiments," *Appl. Opt.*, vol. **49**, pp. 1-11, 2010.
- [11] M. Brydegaard, Z. Guan, *et al.*, "Insect monitoring with fluorescence lidar techniques: feasibility study," *Appl. Opt.*, vol. **48**, pp. 5668-5677, 2009.
- [12] P. Lundin, P. Samuelson, *et al.*, "Remote nocturnal bird classification by spectroscopy in extended wavelength ranges," *Appl. Opt.*, vol. **50**, 2011.
- [13] M. Brydegaard, A. Gebru, and S. Svanberg, "Super resolution laser radar with blinking atmospheric particles – application to interacting flying insects.," *Prog. Electromagn. Res.*, vol. **147**, pp. 141-151, 2014.
- [14] M. Brydegaard, E. Malmqvist, *et al.*, "The Scheimpflug Lidar Method," *SPIE Lidar Remote Sensing for Environmental Monitoring*, vol. **10406**, 2017.
- [15] J. B. Silver, *Mosquito ecology: field sampling methods*: Springer Science & Business Media, 2007.
- [16] A. Moore and R. H. Miller, "Automated identification of optically sensed aphid (*Homoptera: Aphidae*) wingbeat waveforms," *Ann. Entomol. Soc. Am.*, vol. **95**, pp. 1-8, 2002.
- [17] E. Malmqvist, S. Jansson, *et al.*, "Effective parameterization of laser radar observations of atmospheric fauna," *Selected Topics in Quantum Electronics, IEEE Journal of*, vol. **22**, pp. 327-334, 2016.
- [18] M. A. Hoy, *Insect molecular genetics: an introduction to principles and applications*: Academic Press, 2003.
- [19] IAEA, *Manual for the Use of Stable Isotopes in Entomology*: IAEA, VIENNA, 2009.
- [20] H. D. Griffiths and C. J. Baker, *An Introduction to Passive Radar*: Artech House, 2017.
- [21] A. Runemark, M. Wellenreuther, *et al.*, "Rare events in remote dark field spectroscopy: an ecological case study of insects," *IEEE JSTQE*, vol. **18**, pp. 1573 - 1582, 2012.
- [22] M. Brydegaard and S. Svanberg, "Photonic monitoring of atmospheric and aquatic fauna," *Laser & Photonics Reviews*, 2018.

- [23] D. H. Theriault, N. W. Fuller, *et al.*, "A protocol and calibration method for accurate multi-camera field videography," *The Journal of experimental biology*, vol. **217**, pp. 1843-1848, 2014.
- [24] M. Brydegaard, A. Merdasa, *et al.*, "Realistic instrumentation platform for active and passive optical remote sensing," *Applied Spectroscopy*, vol. **70**, pp. 372-385, 2016.
- [25] A. Gebru, S. Jansson, *et al.*, "Remote species and sex determination of disease vector mosquitoes using multispectral polarimetric techniques," 2017.
- [26] A. Gebru, M. Brydegaard, *et al.*, "Probing insect backscatter cross-section and melanization using kHz optical remote detection system," *SPIE, Journal of Applied Remote Sensing*, vol. **16611P**, 2016.
- [27] P. Lundin, M. Brydegaard, *et al.*, "Passive unmanned sky spectroscopy for remote bird classification," *Conference on Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII/18th Internat. Symposium on Remote* vol. **8174**, pp. 81740J-81740J-11, Sep 19-21, 2011 2011.
- [28] S. Jansson and M. Brydegaard, "Passive Entomological Lidar for Stand-Off Modulation Spectroscopy and Heading Assessment," *Animal Telemetry*, 2018.
- [29] L. Mei, Z. G. Guan, *et al.*, "Agricultural pest monitoring using fluorescence lidar techniques," *Appl. Phys. B*, vol. **106**, pp. 733-740, 2011.
- [30] S. Jansson, M. Brydegaard, *et al.*, "Exploitation of an atmospheric lidar network node in single-shot mode for the classification of aerofauna," *JARS*, 2017.
- [31] M. Andersson, "Short-Wave Infrared Lidar on Atmospheric Aerosols and Insects," Msc., Physics, Lund University, 2017.
- [32] M. Brydegaard, A. Gebru, *et al.*, "Daily evolution of the insect biomass spectrum in an agricultural landscape accessed with lidar," in *ILRC 27*, New York, USA, 2015.
- [33] S. Zhu, E. Malmqvist, *et al.*, "Insect abundance over Chinese rice fields in relation to environmental parameters, studied with a polarization-sensitive CW near-IR lidar system," *Applied Physics B*, vol. **123**, 2017.
- [34] E. Malmqvist, S. Jansson, *et al.*, "The bat-bird-bug battle: daily flight activity of insects and their predators over a rice field revealed by high resolution Scheimpflug Lidar " *Royal Society Open Science*, 2018.
- [35] L. Mei and P. Guan, "Development of an atmospheric polarization Scheimpflug lidar system based on a time-division multiplexing scheme," *Optics Letters*, vol. **42**, pp. 3562-3565, 2017/09/15 2017.
- [36] M. Brydegaard, J. Larsson, *et al.*, "Short-wave infrared atmospheric Scheimpflug lidar," *EDP Sciences*, vol. **176**, 2018.
- [37] G. Zhao, E. Malmqvist, *et al.*, "Dual-band continuous-wave lidar system employed for particle classification," *IEEE sensors*, 2017.
- [38] G. Zhao, M. Ljungholm, *et al.*, "Inelastic Hyperspectral Lidar for Profiling Aquatic Ecosystems," *Laser & Photonics Reviews*, vol. **10**, pp. 807-813, 2016.
- [39] J. R. Hagler and C. G. Jackson, "Methods for marking insects: current techniques and future prospects," *Annual review of entomology*, vol. **46**, pp. 511-543, 2001.
- [40] K. Rydhmer and A. Strand, "Applied hyperspectral LIDAR for monitoring fauna dispersal in aquatic environments," 2016.
- [41] S. Jansson, P. Atkinson, and M. Brydegaard, "First Polarimetric Investigation of Malaria Mosquitos as Lidar Targets," *IEEE JSTQE Biophotonics*, 2018.
- [42] J. Riley, "Angular and temporal variations in the radar cross-sections of insects," in *Proceedings of the Institution of Electrical Engineers*, 1973, pp. 1229-1232.
- [43] M. Brydegaard, "Towards quantitative optical cross sections in entomological laser radar – Potential of temporal and spherical parameterizations for identifying atmospheric fauna," *PLoS One*, vol. **10**, p. e0135231, 2015.
- [44] A. Gebru, S. Jansson, *et al.*, "Multiband modulation spectroscopy for determination of sex and species of mosquitoes in flight," *J. Biophotonics*, vol. **J. Biophotonics**, 2018.
- [45] M. Brydegaard, "Advantages of shortwave infrared LIDAR entomology," in *Imaging and Applied Optics 2014*, Seattle, Washington, 2014, p. LW2D.6.
- [46] M. Brydegaard, S. Jansson, *et al.*, "Can the narrow red bands of dragonflies be used to perceive wing interference patterns?," *Ecology and Evolution*, 2018.
- [47] S. Török, "Kilohertz electro-optics for remote sensing of insect dispersal," Master Msc thesis, Department of Physics, Lund University, 2013.
- [48] C. Kirkeby, M. Wellenreuther, and M. Brydegaard, "Observations of movement dynamics of flying insects using high resolution lidar," *Scientific Reports*, vol. **6**, p. 29083, 07/04/online 2016.
- [49] D. F. Silva, V. M. Souza, *et al.*, "Exploring low cost laser sensors to identify flying insect species," *Journal of Intelligent & Robotic Systems*, vol. **80**, pp. 313-330, 2015.
- [50] I. Potamitis, I. Rigakis, and K. Fysarakis, "Insect Biometrics: Optoacoustic Signal Processing and Its Applications to Remote Monitoring of McPhail Type Traps," *PLoS ONE*, vol. **10**, p. e0140474, 2015.
- [51] T. Ganchev, *Computational Bioacoustics: Biodiversity Monitoring and Assessment*: Walter de Gruyter GmbH & Co KG, 2017.
- [52] A. P. Genoud, R. Basistyy, *et al.*, "Optical remote sensing for monitoring flying mosquitoes, gender identification and discussion on species identification," *Applied Physics B*, vol. **124**, p. 46, February 17 2018.
- [53] S. M. Villarreal, O. Winokur, and L. Harrington, "The Impact of Temperature and Body Size on Fundamental Flight Tone Variation in the Mosquito Vector *Aedes aegypti* (Diptera: Culicidae): Implications for Acoustic Lures," *Journal of Medical Entomology*, vol. **54**, pp. 1116-1121, 2017.
- [54] A. Andersson, "Unbiasing entomological kHz Scheimpflug LIDAR data," Msc., Dept. Physics, Lund University, 2018.
- [55] O. Ovaskainen, A. D. Smith, *et al.*, "Tracking butterfly movements with harmonic radar reveals an effect of population age on movement distance," *PNAS*, vol. **105**, pp. 19090-19095, December 9, 2008 2008.
- [56] V. A. Drake, J. W. Chapman, *et al.*, "Ventral-aspect radar cross sections and polarization patterns of insects at X band and their relation to size and form," *International Journal of Remote Sensing*, vol. **38**, pp. 5022-5044, 2017/09/17 2017.
- [57] T. Alerstam, J. Backman, *et al.*, "A polar system of intercontinental bird migration," *Proc. R. Soc. B*, vol. **274**, pp. 2523-30, Oct 22 2007.
- [58] S. Jansson and M. Brydegaard, "Passive Entomological Lidar," *IEEE Photonics*, 2017.
- [59] S. Butail, *Motion reconstruction of animal groups: From schooling fish to swarming mosquitoes*: University of Maryland, College Park, 2012.
- [60] E. R. Mullen, P. Rutschman, *et al.*, "Laser system for identification, tracking, and control of flying insects," *Optics Express*, vol. **24**, pp. 11828-11838, 2016/05/30 2016.
- [61] J. K. Westbrook, R. S. Eyster, and W. W. Wolf, "WSR-88D Doppler radar detection of corn earworm moth migration," *International Journal of Biometeorology*, vol. **58**, pp. 931-940, 2014/07/01 2014.
- [62] B. Torvik, K. E. Olsen, and H. Griffiths, "K-band radar signature analysis of a flying mallard duck," in *Radar Symposium (IRS), 2013 14th International*, 2013, pp. 584-591.
- [63] C. E. Wainwright, P. M. Stepanian, *et al.*, "The movement of small insects in the convective boundary layer: linking patterns to processes," *Scientific Reports*, vol. **7**, p. 5438, 2017.
- [64] C. V. Nguyen, D. R. Lovell, *et al.*, "Capturing Natural-Colour 3D Models of Insects for Species Discovery and Diagnostics," *PLOS ONE*, vol. **9**, p. e94346, 2014.