

Congenital Anomalies in China

Austin Carter

Introduction

National achievement of SDG. Subnational achievement? Reference to 2016 paper to demonstrate growing importance of birth defects in U5M (Cui, He, Kang, Li, Miao, Shen, Zhu, Li, and Wang, 2016). Link the impact of birth defects to traditional measures of health (LE and DALYs)

Methods

Data

Mortality

China Disease Surveillance Points (DSP) for 1991-2002, sample registration system (SRS) data coded as verbal autopsy (VA). VA is method for determining the cause of death through an interview of relatives or associates of the deceased. China DSP for 2004-2012 coded as vital registration (VR) data. China National Maternal and Child Health Surveillance system for 1996-2012.

Non-fatal

China National Maternal and Child Health Surveillance System for Congenital Anomalies for 1996-2012. China hospital data for 2013-2015.

Analysis

Life Expectancy Decomposition

In order to summarize the impact of changes in mortality from CAs, we decompose improvements in life expectancy across age groups and causes of death. Specifically, we assess the proportion of gains in life expectancy resulting from the improvements in CA mortality and calculate expected life expectancy after removal of CA mortality in infants and children 1-4 years old. We implement the Pollard method for decomposition using life tables and cause-specific mortality from GBD 2016. This method assumes independence of causes of death, enabling the construction of a cause-deleted life table through calculation of the probability of survival in age group a in the absence of CA mortality:

$$p_{-i} = \frac{p(a)}{p_i(a)}$$

where $p(a)$ is probability of survival in the all-cause life table and $p_i(a) = e^{\mu_i(a)}$, $\mu_i(a)$ being age-specific mortality from CA. Setting the life table radix, l_0 , to 1. Update a_x in under-1, 1-4, 5-9, and the terminal age group to:

$$a_{x,-i} = n + \frac{(1 - (\frac{\mu_i(x)}{\mu(x)})) * (1 - p_x)}{(1 - p_{x,-i}) * (a_x - n)}$$

and total life years lived in the terminal age group is set to:

$${}_nL_{x,-i} = \frac{e_x}{1 - \frac{\mu_i(x)}{\mu(x)}} * l_{x,-i}$$

The difference in life expectancy between years can be decomposed into the contributions of each cause of deaths according to:

$$e^*(0) - e(0) = \sum_{i=1}^n \sum_{x=0}^{\omega} ({}_nL_{x,i}^* - {}_nL_{x,i}) \left(\frac{{}_nL_{x,-i}^* + {}_nL_{x,-i}}{2n} \right)$$

DALY Decomposition

$$T_{asgt} = A_{sgt} * B_{asgt} * C_{asgt}$$

$$E_A = (A_2 - A_1) \left(\frac{B_1 * C_1 + B_2 * C_2}{3} + \frac{B_1 * C_2 + B_2 * C_1}{6} \right)$$

Results

LE Decomposition

DALY Decomposition

Discussion

References

```
## [1] H. Cui, C. He, L. Kang, et al. "Under-5-Years Child Mortality
## Due to Congenital Anomalies". In: _American Journal of Preventive
## Medicine_ 50.5 (May. 2016), pp. 663-671. ISSN: 0749-3797. DOI:
## 10.1016/j.amepre.2015.12.013. <URL:
## http://www.sciencedirect.com/science/article/pii/S074937971500793X>.
```