


ME 535 Homework Assignment No 1

Debabrata Auddya

Question 1: (Exercise 1.1)


a. Draw 20 points on an ellipse

```
theta = 0:pi/10:2*pi;  
x = 5 + 5*cos(theta);  
y = 3 + 3*sin(theta);  
plot(x,y,'r:o')  
xlabel("X axis")  
ylabel("Y axis")  
title("Plotting 20 points on an ellipse")
```



b. Draw a polygon connecting the above points


```
plot(x,y,'m')
```



Question 2: (Exercise 2.1)

Demonstrate how unequal tangent vector magnitude with the symmetric slope end condition would affect the resulting Hermite curves for the following coefficients, where $k = 0.5, 1, 1.5$ and 2 . Display the results graphically.

```
k = [0.5 1 1.5 2];
figure(1); cla;
colorstring = 'kbgr';
u = 0:0.01:1;
for i = 1:size(k,2)
    p = [4 4 0;24 4 0;0.8320*k(i) 0.5547*k(i) 0;0.8320 -0.5547 0];
    for j = 1:size(u,2)
        g = [u(j)^3 u(j)^2 u(j) 1];
        A = [2 -2 1 1; -3 3 -2 -1; 0 0 1 0; 1 0 0 0];
        x(j) = g*A*p(:,1);
        y(j) = g*A*p(:,2);
    end
    plot(x,y, 'Color', colorstring(i))
    xlabel("X axis")
    ylabel("Y axis")
    hold on
end
legend("k = 0.5","k = 1","k = 1.5","k = 2")
hold off
```



Question 3: (Exercise 2.2)

A Hermite curve is defined by two points. What is the equation for the curve, i.e. the polynomial form? Plot the curve.

```
n = 100;
m = 1/(n-1);
t = 0:m:1;
x = -6*t.^3 + 9*t.^2 + t;
y = -4*t.^2 + 4*t;
plot(x,y,'r')
xlabel("X axis")
ylabel("Y axis")
```

