


CONSTRAINTS ON SOUTH POLE SEDIMENTARY BASIN THICKNESS: A MODIFIED RECEIVER FUNCTION APPROACH USING FORWARD MODELING


A.M. REID-MCLAUGHLIN¹, M. KARRENBACH¹, & Z. ZHAN¹
¹ CALIFORNIA INSTITUTE OF TECHNOLOGY, ² OPTASense



INTRODUCTION




RECEIVER FUNCTIONS ON ICE SHEETS



REFERENCES

- Anandkrishnan, S., & Whittall, J. P. (2004). Anisotropic geological sedimentary layer thickness from receiver function analysis. *Journal of Geophysical Research*, 109, B07307. <https://doi.org/10.1029/2003JB002307>
- Chen, J., & Li, X. (2019). A novel method for estimating the thickness of the sedimentary layer beneath the South Pole region. *Journal of Geophysical Research*, 124, 10, 10, 10, 10. <https://doi.org/10.1029/2019JB017000>
- Yeck, J. L., & Whittall, J. P. (2019). Sequential layer stripping to obtain accurate crustal thickness beneath sedimentary basins. *Journal of Geophysical Research*, 124, 10, 10, 10. <https://doi.org/10.1029/2019JB017000>
- Yu, X., Li, X., & Li, X. (2013). Determining crustal structure beneath active plateaus: applying a new velocity anisotropy layer using receiver functions. *Journal of Geophysical Research*, 118, 10, 10, 10. <https://doi.org/10.1029/2012JB009100>

RESULTS



CONCLUSION

