RWorksheet_Barrientos#3b.Rmd

Auderie Josh Barrientos

2024-10-13

#1.
#a.

```r
data <- data.frame(
  Respondents = 1:20,
  Sex = c(2, 1, 2, 2, 1, 1, 2, 2, 2, 1, 1, 2, 2, 1, 2, 2, 1, 2, 1, 2),
  FatherOccupation = c(1, 2, 3, 1, 2, 1, 3, 2, 3, 3, 1, 3, 2, 1, 3, 1, 3, 3, 1, 1),
  Persons_at_Home = c(5, 7, 3, 5, 5, 3, 6, 6, 7, 7, 3, 7, 4, 7, 8, 8, 3, 11, 8, 6),
  Siblings_at_School = c(6, 4, 3, 2, 3, 3, 5, 5, 4, 5, 3, 7, 5, 2, 1, 3, 1, 5, 3, 2),
  Types_of_Houses = c(1, 2, 3, 1, 3, 1, 3, 3, 3, 1, 3, 3, 3, 1, 3, 3, 3, 3, 3, 2)
)
data
```

```
##    Respondents Sex FatherOccupation Persons_at_Home Siblings_at_School
## 1            1   2                1               5                  6
## 2            2   1                2               7                  4
## 3            3   2                3               3                  3
## 4            4   2                1               5                  2
## 5            5   1                2               5                  3
## 6            6   1                1               3                  3
## 7            7   2                3               6                  5
## 8            8   2                2               6                  5
## 9            9   2                3               7                  4
## 10          10   1                3               7                  5
## 11          11   1                1               3                  3
## 12          12   2                3               7                  7
## 13          13   2                2               4                  5
## 14          14   1                1               7                  2
## 15          15   2                3               8                  1
## 16          16   2                1               8                  3
## 17          17   1                3               3                  1
## 18          18   2                3              11                  5
## 19          19   1                1               8                  3
## 20          20   2                1               6                  2
##    Types_of_Houses
## 1                1
## 2                2
## 3                3
## 4                1
## 5                3
## 6                1
## 7                3
## 8                3
```

```
## 9                  3
## 10                 1
## 11                 3
## 12                 3
## 13                 3
## 14                 1
## 15                 3
## 16                 3
## 17                 3
## 18                 3
## 19                 3
## 20                 2
```

#b.

```
str(data)
```

```
## 'data.frame':    20 obs. of  6 variables:
##  $ Respondents      : int  1 2 3 4 5 6 7 8 9 10 ...
##  $ Sex              : num  2 1 2 2 1 1 2 2 2 1 ...
##  $ FatherOccupation : num  1 2 3 1 2 1 3 2 3 3 ...
##  $ Persons_at_Home  : num  5 7 3 5 5 3 6 6 7 7 ...
##  $ Siblings_at_School: num  6 4 3 2 3 3 5 5 4 5 ...
##  $ Types_of_Houses  : num  1 2 3 1 3 1 3 3 3 1 ...
```

```
summary(data)
```

```
##    Respondents          Sex       FatherOccupation Persons_at_Home
##  Min.   : 1.00   Min.   :1.0   Min.   :1       Min.   : 3.00
##  1st Qu.: 5.75   1st Qu.:1.0   1st Qu.:1       1st Qu.: 4.75
##  Median :10.50   Median :2.0   Median :2       Median : 6.00
##  Mean   :10.50   Mean   :1.6   Mean   :2       Mean   : 5.95
##  3rd Qu.:15.25   3rd Qu.:2.0   3rd Qu.:3       3rd Qu.: 7.00
##  Max.   :20.00   Max.   :2.0   Max.   :3       Max.   :11.00
##  Siblings_at_School Types_of_Houses
##  Min.   :1.00       Min.   :1.00
##  1st Qu.:2.75       1st Qu.:1.75
##  Median :3.00       Median :3.00
##  Mean   :3.60       Mean   :2.40
##  3rd Qu.:5.00       3rd Qu.:3.00
##  Max.   :7.00       Max.   :3.00
```

#c.

```
mean(data$Siblings_at_School)
```

```
## [1] 3.6
```

#d.

```
data[1:2, ]
```

```
##   Respondents Sex FatherOccupation Persons_at_Home Siblings_at_School
## 1           1   2                1               5                  6
## 2           2   1                2               7                  4
##   Types_of_Houses
## 1               1
## 2               2
```

```
#e.
data[c(3, 5), c(2, 4)]
```

```
##   Sex Persons_at_Home
## 3   2               3
## 5   1               5
```

```
#f.
types_houses <- data$Types_of_Houses
types_houses
```

```
##  [1] 1 2 3 1 3 1 3 3 3 1 3 3 3 1 3 3 3 3 3 2
```

```
#g.
Male_Farmers <- subset(data, Sex == 1 & FatherOccupation == 1)
Male_Farmers
```

```
##     Respondents Sex FatherOccupation Persons_at_Home Siblings_at_School
## 6             6   1                1               3                  3
## 11           11   1                1               3                  3
## 14           14   1                1               7                  2
## 19           19   1                1               8                  3
##     Types_of_Houses
## 6                 1
## 11                3
## 14                1
## 19                3
```

```
#h.
Female_Siblings <- subset(data, Sex == 2 & Siblings_at_School >= 5)
Female_Siblings
```

```
##     Respondents Sex FatherOccupation Persons_at_Home Siblings_at_School
## 1             1   2                1               5                  6
## 7             7   2                3               6                  5
## 8             8   2                2               6                  5
## 12           12   2                3               7                  7
## 13           13   2                2               4                  5
## 18           18   2                3              11                  5
##     Types_of_Houses
## 1                 1
## 7                 3
## 8                 3
## 12                3
## 13                3
## 18                3
```

```
#2.
#a.
df = data.frame(Ints=integer(),

Doubles=double(), Characters=character(),
Logicals=logical(),
Factors=factor(),
```

```
stringsAsFactors=FALSE)

print("Structure of the empty dataframe:")

## [1] "Structure of the empty dataframe:"
print(str(df))

## 'data.frame':    0 obs. of  5 variables:
##  $ Ints      : int
##  $ Doubles   : num
##  $ Characters: chr
##  $ Logicals  : logi
##  $ Factors   : Factor w/ 0 levels:
## NULL
```

#3.
#a.

```
datac <- read.csv("HouseholdData.csv")
datac
```

```
##    Respondents    Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 1            1   Male                 1             5                2
## 2            2 Female                 2             7                3
## 3            3 Female                 3             3                0
## 4            4   Male                 3             8                5
## 5            5   Male                 1             6                2
## 6            6 Female                 2             4                3
## 7            7 Female                 2             4                1
## 8            8   Male                 3             2                2
## 9            9 Female                 1            11                6
## 10          10   Male                 3             6                2
##    TypesOfHouses
## 1           Wood
## 2       Congrete
## 3       Congrete
## 4           wood
## 5  Semi-congrete
## 6  Semi-congrete
## 7           Wood
## 8  Semi-congrete
## 9  Semi-congrete
## 10      Congrete
```

#b.

```
datac$Sex <- factor(datac$Sex, levels = c("Male", "Female"), labels = c(1,2))
datac
```

```
##    Respondents Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 1            1   1                 1             5                2
## 2            2   2                 2             7                3
## 3            3   2                 3             3                0
## 4            4   1                 3             8                5
## 5            5   1                 1             6                2
## 6            6   2                 2             4                3
```

```
## 7                7   2                    2              4                 1
## 8                8   1                    3              2                 2
## 9                9   2                    1             11                 6
## 10              10   1                    3              6                 2
##    TypesOfHouses
## 1           Wood
## 2       Congrete
## 3       Congrete
## 4           wood
## 5  Semi-congrete
## 6  Semi-congrete
## 7           Wood
## 8  Semi-congrete
## 9  Semi-congrete
## 10      Congrete
```

#c.

```
datac$TypesOfHouses <- factor(datac$TypesOfHouses, levels = c("Wood", "Congrete", "Semi-congrete"), lab
datac
```

```
##    Respondents Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 1            1   1                 1             5                2
## 2            2   2                 2             7                3
## 3            3   2                 3             3                0
## 4            4   1                 3             8                5
## 5            5   1                 1             6                2
## 6            6   2                 2             4                3
## 7            7   2                 2             4                1
## 8            8   1                 3             2                2
## 9            9   2                 1            11                6
## 10          10   1                 3             6                2
##    TypesOfHouses
## 1              1
## 2              2
## 3              2
## 4           <NA>
## 5              3
## 6              3
## 7              1
## 8              3
## 9              3
## 10             2
```

#d.

```
datac$FathersOccupation <- factor(datac$FathersOccupation, levels = c(1,2,3), labels = c("Farmer", "Driv
datac
```

```
##    Respondents Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 1            1   1            Farmer             5                2
## 2            2   2            Driver             7                3
## 3            3   2            Others             3                0
## 4            4   1            Others             8                5
## 5            5   1            Farmer             6                2
## 6            6   2            Driver             4                3
## 7            7   2            Driver             4                1
```

```
## 8            8  1         Others            2              2
## 9            9  2         Farmer           11              6
## 10          10  1         Others            6              2
##    TypesOfHouses
## 1             1
## 2             2
## 3             2
## 4          <NA>
## 5             3
## 6             3
## 7             1
## 8             3
## 9             3
## 10            2
```

#e.

```
FemaleDriverDad <- subset(datac, Sex == 2 & FathersOccupation =="Driver")
FemaleDriverDad
```

```
##   Respondents Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 2           2   2            Driver             7                3
## 6           6   2            Driver             4                3
## 7           7   2            Driver             4                1
##   TypesOfHouses
## 2             2
## 6             3
## 7             1
```

#f.

```
manysiblings <- subset(datac, Respondents & SiblingsAtSchool >= 5)
manysiblings
```

```
##   Respondents Sex FathersOccupation PersonsAtHome SiblingsAtSchool
## 4           4   1            Others             8                5
## 9           9   2            Farmer            11                6
##   TypesOfHouses
## 4          <NA>
## 9             3
```

#4. The graph shows thw Sentiments Of Tweets Per Day, from July 14, 2020 to July 21, 2020.The red shows negative, orange shows neutral, and blue shows a positive.As the graph shows that the negative tweets is the highest of the all.