

# Introduction to linear regression

Peter Gatica

```
{r global_options, include=FALSE} #knitr::opts_chunk$set(eval
= TRUE, results = TRUE, fig.show = "show", message = FALSE,
warning = FALSE) #
```

The Human Freedom Index is a report that attempts to summarize the idea of “freedom” through a bunch of different variables for many countries around the globe. It serves as a rough objective measure for the relationships between the different types of freedom - whether it’s political, religious, economical or personal freedom - and other social and economic circumstances. The Human Freedom Index is an annually co-published report by the Cato Institute, the Fraser Institute, and the Liberales Institut at the Friedrich Naumann Foundation for Freedom.

In this lab, you’ll be analyzing data from Human Freedom Index reports from 2008-2016. Your aim will be to summarize a few of the relationships within the data both graphically and numerically in order to find which variables can help tell a story about freedom.

## Getting Started

### Load packages

In this lab, you will explore and visualize the data using the **tidyverse** suite of packages. The data can be found in the companion package for OpenIntro resources, **openintro**.

Let’s load the packages.

```
library(DATA606)
```

```
## Loading required package: shiny
```

```
## Loading required package: openintro
```

```
## Loading required package: airports
```

```
## Loading required package: cherryblossom
```

```
## Loading required package: usdata
```

```
## Loading required package: OIdata
```

```
## Loading required package: RCurl
```

```

## Loading required package: maps

## Loading required package: ggplot2

## Loading required package: markdown

##
## Welcome to CUNY DATA606 Statistics and Probability for Data Analytics
## This package is designed to support this course. The text book used
## is OpenIntro Statistics, 3rd Edition. You can read this by typing
## vignette('os3') or visit www.OpenIntro.org.
##
## The getLabs() function will return a list of the labs available.
##
## The demo(package='DATA606') will list the demos that are available.

##
## Attaching package: 'DATA606'

## The following objects are masked from 'package:openintro':
##
##   calc_streak, present, qqnormsim

## The following object is masked from 'package:utils':
##
##   demo

library(tidyverse)

## -- Attaching packages ----- tidyverse 1.3.0 --

## v tibble  3.1.1    v dplyr   1.0.2
## v tidyr   1.1.2    v stringr 1.4.0
## v readr   1.4.0    v forcats 0.5.1
## v purrr   0.3.4

## -- Conflicts ----- tidyverse_conflicts() --
## x tidyr::complete() masks Rcurl::complete()
## x dplyr::filter()   masks stats::filter()
## x dplyr::lag()      masks stats::lag()
## x purrr::map()      masks maps::map()

library(openintro)
library(ggplot2)

```

## Creating a reproducible lab report

To create your new lab report, in RStudio, go to New File -> R Markdown... Then, choose From Template and then choose Lab Report for OpenIntro Statistics Labs from the list of templates.

## The data

The data we're working with is in the openintro package and it's called `hfi`, short for Human Freedom Index.

```
data("hfi")
```

```
head(hfi,10)
```

```
## # A tibble: 10 x 123
##   year ISO_code countries region pf_rol_procedur~ pf_rol_civil pf_rol_criminal
##   <dbl> <chr>    <chr>    <chr>          <dbl>         <dbl>         <dbl>
## 1  2016 ALB      Albania  Easte~          6.66          4.55          4.67
## 2  2016 DZA      Algeria  Middl~          NA           NA           NA
## 3  2016 AGO      Angola   Sub-S~          NA           NA           NA
## 4  2016 ARG      Argentina Latin~          7.10          5.79          4.34
## 5  2016 ARM      Armenia  Cauca~          NA           NA           NA
## 6  2016 AUS      Australia Ocean~          8.44          7.53          7.36
## 7  2016 AUT      Austria  Weste~          8.97          7.87          7.67
## 8  2016 AZE      Azerbaij~ Cauca~          NA           NA           NA
## 9  2016 BHS      Bahamas Latin~          6.93          6.01          6.26
## 10 2016 BHR      Bahrain  Middl~          NA           NA           NA
## # ... with 116 more variables: pf_rol <dbl>, pf_ss_homicide <dbl>,
## #   pf_ss_disappearances_disap <dbl>, pf_ss_disappearances_violent <dbl>,
## #   pf_ss_disappearances_organized <dbl>,
## #   pf_ss_disappearances_fatalities <dbl>, pf_ss_disappearances_injuries <dbl>,
## #   pf_ss_disappearances <dbl>, pf_ss_women_fgm <dbl>,
## #   pf_ss_women_missing <dbl>, pf_ss_women_inheritance_widows <dbl>,
## #   pf_ss_women_inheritance_daughters <dbl>, pf_ss_women_inheritance <dbl>,
## #   pf_ss_women <dbl>, pf_ss <dbl>, pf_movement_domestic <dbl>,
## #   pf_movement_foreign <dbl>, pf_movement_women <dbl>, pf_movement <dbl>,
## #   pf_religion_estop_establish <dbl>, pf_religion_estop_operate <dbl>,
## #   pf_religion_estop <dbl>, pf_religion_harassment <dbl>,
## #   pf_religion_restrictions <dbl>, pf_religion <dbl>,
## #   pf_association_association <dbl>, pf_association_assembly <dbl>,
## #   pf_association_political_establish <dbl>,
## #   pf_association_political_operate <dbl>, pf_association_political <dbl>,
## #   pf_association_prof_establish <dbl>, pf_association_prof_operate <dbl>,
## #   pf_association_prof <dbl>, pf_association_sport_establish <dbl>,
## #   pf_association_sport_operate <dbl>, pf_association_sport <dbl>,
## #   pf_association <dbl>, pf_expression_killed <dbl>,
## #   pf_expression_jailed <dbl>, pf_expression_influence <dbl>,
## #   pf_expression_control <dbl>, pf_expression_cable <dbl>,
## #   pf_expression_newspapers <dbl>, pf_expression_internet <dbl>,
## #   pf_expression <dbl>, pf_identity_legal <dbl>,
## #   pf_identity_parental_marriage <dbl>, pf_identity_parental_divorce <dbl>,
## #   pf_identity_parental <dbl>, pf_identity_sex_male <dbl>,
## #   pf_identity_sex_female <dbl>, pf_identity_sex <dbl>,
## #   pf_identity_divorce <dbl>, pf_identity <dbl>, pf_score <dbl>,
## #   pf_rank <dbl>, ef_government_consumption <dbl>,
## #   ef_government_transfers <dbl>, ef_government_enterprises <dbl>,
## #   ef_government_tax_income <dbl>, ef_government_tax_payroll <dbl>,
## #   ef_government_tax <dbl>, ef_government <dbl>, ef_legal_judicial <dbl>,
## #   ef_legal_courts <dbl>, ef_legal_protection <dbl>, ef_legal_military <dbl>,
```

```
## # ef_legal_integrity <dbl>, ef_legal_enforcement <dbl>,
## # ef_legal_restrictions <dbl>, ef_legal_police <dbl>, ef_legal_crime <dbl>,
## # ef_legal_gender <dbl>, ef_legal <dbl>, ef_money_growth <dbl>,
## # ef_money_sd <dbl>, ef_money_inflation <dbl>, ef_money_currency <dbl>,
## # ef_money <dbl>, ef_trade_tariffs_revenue <dbl>,
## # ef_trade_tariffs_mean <dbl>, ef_trade_tariffs_sd <dbl>,
## # ef_trade_tariffs <dbl>, ef_trade_regulatory_nontariff <dbl>,
## # ef_trade_regulatory_compliance <dbl>, ef_trade_regulatory <dbl>,
## # ef_trade_black <dbl>, ef_trade_movement_foreign <dbl>,
## # ef_trade_movement_capital <dbl>, ef_trade_movement_visit <dbl>,
## # ef_trade_movement <dbl>, ef_trade <dbl>,
## # ef_regulation_credit_ownership <dbl>, ef_regulation_credit_private <dbl>,
## # ef_regulation_credit_interest <dbl>, ef_regulation_credit <dbl>,
## # ef_regulation_labor_minwage <dbl>, ef_regulation_labor_firing <dbl>,
## # ef_regulation_labor_bargain <dbl>, ef_regulation_labor_hours <dbl>, ...
```

1. What are the dimensions of the dataset?


Answer: The dimensions in the dataset are year, ISO\_code, countries and region.

2. What type of plot would you use to display the relationship between the personal freedom score, `pf_score`, and one of the other numerical variables? Plot this relationship using the variable `pf_expression_control` as the predictor. Does the relationship look linear? If you knew a country's `pf_expression_control`, or its score out of 10, with 0 being the most, of political pressures and controls on media content, would you be comfortable using a linear model to predict the personal freedom score?

Answer: I would use a scatterplot to display the relationship between the `pf_score` and another numerical variable.

```
ggplot(data = hfi, aes(x = pf_expression_control, y = pf_score)) + geom_point()
```

```
## Warning: Removed 80 rows containing missing values (geom_point).
```



If the relationship looks linear, we can quantify the strength of the relationship with the correlation coefficient.

```
hfi %>%
  summarise(cor(pf_expression_control, pf_score, use = "complete.obs"))
```

```
## # A tibble: 1 x 1
##   'cor(pf_expression_control, pf_score, use = "complete.obs")'
##                                     <dbl>
## 1                                     0.796
```

Here, we set the `use` argument to “complete.obs” since there are some observations of NA.

## Sum of squared residuals

In this section, you will use an interactive function to investigate what we mean by “sum of squared residuals”. You will need to run this function in your console, not in your markdown document. Running the function also requires that the `hfi` dataset is loaded in your environment.

Think back to the way that we described the distribution of a single variable. Recall that we discussed characteristics such as center, spread, and shape. It’s also useful to be able to describe the relationship of two numerical variables, such as `pf_expression_control` and `pf_score` above.


1. Looking at your plot from the previous exercise, describe the relationship between these two variables. Make sure to discuss the form, direction, and strength of the relationship as well as any unusual observations.

Answer: There is a positive correlation between `pf_expression_control` and `pf_score`. As you can see from the positive upward trend of the observations. The correlation is also strong at almost 80%

Just as you've used the mean and standard deviation to summarize a single variable, you can summarize the relationship between these two variables by finding the line that best follows their association. Use the following interactive function to select the line that you think does the best job of going through the cloud of points.

```
hfi_no_na <- hfi[!is.na(hfi$pf_expression_control),]
hfi_no_na <- hfi[!is.na(hfi$pf_score),]
```

```
#this isn't working, wondering if we should just remove rows with NA
plot_ss(x = hfi_no_na$pf_expression_control, y = hfi_no_na$pf_score)
```




```
## Click two points to make a line.
## Call:
## lm(formula = y ~ x, data = pts)
##
## Coefficients:
## (Intercept)          x
##      4.6171      0.4914
##
## Sum of Squares:  952.153
```

After running this command, you'll be prompted to click two points on the plot to define a line. Once you've done that, the line you specified will be shown in black and the residuals in blue. Note that there are 30 residuals, one for each of the 30 observations. Recall that the residuals are the difference between the observed values and the values predicted by the line:

$$e_i = y_i - \hat{y}_i$$

The most common way to do linear regression is to select the line that minimizes the sum of squared residuals. To visualize the squared residuals, you can rerun the plot command and add the argument `showSquares = TRUE`.

```
plot_ss(x = hfi_no_na$pf_expression_control, y = hfi_no_na$pf_score, showSquares = TRUE)
```



```
## Click two points to make a line.
## Call:
## lm(formula = y ~ x, data = pts)
##
## Coefficients:
## (Intercept)      x
##      4.6171    0.4914
##
## Sum of Squares:  952.153
```

Note that the output from the `plot_ss` function provides you with the slope and intercept of your line as well as the sum of squares.

1. Using `plot_ss`, choose a line that does a good job of minimizing the sum of squares. Run the function several times. What was the smallest sum of squares that you got? How does it compare to your neighbors?

Answer: I ran the `plot_ss` in an `r` script in RStudio and the smallest Sum of Squares that I got was 952.153.

## The linear model

It is rather cumbersome to try to get the correct least squares line, i.e. the line that minimizes the sum of squared residuals, through trial and error. Instead, you can use the `lm` function in R to fit the linear model (a.k.a. regression line).

```
(m1 <- lm(pf_score ~ pf_expression_control, data = hfi_no_na))
```

```
##
## Call:
## lm(formula = pf_score ~ pf_expression_control, data = hfi_no_na)
##
## Coefficients:
##           (Intercept)  pf_expression_control
##                4.6171                0.4914
```

The first argument in the function `lm` is a formula that takes the form  $y \sim x$ . Here it can be read that we want to make a linear model of `pf_score` as a function of `pf_expression_control`. The second argument specifies that R should look in the `hfi` data frame to find the two variables.

The output of `lm` is an object that contains all of the information we need about the linear model that was just fit. We can access this information using the summary function.

```
summary(m1)
```

```
##
## Call:
## lm(formula = pf_score ~ pf_expression_control, data = hfi_no_na)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.8467 -0.5704  0.1452  0.6066  3.2060
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    4.61707    0.05745   80.36  <2e-16 ***
## pf_expression_control 0.49143    0.01006   48.85  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.8318 on 1376 degrees of freedom
## Multiple R-squared:  0.6342, Adjusted R-squared:  0.634
## F-statistic: 2386 on 1 and 1376 DF, p-value: < 2.2e-16
```

Let's consider this output piece by piece. First, the formula used to describe the model is shown at the top. After the formula you find the five-number summary of the residuals. The "Coefficients" table shown next is key; its first column displays the linear model's y-intercept and the coefficient of `at_bats`. With this table, we can write down the least squares regression line for the linear model:


$$\hat{y} = 4.61707 + 0.49143 \times pf\_expression\_control$$

One last piece of information we will discuss from the summary output is the Multiple R-squared, or more simply,  $R^2$ . The  $R^2$  value represents the proportion of variability in the response variable that is explained by the explanatory variable. For this model, 63.42% of the variability in runs is explained by `at_bats`.

1. Fit a new model that uses `pf_expression_control` to predict `hf_score`, or the total human freedom score. Using the estimates from the R output, write the equation of the regression line. What does the slope tell us in the context of the relationship between human freedom and the amount of political pressure on media content?



```
plot_ss(x = hfi_no_na$pf_expression_control, y = hfi_no_na$pf_score)
```




```
## Click two points to make a line.
## Call:
## lm(formula = y ~ x, data = pts)
##
## Coefficients:
## (Intercept)          x
##      4.6171      0.4914
##
## Sum of Squares:  952.153
```

## Prediction and prediction errors

Let's create a scatterplot with the least squares line for m1 laid on top.

```
ggplot(data = hfi_no_na, aes(x = pf_expression_control, y = pf_score)) +
  geom_point() +
  stat_smooth(method = "lm", se = FALSE)
```

```
## 'geom_smooth()' using formula 'y ~ x'
```



Here, we are literally adding a layer on top of our plot. `geom_smooth` creates the line by fitting a linear model. It can also show us the standard error `se` associated with our line, but we'll suppress that for now.

This line can be used to predict  $y$  at any value of  $x$ . When predictions are made for values of  $x$  that are beyond the range of the observed data, it is referred to as *extrapolation* and is not usually recommended. However, predictions made within the range of the data are more reliable. They're also used to compute the residuals.

1. If someone saw the least squares regression line and not the actual data, how would they predict a country's personal freedom score for one with a 6.7 rating for `pf_expression_control`? Is this an overestimate or an underestimate, and by how much? In other words, what is the residual for this prediction?


Answer: Estimate where 6.7 lies and then mark where the line intersects the x and y axis. This is your predicted value or 8. I would think that this is overestimated. The residual would be the difference between the observed residual and the predicted residual.

## Model diagnostics

To assess whether the linear model is reliable, we need to check for (1) linearity, (2) nearly normal residuals, and (3) constant variability.

**Linearity:** You already checked if the relationship between `pf_score` and '`pf_expression_control`' is linear using a scatterplot. We should also verify this condition with a plot of the residuals vs. fitted (predicted) values.

```
ggplot(data = m1, aes(x = .fitted, y = .resid)) +
  geom_point() +
  geom_hline(yintercept = 0, linetype = "dashed") +
  xlab("Fitted values") +
  ylab("Residuals")
```




Notice here that `m1` can also serve as a data set because stored within it are the fitted values ( $\hat{y}$ ) and the residuals. Also note that we're getting fancy with the code here. After creating the scatterplot on the first layer (first line of code), we overlay a horizontal dashed line at  $y = 0$  (to help us check whether residuals are distributed around 0), and we also rename the axis labels to be more informative.

1. Is there any apparent pattern in the residuals plot? What does this indicate about the linearity of the relationship between the two variables?

Answer: There is an apparent pattern in the residual plot where one can see that both positive and negative residuals are tightly packed around the least of squares line which suggests a high correlation score which suggests a strong correlation to a positive linear trend.


**Nearly normal residuals:** To check this condition, we can look at a histogram

```
ggplot(data = m1, aes(x = .resid)) +
  geom_histogram(binwidth = 25) +
  xlab("Residuals")
```



or a normal probability plot of the residuals.

```
ggplot(data = m1, aes(sample = .resid)) +  
  stat_qq()
```



Note that the syntax for making a normal probability plot is a bit different than what you're used to seeing: we set `sample` equal to the residuals instead of `x`, and we set a statistical method `qq`, which stands for “quantile-quantile”, another name commonly used for normal probability plots.

1. Based on the histogram and the normal probability plot, does the nearly normal residuals condition appear to be met?

Answer: Yes it does. There are not noticeable outliers that are influential points to affect the linearity of the observations.

#### Constant variability:

1. Based on the residuals vs. fitted plot, does the constant variability condition appear to be met?

Answer: Yes, constant variability is met because there is no fanning out of the observations.

---

### More Practice

- Choose another freedom variable and a variable you think would strongly correlate with it. Produce a scatterplot of the two variables and fit a linear model. At a glance, does there seem to be a linear relationship?

Answer: I have chosen `pf_religion`. There seems to be a positive linear relationship upwards concentrated on the right side of the graph and you would think this to be an obvious relationship since persons `pf_score` would be higher the more religious freedom a person has. However, the correlation which measures how good the model. In this case it is mediocre at 49.7% , I expected higher.


```
hfi_no_na <- hfi[!is.na(hfi$pf_religion),]
hfi_no_na <- hfi[!is.na(hfi$pf_expression),]
hfi_no_na <- hfi[!is.na(hfi$pf_score),]
```

```
ggplot(data = hfi_no_na, aes(x = pf_religion, y = pf_score)) +
  geom_point() +
  stat_smooth(method = "lm", se = FALSE)
```

```
## 'geom_smooth()' using formula 'y ~ x'
```

```
## Warning: Removed 10 rows containing non-finite values (stat_smooth).
```

```
## Warning: Removed 10 rows containing missing values (geom_point).
```



```
hfi_no_na %>%
  summarise(cor(pf_religion, pf_score, use = "complete.obs"))
```

```
## # A tibble: 1 x 1
##   'cor(pf_religion, pf_score, use = "complete.obs")'
##                                     <dbl>
## 1                                     0.497
```


```
(m3 <- lm(pf_score ~ pf_religion, data = hfi_no_na))
```

```
##  
## Call:  
## lm(formula = pf_score ~ pf_religion, data = hfi_no_na)  
##  
## Coefficients:  
## (Intercept)  pf_religion  
##      3.3098      0.4969
```

```
summary(m3)
```

```
##  
## Call:  
## lm(formula = pf_score ~ pf_religion, data = hfi_no_na)  
##  
## Residuals:  
##      Min       1Q   Median       3Q      Max   
## -4.3515 -0.8728 -0.0744  1.0532  2.3520   
##  
## Coefficients:  
##              Estimate Std. Error t value Pr(>|t|)      
## (Intercept)  3.30981    0.18744   17.66  <2e-16 ***   
## pf_religion  0.49689    0.02346   21.18  <2e-16 ***   
## ---  
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## Residual standard error: 1.17 on 1366 degrees of freedom  
##    (10 observations deleted due to missingness)  
## Multiple R-squared:  0.2472, Adjusted R-squared:  0.2467   
## F-statistic: 448.6 on 1 and 1366 DF,  p-value: < 2.2e-16
```

```
ggplot(data = m3, aes(x = .fitted, y = .resid)) +  
  geom_point() +  
  geom_hline(yintercept = 0, linetype = "dashed") +  
  xlab("Fitted values") +  
  ylab("Residuals")
```



- How does this relationship compare to the relationship between `pf_expression_control` and `pf_score`? Use the  $R^2$  values from the two model summaries to compare. Does your independent variable seem to predict your dependent one better? Why or why not?

Answer: The `pf_expression_control` to `pf_score` has a much more positive linear relationship than the model of `pf_religion` to `pf_score`. It surprises me because I would think that having more religious freedom would translate to a stronger linear relationship but the  $R^2$  seems to not indicate that with an  $R^2$  of 49.7%.

- What's one freedom relationship you were most surprised about and why? Display the model diagnostics for the regression model analyzing this relationship.