

Standard 13: Triple Integrals - Rectangular

Triple Integrals

The next step up from integrating over a two-dimensional region is integrating over a three-dimensional region. It should come as no surprise that we use a triple integral to integrate over a three-dimensional region. The general triple integral is $\iiint_E f(x,y,z) dV$.

rectangular-prism region

Over the box $B = [a,b] \times [c,d] \times [r,s] = \{(x,y,z) \mid a \leq x \leq b, c \leq y \leq d, r \leq z \leq s\}$, the triple integral $\iiint_B f(x,y,z) dV = \int_a^b \int_c^d \int_r^s f(x,y,z) dx dy dz$.

Note: we could rewrite this integral six different ways, $dx dy dz, dx dz dy, dy dx dz, dy dz dx, dz dx dy, dz dy dx$.

example. Evaluate $\iiint_B 8xyz dV$ for $B = [2,3] \times [1,2] \times [0,1]$.

$$\begin{aligned} \iiint_E 8xyz dx dy dz &= \int_2^3 \int_1^2 \int_0^1 8xyz dz dy dx \\ &= \int_2^3 \int_1^2 4xy z^2 \Big|_0^1 dy dx \\ &= \int_2^3 \int_1^2 4xy dy dx \\ &= \int_2^3 2xy^2 \Big|_1^2 dx \\ &= \int_2^3 2x(2^2 - 1^2) dx \\ &= \int_2^3 6x dx \\ &= 3x^2 \Big|_2^3 \\ &= 3(3^2) - 3(2^2) \\ &= 27 - 12 \\ &= 15 \end{aligned}$$

$$\begin{aligned} &= \int_1^2 \int_2^3 \int_0^1 8xyz dz dx dy \\ &= \int_1^2 \int_2^3 4xy z^2 \Big|_0^1 dx dy \\ &= \int_1^2 \int_2^3 4xy dx dy \\ &= \int_1^2 2xy^2 \Big|_2^3 dy \\ &= \int_1^2 2x(3^2 - 2^2) dy \\ &= \int_1^2 18x - 8x dy \\ &= \int_1^2 10x dy \\ &= 5x^2 \Big|_1^2 \\ &= 5(2^2) - 5(1^2) \\ &= 15 \end{aligned}$$


$$\begin{aligned} &= \int_0^1 \int_1^2 \int_2^3 8xyz dx dy dz \\ &= \int_0^1 \int_1^2 4x^2 y z \Big|_2^3 dy dz \\ &= \int_0^1 \int_1^2 4(3^2)yz - 4(2^2)yz dy dz \\ &= \int_0^1 \int_1^2 36yz - 16yz dy dz \\ &= \int_0^1 20 yz \Big|_1^2 dz \\ &= \int_0^1 10(z^2 z - 10(1)^2 z) dz \\ &= \int_0^1 30z dz \\ &= 15z \Big|_0^1 \\ &= 15 \end{aligned}$$

Recall the calculus I explanation of the double integral: $A = \int_a^b g_2(x) - g_1(x) dx = \int_a^b \int_{g_1(x)}^{g_2(x)} 1 dy dx$. We can make a similar assertion for triple integrals: $V = \iint_R u_2(x,y) - u_1(x,y) dA = \iint_R \int_{u_1(x,y)}^{u_2(x,y)} 1 dz dA = \iiint_E 1 dV$.

general region

We have three ways to describe a general region in three-dimension:


(i) $E = \{(x,y,z) \mid (x,y) \in D, u_1(x,y) \leq z \leq u_2(x,y)\}$



D is the shadow region in the xy-plane


$$\iiint_E f(x,y,z) dV = \iint_D \left[\int_{u_1(x,y)}^{u_2(x,y)} f(x,y,z) dz \right] dA$$

example. Evaluate $\iiint_E 2x dV$ where E is the region under the plane $2x+3y+z=6$ that lies in the first octant.



$$\begin{aligned} \iiint_E 2x dV &= \iint_D \left[\int_0^{6-2x-3y} 2x dz \right] dA \\ &= \int_0^3 \int_{-2/3x+2}^0 2x dz dy dx \end{aligned}$$


(ii) $E = \{(x,y,z) \mid (y,z) \in D, u_1(y,z) \leq x \leq u_2(y,z)\}$



D is the shadow region in the yz-plane


$$\iiint_E f(x,y,z) dV = \iint_D \left[\int_{u_1(y,z)}^{u_2(y,z)} f(x,y,z) dx \right] dA$$

example. Determine the volume of the region that lies behind the plane $x+y+z=8$ and in front of the region in the yz-plane that is bounded by $z=3/2\sqrt{y}$ and $z=3/4y$.



$$\begin{aligned} \iiint_E 1 dV &= \iint_D \left[\int_0^{8-y-z} 1 dx \right] dA \\ &= \int_0^4 \int_{3/4y}^{3/2\sqrt{y}} \int_0^{8-y-z} 1 dx dz dy \end{aligned}$$


(iii) $E = \{(x,y,z) \mid (x,z) \in D, u_1(x,z) \leq y \leq u_2(x,z)\}$



D is the shadow region in the xz-plane

$$\iiint_E f(x,y,z) dV = \iint_D \left[\int_{u_1(x,z)}^{u_2(x,z)} f(x,y,z) dy \right] dA$$

example. Evaluate $\iiint_E \sqrt{3x^2+3z^2} dV$ where E is the solid bounded by $y=2x^2+2z^2$ and the plane $y=8$.



$$\begin{aligned} \iiint_E \sqrt{3x^2+3z^2} dV &= \iint_D \left[\int_0^{8-2x^2-2z^2} \sqrt{3x^2+3z^2} dy \right] dA \\ &= \int_0^{2\pi} \int_0^2 \left[\int_0^{8-2r^2} \sqrt{3r^2} dy \right] r dr d\theta \end{aligned}$$