

Coping with climate change.

Implications of the Yellow-Bellied Marmot's (*Marmota flaviventris*) body mass evolution in the last half-century.

Augustin Birot (300444988)

2024-12-02

Table of contents

Introduction	2
Climate change (C.C.)	2
Body size as a Life-History Traits (LHT) and expected effect of global warming . . .	2
Q.G. and animal models	3
LHT coevolution	3
POLS	3
Phenotypic plasticity vs microevolution	3
I * E, G * E (individual variation in their plasticity)	3
Species and study site	3
Phenological mismatch??	3
Chapter 1 - Mechanisms	4
Chapter 2 - Methodology	5
Chapter 3 - Triggers	6
Chapter 4 - Implications	7
References	8

Introduction

Climate change (C.C.)

- The importance of C.C. and its impact in the near future is no longer in doubt (Intergovernmental Panel On Climate Change (Ipcc) 2023)
- Broadly, C.C. is [...] (Polar melting, etc.
- Which even impact human society (e.g., winter in Ottawa isn't the same anymore: Rideau Canal ice skating future is in jeopardy, the number of days with under -20°C is expected to severely decrease in the near future, etc. (!!! + FACT CHECK everything!))
- Main/Precise impacts of C.C. in natural environments
 - **Raising T°**: Explain + study case (!!!)
 - **Changing season length**: Explain + study case (try to find something at RMBL !!!)
 - **Environmental predictability**: Explain + study case (!!!)
 - **Drought events**: Explain + study case (!!!)
 - **Extreme weather events**: Explain + study case (!!!)
 - etc.

Ok, so, now, how does C.C. and these precise perturbations impacts concretely natural population? *Study cases* (!!!)

universal C.C. responses (C.C.)

Body size as a Life-History Traits (LHT) and expected effect of global warming

C.C. is expected to impact life history traits (LHT, i.e., traits impacting directly survival and reproduction, so individual's fitness Roff 1992).

Body mass is a LHT as in many species it has direct impact on survival and reproduction (explain + !!!)

(Daufresne et al. 2009), (Gardner et al. 2011), (Guillemain et al. 2010), (Sheridan and Bickford 2011), (Yom-Tov et al. 2008), (Ozgul et al. 2010)

And some authors argue that a decreasing body size could be one the universal C.C. response (!!!)

Because Bergmann's rules, which state that smaller body size should be expected in warmer environment as it raises the surface to volume ratio, thus favoring heat dissipation (Bergmann, 1847).

Thus, a general shrinking body size is expected with global warming. However opposite results at higher latitude yield objections of this theory and raise the need of more general study about that. Furthermore, these opposite results (i.e., increasing body mass at higher latitudes) can also be explained as C.C. is synonym to milder conditions in those latitudes. This change allows individuals to forage more and thus gain mass.

Q.G. and animal models

(Kruuk 2004) (Charmanier et al. 2014)

LHT coevolution

Traits can't evolve alone Gould & Lewontin (1979)

Need to show that with multivariate animal model, but no one has enough power for the models (Teplitsky et al. 2014)

POLS

(Dammhahn et al. 2018)

Phenotypic plasticity vs microevolution

I * E, G * E (individual variation in their plasticity)

(Nussey et al. 2007)

Link with body mass, individual can vary in their growing speed

I * A and G * A

Species and study site

Phenological mismatch??

Chapter 1 - Mechanisms

Marmot's Biology: What mechanisms are behind the body mass increase?

Growth? Baseline? Both?

Double random (Intercept, Slope)




Figure 1: Body mass trend over time cohort for females (black) and males (red) juveniles compared between a) the beginning of the active season (birth weight) and b) the end of the season (mass on August 15th).

Chapter 2 - Methodology

*Methodology: I * E detection with double random mixed models*

(Nussey et al. 2007) → double random

So we're doing something different → examining the residuals of the model (if I * E, still a lot of residual variance ?)

**Look at this one: (Westneat et al. 2015)

Chapter 3 - Triggers

Marmot's Biology: Which environmental factors have triggered the phenotypic shift?

E1 - E10 (T°, Precipitation, ...), Seasonal Gradient

Predators, Diet?

Chapter 4 - Implications

Marmot's Biology: What could be the implications of that for the population's future?

Manuscript models Body Mass/active season with survival

References

- Bergmann, C (1847), “About the relationships between heat conservation and body size of animals,” *Goett Stud*, 1, 595–708.
- Charmantier, A., Garant, D., and Kruuk, L. E. B. (2014), *Quantitative genetics in the wild*, Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199674237.001.0001>.
- Dammhahn, M., Dingemanse, N. J., Niemelä, P. T., and Réale, D. (2018), “Pace-of-life syndromes: A framework for the adaptive integration of behaviour, physiology and life history,” *Behavioral Ecology and Sociobiology*, 72, 62, s00265-018-2473-y. <https://doi.org/10.1007/s00265-018-2473-y>.
- Daufresne, M., Lengfellner, K., and Sommer, U. (2009), “Global warming benefits the small in aquatic ecosystems,” *Proceedings of the National Academy of Sciences*, 106, 12788–12793. <https://doi.org/10.1073/pnas.0902080106>.
- Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., and Heinsohn, R. (2011), “Declining body size: A third universal response to warming?” *Trends in Ecology & Evolution*, 26, 285–291. <https://doi.org/10.1016/j.tree.2011.03.005>.
- Guillemain, M., Elmberg, J., Gauthier-Clerc, M., Massez, G., Hearn, R., Champagnon, J., and Simon, G. (2010), “Wintering French Mallard and Teal Are Heavier and in Better Body Condition than 30 Years Ago: Effects of a Changing Environment?” *AMBIO*, 39, 170–180. <https://doi.org/10.1007/s13280-010-0020-9>.
- Intergovernmental Panel On Climate Change (Ipcc) (2023), *Climate Change 2022 – Impacts, Adaptation and Vulnerability: Working Group II Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press. <https://doi.org/10.1017/9781009325844>.
- Kruuk, L. E. B. (2004), “Estimating genetic parameters in natural populations using the ‘animal model’,” *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 359, 873–890. <https://doi.org/10.1098/rstb.2003.1437>.
- Nussey, D. H., Wilson, A. J., and Brommer, J. E. (2007), “The evolutionary ecology of individual phenotypic plasticity in wild populations,” *Journal of Evolutionary Biology*, 20, 831–844. <https://doi.org/10.1111/j.1420-9101.2007.01300.x>.
- Ozgul, A., Childs, D. Z., Oli, M. K., Armitage, K. B., Blumstein, D. T., Olson, L. E., Tuljapurkar, S., and Coulson, T. (2010), “Coupled dynamics of body mass and population growth in response to environmental change,” *Nature*, 466, 482–485. <https://doi.org/10.1038/nature09210>.
- Roff, D. A. (1992), “The evolution of life histories : Theory and analysis.”
- Sheridan, J. A., and Bickford, D. (2011), “Shrinking body size as an ecological response to climate change,” *Nature Climate Change*, 1, 401–406. <https://doi.org/10.1038/nclimate1259>.
- Teplitsky, C., Robinson, M. R., and Merilä, J. (2014), “Evolutionary potential and constraints in wild populations,” in *Quantitative Genetics in the Wild*, eds. A. Charmantier, D. Garant, and L. E. B. Kruuk, Oxford University Press Oxford, pp. 190–208. <https://doi.org/10.1093/acprof:oso/9780199674237.003.0012>.
- Westneat, D. F., Wright, J., and Dingemanse, N. J. (2015), “The biology hidden inside residual

within-individual phenotypic variation,” *Biological Reviews*, 90, 729–743. <https://doi.org/10.1111/brv.12131>.

Yom-Tov, Y., Yom-Tov, S., and Jarrell, G. (2008), “Recent increase in body size of the American marten *Martes americana* in Alaska: GLOBAL WARMING AND BODY SIZE OF THE AMERICAN MARTEN,” *Biological Journal of the Linnean Society*, 93, 701–707. <https://doi.org/10.1111/j.1095-8312.2007.00950.x>.