

Design and Usability Testing of a Mobile Phone-Based Patient Management System for Women in Rural Kenya

Amogh Karnik
M.Sc. Candidate

Duke Global Health Institute

March 29, 2014

OVERVIEW

Introduction

Maternal Mortality

Methods

Setting

Human-Centered Design

Results

Qualitative Results

Quantitative Results

Discussion

Frame 1

Every day, 800 women die of pregnancy-related complications around the world.

Every day, 800 women die of pregnancy-related complications around the world.

The most common causes of maternal death are hemorrhage and infection.

Every day, 800 women die of pregnancy-related complications around the world.

The most common causes of maternal death are hemorrhage and infection.

Most of these deaths take place during or within the first few days after childbirth.


Every day, 800 women die of pregnancy-related complications around the world.

The most common causes of maternal death are hemorrhage and infection.

Most of these deaths take place during or within the first few days after childbirth.

In Sub-Saharan Africa, 1 of 16 women will die of pregnancy-related complications.

Most maternal deaths are avoidable.



RESEARCH SITE

Specifics about Sinoko clinic, Ndivisi, Bungoma County

HUMAN-CENTERED DESIGN

Explanation of the HCD framework.

HEAR PHASE

This is the hear phase.

CREATE PHASE

This is the create phase.

DELIVER PHASE

This is the deliver phase.

CREATE PHASE

These are the results of the create phase.

DELIVER PHASE

FRAME 1