

Progress Report

Modeling the mechanism of topoisomerase

Aukkawut Ammartayakun

June 23, 2022

Worcester Polytechnic Institute

1. Goals
2. Literature Review

Goals



Overall goal

- Understand the interaction of *TOP2* and the topology of DNA.

This week goal

- Continue to experiment on the minimum force required to reduce the linking number.

Literature Review

- Let say there is an exist of the energy-state mapping $E : \mathcal{S} \rightarrow \Omega$ for state domain \mathcal{S} .
- with our assumption last time, we can estimate the energy required to transform a state $s_1 \in \mathcal{S}$ to $s_2 \in \mathcal{S}$. Essentially it is $\Delta E = E(s_1) - E(s_2)$.
- Ziraldo *et al.* (2019) suggests that there is a dynamic that can identify the path given the state $(K, \Delta Lk)$ for knot type K .
 - The energy landscape is defined as $E(K, \Delta Lk) = -k_B T \ln(P(K, \Delta Lk))$
 - That is, in **static** system, we can estimate the work required for *TOP2*

How can we know $P(K, \Delta Lk)$?

From Bayes' law,

$$P(K, \Delta Lk) = P(K)P(\Delta Lk|K)$$

How can we know $P(K, \Delta Lk)$?

From Bayes' law,

$$P(K, \Delta Lk) = P(K)P(\Delta Lk|K)$$

$P(K)$ can be simulated with Monte Carlo method (i.e. generating the distribution for knot type) with the known data.

How can we know $P(K, \Delta Lk)$?

From Bayes' law,

$$P(K, \Delta Lk) = P(K)P(\Delta Lk|K)$$

$P(K)$ can be simulated with Monte Carlo method (i.e. generating the distribution for knot type) with the known data.

$P(\Delta Lk|K)$ can be estimated from convolution of torsional fluctuation (i.e. distribution of twist) along with the relationship $\Delta Tw = \Delta Lk - Wr$.

How can we know $P(K, \Delta Lk)$?

From Bayes' law,

$$P(K, \Delta Lk) = P(K)P(\Delta Lk|K)$$

$P(K)$ can be simulated with Monte Carlo method (i.e. generating the distribution for knot type) with the known data.

$P(\Delta Lk|K)$ can be estimated from convolution of torsional fluctuation (i.e. distribution of twist) along with the relationship $\Delta Tw = \Delta Lk - Wr$.

Podtelezhnikov *et al.* (1999) come up with the approximation for $P(\Delta Lk|K)$

$$P(\Delta Lk|K) \stackrel{\text{approx}}{\sim} \mathcal{N}(c_K, \sigma_K^2)$$

where $c_K = \mathbb{E}_K(Wr)$

Free energy minimization (con't)




Figure 1: Energy minimization pathway

Free energy minimization (con't)







Figure 2: Topological simplification pathway

-  PODTELEZHNIKOV, A. A., COZZARELLI, N. R., AND VOLOGODSKII, A. V.
Equilibrium distributions of topological states in circular dna: Interplay of supercoiling and knotting.
Proceedings of the National Academy of Sciences 96, 23 (1999), 12974–12979.
-  STOLZ, R., YOSHIDA, M., BRASHER, R., FLANNER, M., ISHIHARA, K., SHERRATT, D. J., SHIMOKAWA, K., AND VAZQUEZ, M.
Pathways of dna unlinking: A story of stepwise simplification.
Scientific Reports 7, 1 (Sep 2017), 12420.

-  ZIRALDO, R., HANKE, A., AND LEVENE, S. D.
Kinetic pathways of topology simplification by Type-II topoisomerases in knotted supercoiled DNA.
Nucleic Acids Research 47, 1 (11 2018), 69–84.