

# Homework 3

Aukkawut Ammartayakun  
CS 539 Machine Learning

Spring 2023

## Problem 1

Consider a hidden Markov model in which the emission densities are represented by a parametric model  $p(x|z, w)$ , such as a linear regression model or a neural network, in which  $w$  is a vector of adaptive parameters. Describe how the parameters  $w$  can be learned from data using maximum likelihood.

**Solution.** Given that  $w$  is the parameters of the parametric model, one can define the likelihood using the structure of HMM (from emission density, which we assume to be a parametric model) as product of the emission density and the transition density. Now, we can just use EM algorithm to find the maximum likelihood of  $w$ .

## Problem 2

Show that the finite sample estimator  $\hat{f}$  defined by (11.2)

$$\hat{f} = \frac{1}{L} \sum_{l=1}^L f(\mathbf{z}^{(l)})$$

has mean equal to  $\mathbb{E}[f]$  and variance given by (11.3)

$$\text{Var}[\hat{f}] = \frac{1}{L} \mathbb{E}[(f - \mathbb{E}[f])^2]$$

**Solution.** It is clear that,

$$\begin{aligned} \mathbb{E}[\hat{f}] &= \mathbb{E} \left[ \frac{1}{L} \sum_{l=1}^L f(\mathbf{z}^{(l)}) \right] \\ &= \frac{1}{L} \sum_{l=1}^L \mathbb{E}[f(\mathbf{z}^{(l)})] \\ &= \frac{1}{L} \sum_{l=1}^L \mathbb{E}[f] \\ &= \mathbb{E}[f] \end{aligned}$$

We can also show that

$$\begin{aligned} \text{Var}[\hat{f}] &= \mathbb{E}[(\hat{f} - \mathbb{E}[\hat{f}])^2] \\ &= \mathbb{E} \left[ \left( \frac{1}{L} \sum_{l=1}^L f(\mathbf{z}^{(l)}) - \mathbb{E}[f] \right)^2 \right] \end{aligned}$$

Since  $\hat{f}$  is an unbiased estimator of  $f$ , the summation terms after distributing the quadratic form, will be  $\frac{1}{L}\mathbb{E}[f^2]$  which then leads to the desired result.

### Problem 3

Suppose that  $z$  has a uniform distribution over the interval  $[0, 1]$ . Show that the variable  $y = b \tan z + c$  has a Cauchy distribution given by (11.16).

$$q(z) = \frac{k}{1 + (z - c)^2/b^2}$$

**Solution.** We will show that  $y$  has a Cauchy distribution. We know that the transformation of  $u(z)$  to  $q(y)$  can be done with the Jacobian determinant. We know that  $y = b \tan z + c$  or  $z = \arctan\left(\frac{y-c}{b}\right)$  and  $z \sim U(0, 1)$ . We can calculate the Jacobian determinant and transform that as

$$\begin{aligned} q(y) &= u(z) \left| \frac{dz}{dy} \right| \\ &= \left| \frac{dz}{dy} \right| \\ &= \frac{1}{b} \frac{1}{1 + \left(\frac{y-c}{b}\right)^2} \end{aligned}$$

Without loss of generality, we can assume that  $\frac{1}{b} = k$  and get what we desired.