# Grundlagen der Betriebssysteme

Tim Luchterhand, Paul Nykiel

21. April 2018

#### 1 Aufgabe

(a)

$$89/2 = 44 \text{ Rest } 1$$
  
 $44/2 = 22 \text{ Rest } 0$   
 $22/2 = 11 \text{ Rest } 0$   
 $11/2 = 5 \text{ Rest } 1$   
 $5/2 = 2 \text{ Rest } 1$   
 $2/2 = 1 \text{ Rest } 0$   
 $1/2 = 0 \text{ Rest } 1$   
 $\Rightarrow 89_{10} = 1011001_2$ 

(b) Zuerst Umrechnung ins Dezimalsystem, um Division zu erleichtern:

$$32_7 = 3 * 7^1 + 2 * 7^0 = 23_{10}$$
  
 $23/5 = 4 \text{ Rest } 3$   
 $4/5 = 0 \text{ Rest } 4$   
 $\Rightarrow 32_7 = 43_5$ 

(c) 
$$4360_{10} = 4096_{10} + 256_{10} + 8_{10} = 2^{12} + 2^8 + 2^3 = 1000100001000_2$$

(d)

$$\begin{array}{rll} & & & 1414215376_8 & & = \\ 001\ 100\ 001\ 100\ 010\ 001\ 101\ 011\ 111\ 1110_2 & = \\ & & 1100\ 0011\ 0001\ 0001\ 1010\ 1111\ 1110_2 & = \\ & & C\ 3\ 1\ 1\ A\ F\ E \end{array}$$

## 2 Aufgabe

(a)  $CAFFEE_{16} = 14\ 12\ 17\ 17\ 16\ 16_8 = 141217171616_8$ 

(b) 
$$3072 = 2048 + 1024 = 2^{11} + 2^{10} = 110000000000_2$$

(c)  $1724656_8 = 001\ 111\ 010\ 100\ 110\ 101\ 110_2 = 01111\ 01010\ 01101\ 01110_2 = F\ A\ D\ E_{32}$ 

(d) 
$$1316_2 = 1024 + 256 + 32 + 4 = 2^{10} + 2^8 + 2^5 + 2^2 = 10100100100_2$$

#### 3 Aufgabe

(a)

$$\begin{array}{r}
1011001_2 \\
+ 11000_2 \\
\hline
11 \\
----
=1110001_2
\end{array}$$

(b)

$$\begin{array}{r}
 101101_2 \\
 + 10111111_2 \\
 \hline
 11111111 \\
 \hline
 = 10001100_2
 \end{array}$$

(c)

$$\begin{array}{r}
 101011_2 \\
 + 110111_2 \\
 \hline
 1111111 \\
 \hline
 = 1100010_2
 \end{array}$$

### 4 Aufgabe

(a)

$$\begin{array}{rcl} 2018_{10} & = & 00000111 \ 11100010_2 \\ -2018_{10} & = & 11111000 \ 00011101_2 + 1 \\ & = & 11111000 \ 00011110_2 \end{array}$$

(b)

$$\begin{array}{rcl} 27346_{10} & = & 01101010 \ 11010011_2 \\ -27346_{10} & = & 10010101 \ 00101100_2 + 1 \\ & = & 1001010100101101_2 \end{array}$$

## 5 Aufgabe

(a)

$$\begin{array}{r}
10110_2 \cdot 111_2 \\
\hline
10110_2 \\
+ 10110_2 \\
+ 10110_2 \\
\hline
10011010_2
\end{array}$$

(b)

#### $10010010_2 \cdot 1001001_2$

 $\begin{array}{rrrr}
10010010_{2} \\
+ & 0_{2} \\
+ & 0_{2} \\
+ & 10010010_{2} \\
+ & 0_{2} \\
+ & 0_{2} \\
+ & 10010010_{2}
\end{array}$ 

 $10100110100010_2\\$ 

(c)

#### $10011110_2\cdot 10101_2$

 $\begin{array}{r}
 10011110_2 \\
 + 0_2 \\
 + 10011110_2 \\
 + 0_2 \\
 + 10011110_2
 \end{array}$ 

 $110011110110_2\\$