

Chapter 2 A Taste of Topology

Problem 6

Determine whether $d_x(p, q) = \sin |p - q|$ on $[0, \frac{\pi}{2})$ is a metric.

Proof: It is a metric. Positive definiteness follows from the fact that the absolute value function is a metric over \mathbb{R} , and sine being one-to-one over the range of possible functions. Symmetry follows for the same reason. The triangle inequality follows from sine being increasing and concave over $[0, \frac{\pi}{2})$.

Specifically, let $p, r \in [0, \frac{\pi}{2})$, and without loss of generality, let $p \leq r$. If $q = p$ or $q = r$, the triangle inequality is trivial.

Let $q \notin [p, r]$. If $q > r$, then $q - p > r - p$ implies

$$d_s(p, q) + d_s(q, r) \geq d_s(p, r) + 0 = d_s(p, r)$$

A similar result holds if $q < p$. If $q \in (p, r)$, imagine p, q , and r arranged on a line, with p at the origin. As x increases from q to r , the increase in sine is less than the corresponding increase from 0 to $r - q$, because sine is concave. Thus $\sin(r - p) < \sin(r - q) + \sin(q - p)$.

□

Problem 12

Let (p_n) be a sequence and $f : \mathbb{N} \rightarrow \mathbb{N}$ be a bijection. The sequence $(q_k)_{k \in \mathbb{N}}$ is a rearrangement of (p_n) with $q_k = p_{f(k)}$.

Part a

Are the limits of a sequence unaffected by rearrangement?

The limits of a sequence are unaffected by rearrangement.

Suppose that $(p_n) \rightarrow p$. Then for arbitrary $\epsilon > 0$, there exists an $n \in \mathbb{N}$ such that $i > n$ implies $d(p_i, p) < \epsilon$. This implies that there are at most finite elements of the sequence (p_n) such that $d(p_n, p) \geq \epsilon$. In the rearrangement (q_n) , let M be the smallest integer such that $f^{-1}(k) \leq n$. M exists because $\{1, 2, 3 \dots n\}$ is finite. Then for all $i > M$, $d(q_i, p) < \epsilon$, and thus $(q_i) \rightarrow p$.

On the other hand, suppose that (p_n) has no limit. Then for arbitrary p , there exists an $\epsilon > 0$ such that for all $n \in \mathbb{N}$, there exists $i > n$ such that $d(p_i, p) > \epsilon$. Letting $A(p)$ be the set of these points. A is infinite, because A being finite implies that A has a maximum element.

By contradiction, assume that that (q_n) has a limit of q . Then for all $\epsilon_q > 0$, there exists $n(\epsilon_q) \in \mathbb{N}$ such that $i > n(\epsilon_q)$ implies $d(q_i, q) < \epsilon_q$.

Let $\epsilon_q = \epsilon$. Then the set of points A_q such that $d(q_i, q) > \epsilon$ is a subset of $\{1, 2 \dots n(\epsilon)\}$, and is thus finite. But because (q_n) is a rearrangement of (p_n) , $A_q = A$. Thus A_q is both finite and infinite, which is a contradiction.

Part b

What if f is an injection?

At first glance, this question seems nonsensical, as for $n \in \mathbb{N}$ which are not in the preimage of $f(\mathbb{N})$, q_n is undefined. If we define such points to have $q_n = 0$, this question becomes intelligible.

Note that if A is an arbitrary set in \mathbb{N} , then $f(A)$ has the same cardinality as A , because the restriction of f on A is a bijection between the two.

The first result in Part a does not hold. Let $(p_n) = 1$, and let $f(x) = 2x$. The second result still holds, since the cardinality of the sets remains unchanged.

Part c

What if f is a surjection?

The first result in Part a holds. The cardinality of the image of a function must be less than or equal to the cardinality of the domain of the function. Letting $A = \{1, 2, \dots, n\}$ and $B = \{n+1, n+2, n+3, \dots\}$, $f(A)$ is finite. Since f is surjective, it covers \mathbb{N} , which is infinite. $A + B = \mathbb{N}$ which is the domain of f . If $f(B)$ is finite, then $f(A) + f(B) = f(A + B) = f(\mathbb{N})$ is finite, which contradicts the assumption that f is surjective. Therefore $f(B)$ is infinite, and the argument in Part a holds.

The second result in Part b does not hold. Let $(p_n) = \{0, 0, 0, 1, 0, 1, 0, 1, \dots\}$ and let $f(x) = 1$ if x is odd or equals 2, and $\frac{1}{2}x$ if x is even and greater than 3. f is clearly surjective, and $(q_n) = \{0, 1, 1, 1, 1, \dots\}$ clearly has a limit of 1.

Problem 13

Show that if $f : M \rightarrow N$ is a function such that (p_n) converging in M implies that $(f(p_n))$ converges in N , then f is continuous.

Note that this is almost the definition of convergence, except for the requirement that $(f(p_n)) \rightarrow f(p)$. Thus showing that f is continuous is equivalent to showing the above requirement.

Proof: Let $p \in M$ be arbitrary. Let (a_n) be the uniform sequence $a_n = p$ for all $n \in \mathbb{N}$. From the properties of f , $(f(a_n))$ converges in N , and it converges to $f(p)$. If (a_n) is the only sequence in M that converges to p for all p in M (such as when the discrete metric is used), then f satisfies the sequential convergence condition and is thus continuous.

Let (b_n) be an arbitrary sequence in M such that $b_n \rightarrow p$. Construct the sequence (c_n) such that $c_n = a_{\lceil n/2 \rceil}$ if n is odd and $c_n = b_{n/2}$ if n is even. (c_n) clearly converges to p .

By the convergence preservation condition of f , $(f(c_n))$ converges in N . The subsequence of $(f(c_n))$ consisting of the odd numbers converges to $f(p)$. All subsequences of a convergent sequence converge to the same limit as the main sequence, so $f(c_n) \rightarrow f(p)$, and since $(f(b_n))$ is a subsequence of $(f(c_n))$, $f(b_n) \rightarrow f(p)$. Thus f preserves sequential convergence, and is thus continuous.

□

Problem 14

Let $f : M \rightarrow N$ be a bijection from one metric space to another that preserves distance, i.e. for all $p, q \in M$

$$d_N(fp, fq) = d_M(p, q)$$

Then f is called an isometry from M to N , and M and N are said to be isometric, $M \equiv N$.

Part a

Prove that every isometry is continuous.

Proof: Let $(p_n) \in M$ be an arbitrary sequence such that $p_n \rightarrow p \in M$. Then for arbitrary $\epsilon > 0$, there exists an $A \in \mathbb{N}$ such that $a > A$ implies that $d_M(p_a, p) < \epsilon$. Moving to the sequence $(f(p_n))$, by the distance preservation property we have that for the same $\epsilon > 0$, for the same $a > A$, we have $d_N(f(p_a), f(p)) < \epsilon$. Thus $f(p_n) \rightarrow f(p) \in N$, so f preserves sequential convergence and is thus continuous.

□

Part b

Prove that every isometry is a homeomorphism.

Proof: Since f is a bijection, its inverse f^{-1} exists. Since we proved that f is continuous, if we prove that f^{-1} is continuous, then f is a homeomorphism between M and N , and thus M and N are homeomorphic.

Let $(p_n) \in N$ be arbitrary such that $p_n \rightarrow p \in N$. Then because f is a bijection, f^{-1} exists, and so $(f^{-1}(p_n))$ and $f^{-1}(p)$ are well defined.

Suppose f^{-1} is not continuous. Then $(f^{-1}(p_n))$ does not converge to $f^{-1}(p)$, and since f is continuous and the inverse of f^{-1} , this implies that $(f(f^{-1}(p_n))) = (p_n)$ does not converge to $f(f^{-1}(p)) = p$. But this contradicts the assumption that $p_n \rightarrow p$. Thus f^{-1} is continuous, and M and N are homeomorphic.

□

Part c

Prove that $[0, 1]$ is not isometric to $[0, 2]$.

Proof: Consider the set of pairs of points in $[0, 2]$ that are distance 1 from each other. Because of the total ordering that $[0, 2]$ inherits from \mathbb{R} , these pairs can be uniquely identified by their left endpoints. Letting A be the set of left endpoints of such pairs of points, A takes the form $A = \{a : [a, a + 1] \subset [0, 2]\} = [0, 1]$.

Let f be an isometry from $[0, 1]$ to $[0, 2]$. By the distance preservation condition of f , for all $a \in A$, the preimage of a under f is a point $b \in [0, 1]$ such that the interval $[b, b + 1] \subset [0, 1]$. However, the only interval of such form is $[0, 1]$. $f(0)$ can not correspond to $[0, 1]$, as this violates f being a function. Thus an isometry from $[0, 1]$ to $[0, 2]$ does not exist. \square

Problem 15

Prove that isometry is an equivalence relation.

Proof: Let M be isometric to N , and $f : M \rightarrow N$ be the isometry. For arbitrary $p, q \in N$, consider $f^{-1}(p), f^{-1}(q) \in M$. By the distance preserving condition of f , $d_M(f^{-1}(p), f^{-1}(q)) = d_N(f(f^{-1}(p)), f(f^{-1}(q))) = d_N(p, q)$. Thus f^{-1} is a bijection that preserves distances from N to M , and thus N is isometric to M .

Let M be a metric space and f be the identity function. f is clearly a bijection from M to M and distance-preserving, so f is an isometry. Therefore M is isometric to itself.

Let $f : M \rightarrow N$ and $g : N \rightarrow P$ be isometries. Consider their composition $H : M \rightarrow P = (g \circ f)(m)$. H is the composition of bijective functions, and thus is bijective itself, and it's clear that H preserves distances between M and P . Thus M is isometric to P . \square

Problem 16

Is the perimeter of a square isometric to the circle? Homeomorphic?

Assuming that the square and the circle are embedded in \mathbb{R}^2 with the usual metric, the two are not isometric. If the length diagonal of the square does not equal the diameter of the circle, the proof is immediate. If the length of the diagonal of the square equals the diameter of the circle, call their common distance d . There are only two pairs of points on the square with distance d from each other, while there are an infinite number of pairs on the circle distance d from each other. Therefore f can not map between them while remaining a function.

The two are homeomorphic, as one can be stretched into the other.

Problem 18

Is \mathbb{R} homeomorphic to \mathbb{Q} ?

No, as the two have different cardinalities, there can not be a bijection between them.

Problem 19

Is \mathbb{Q} homeomorphic to \mathbb{N} ?

No. Let $p \in \mathbb{N}$. All convergent sequences $p_n \rightarrow p$ in \mathbb{N} eventually end in repeating p . On the other hand, for $q \in \mathbb{Q}$, there exist sequences $(q_n), (r_n) \in \mathbb{Q}$ such that for all n , $q_n \neq r_n \neq q$.

Suppose that f is a homeomorphism from \mathbb{Q} to \mathbb{N} such that $f(q) = p \in \mathbb{N}$. Since f is bicontinuous, it preserves sequential convergence. Thus $(f(q_n))$ and $(f(r_n))$ both converge to p . Because of the nature of continuous sequences in \mathbb{N} , there exists an $N \in \mathbb{N}$ such that $n > N$ implies that $f(q_n) = f(r_n) = p$. But this implies that f is not injective, and thus not a bijection, which contradicts the assumption that f is a homeomorphism. Thus there exists no homeomorphism between \mathbb{Q} and \mathbb{N} .

Problem 20

What function is a homeomorphism from $(-1, 1)$ to \mathbb{R} ? Is every open interval homeomorphic to $(0, 1)$?

The function $\tan(\frac{\pi}{2}x)$ is a homeomorphism from $(-1, 1)$ to \mathbb{R} . It is bijective, continuous, and its inverse $\frac{2}{\pi} \tan^{-1}(x)$ is continuous.

Every open interval (a, b) is homeomorphic to $(0, 1)$. The function $\frac{x-a}{b-a}$ is bijective and bicontinuous.

Problem 22

If every closed and bounded subset of a metric space M is compact, does it follow that M is complete?

No. Let $M = \{x \in \mathbb{R} : x > 0\}$ with the usual metric. The closed and bounded subsets of M are compact, by the same reasoning as in \mathbb{R} . However, the Cauchy sequence $(p_n) = \frac{1}{n}$ has no limit in M .

Problem 24

For which intervals $[a, b]$ in \mathbb{R} is the intersection $[a, b] \cap \mathbb{Q}$ a clopen subset of the metric space \mathbb{Q} ?

The intervals where $a \leq b$ and a, b are irrational numbers. Since \mathbb{Q} is a metric subspace of \mathbb{R} , it inherits its open and closed sets from \mathbb{R} . Since all intervals of the form $[a, b]$ are closed in \mathbb{R} , their intersection with \mathbb{Q} is closed.

These intersections are also open. In \mathbb{R} , the interval (a, b) is open, and because a, b are irrational, $a, b \notin \mathbb{Q}$, so $[a, b] \cap \mathbb{Q} = (a, b) \cap \mathbb{Q}$. Thus $[a, b] \cap \mathbb{Q}$ is open.

The above does not hold if a or b is rational. Without any loss of generality, suppose a is rational. Then $a \in [a, b] \cap \mathbb{Q}$, and there is no open ball that contains a and is a subset of $[a, b] \cap \mathbb{Q}$. Thus $[a, b] \cap \mathbb{Q}$ is not open.

Problem 26

Prove that a set $U \subset M$ is open if and only if none of its points are the limits of its complement.

For the forward, suppose $x \in U$ is a limit point of U^C . Then $x \in \bar{U}^C$, and since for all sets $A \subset \bar{A}$, $x \in U^C$, which contradicts the assumption that $x \in U$. Thus U contains none of the limits of its complement.

For the reverse, suppose that U is not open. Then there exists a $x \in U$ such that for all $r > 0$, the open ball $B_U(x, r)$ is not fully contained in U . Thus, for $r = \frac{1}{n}$, we can choose a sequence $x_n \in B_U(x, r) \cup U^C$. $x_n \rightarrow x$ and $x_n \in U^C$ for all n , so x is a limit point of U^C . This contradicts the assumption that none of U 's points are the limit points of its complement, so U is open.

Problem 27

If $S, T \subset M$, a metric space, and $S \subset T$, prove that

Part a

$$\bar{S} \subset \bar{T}$$

Proof: Let $x \in \bar{S}$. Then there exists a sequence $(x_n) \in S$ such that $x_n \rightarrow x$. Since $S \subset T$, $(x_n) \in T$, so $x_n \rightarrow x$ implies that $x \in \bar{T}$. \square

Part b

$$\text{int}(S) \subset \text{int}(T)$$

Proof: $\text{int}(S) \subset S \subset T$. Because $\text{int}(S)$ is open and a subset of T , it must be contained in the largest open subset of T , which is $\text{int}(T)$. \square

Problem 28

A map $f : M \rightarrow N$ is open if U being an open subset of M implies that $f(U)$ is an open subset of N .

Part a

If f is open, is it continuous?

No. Let f be the floor function from \mathbb{R} to \mathbb{Z} . Every set in \mathbb{Z} is open, since singleton sets are open in \mathbb{Z} and the union of open sets is open, whether finite or infinite. However, the floor function is not continuous, as the example of $p_n = 1 - \frac{1}{n}$ shows.

Part b

If f is a homeomorphism, is it open?

Yes. Let $f : M \rightarrow N$. Let $A \subset M$ be an arbitrary open set, and $f(A) \subset N$ be its image. Because f is a bijection, it has a continuous inverse function f^{-1} . Since $f(A)$ is the preimage of A , $f^{-1}(f(A)) = A$ being open implies $f(A)$ is open. Thus open sets are mapped to open sets, and f is open.

Part c

If f is an open, continuous bijection, is it a homeomorphism?

Yes. By similar reasoning to Part b, let $A \subset M$ be an arbitrary open set. Because f is a bijection, f^{-1} exists, and A is the image of some set $f(A) \subset N$. Because f is open, $f(A)$ is open. Thus the preimage of open sets of f^{-1} is an open set, and f^{-1} is continuous. Since f is a bicontinuous bijection, it is a homeomorphism.

Part d

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous surjection, must it be open?

No. Let f be defined as $f(x) = x$ if $x \in \mathbb{Z}$. For $x \in \mathbb{R} - \mathbb{Z}$, define f as

$$f(x) = \begin{cases} x & x \in \mathbb{Z} \\ \lfloor x \rfloor + 2\text{Remainder}(x) & \text{Remainder}(x) \in (0, \frac{1}{2}) \\ \lceil x \rceil & \text{Remainder}(x) \in [\frac{1}{2}, 1) \end{cases}$$

where $\text{Remainder}(x)$ is the non-integer part of x . $f(x)$ is essentially a piecewise staircase with a slope of 2 on $[0, \frac{1}{2})$ pieces and a slope of 0 on $(\frac{1}{2}, 1)$ pieces. The function is continuous and surjective, but not open. For example, the open set $(\frac{1}{2}, \frac{3}{4})$ maps to the singleton 1, which is not open.

Part e

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous, open surjection, must it be a homeomorphism?

Yes. Note that if f is injective, then f is a bijection and Part c implies that it is a homeomorphism.

Suppose that f is not injective. Then there exists $a, b \in \mathbb{R}, a \neq b$ such that $f(a) = f(b)$. Without loss of generality, let $a < b$. Then by continuity, f achieves a maximum value on $[a, b]$.

If $f(a) = M = m$, then f is constant on $[a, b]$. This contradicts the assumption that f is open. For example, the image of the open ball $(a + \frac{b-a}{4}, b - \frac{b-a}{4})$ is a singleton, and thus not open.

If $f(a) < M$, then there exists $c \in (a, b)$ such that $f(c) = M$. Thus there exists $\delta = \min(\frac{c-a}{2}, \frac{b-a}{2}) > 0$. Consider the open ball $(c - \delta, c + \delta)$ and its image under f . Because f is bounded, the image is bounded, and because f is continuous, the image obtains all intermediate values. Thus $f((c - \delta, c + \delta))$ is an interval. Let x, y be the left and right endpoints of the interval, respectively.

Since $f(c)$ is the maximum of f , the right endpoint of the interval is closed. Thus $f((c - \delta, c + \delta))$ has the form $(x, y]$ or $[x, y]$, neither of which is open. Thus, the image of an open set under f is not open, which contradicts the assumption that f is open. A similar argument holds for the minimum. This, f is injective, bijective, and a homeomorphism.

Part f

What happens in Part e if \mathbb{R} is replaced by the unit circle S^1 ?

The result does not hold. Parameterize S^1 by the angle $\theta = \arctan(x)$, and let $f : S^1 \rightarrow S^1$ map a point x to the corresponding point at 2θ . f is obviously surjective, and it is continuous because it preserves sequential limits. It is open because open sets in S^1 consist of open line segments or their unions, and the image of line segments is either another union of line segments, or the entire metric space S^1 . However, f is not injective. Letting a be the point at $\theta = 0$ and b the point at $\theta = \pi$, $a \neq b$, but $f(a) = f(b)$.

Problem 30

Consider a two-point set $M = \{a, b\}$ whose topology consists of two sets, M and the empty set. Why does this topology not arise from a metric on M ?

Let d be a metric on M . By the properties of metrics, $d(a, b) = d(b, a) = c > 0$, and $d(a, a) = d(b, b) = 0$. The singleton sets a, b are open, since the open ball $B_d(a, c/2) = a$ and $B_d(b, c/2) = b$ are contained within themselves. However, the singleton sets are not contained in the topology.

Problem 31

Prove the following.

Part a

If U is an open subset of \mathbb{R} then it consists of countably many disjoint intervals $U = \sqcup U_i$. (Unbounded intervals $(-\infty, b)$, (a, ∞) , and $(-\infty, \infty)$ are allowed).

Lemma 1 *The bounded, connected sets on \mathbb{R} are intervals.*

Proof: *Let A be bounded and connected. By connectedness, it has the intermediate value property. By boundedness, it has a l.u.b. and g.l.b. Letting a be the g.l.b. and b be the l.u.b. and using the intermediate value property, $(a, b) \subset A$. By the l.u.b., $A \subset [a, b]$. Thus A is an interval. The reverse was shown in the book. \square*

Corollary 2 *The bounded, connected, open sets on \mathbb{R} are open intervals.*

Proof: The connected sets on \mathbb{R} are intervals, and the (topologically) open intervals are (colloquial) open intervals. \square

Lemma 3 Let A and B be overlapping bounded open intervals. Then $A \cup B$ is an open interval.

Proof: A and B are connected and share a common point, so $A \cup B$ is connected. Since $A \cup B$ is bounded, $A \cup B$ is an interval. Since $A \cup B$ is open, $A \cup B$ is an open interval. \square

Corollary 4 Let A_i be a collection of disjoint bounded open intervals of \mathbb{R} . Then there exists disjoint open intervals B_i such that $\cup A_i = \cup B_i$

Proof: If the A_i are disjoint, the proof is obvious. If not, then there exist i, j such that A_i and A_j overlap. By Lemma 3, $B_i = A_i \cup A_j$ is an open interval. Substitute A_i and A_j with B_i , and repeat while there are still overlapping intervals in A . \square

(I'm not sure that this proof is rigorous. It seems like I'm implicitly assuming that A_i is countable, but I don't know enough to be sure.)

Lemma 5 If A is an open subset in \mathbb{R} , then there exist bounded open intervals A_i such that $\cup A_i = A$.

Proof: Because A is open, for all $a \in A$, there exists $r_a > 0$ such that $(a - r_a, a + r_a) \subset A$. Do this for all points in A . \square

Lemma 6 Let $\epsilon > 0$. Let A_i be a collection of disjoint open intervals such that $\text{length}(A_i) > \epsilon$ for all i . Then A_i is countable.

Proof: Let B_i be the intervals of A_i dilated by a factor of $1/\epsilon$. That is, if $A_i = (x, y)$, $B_i = (x/\epsilon, y/\epsilon)$. Because all of the A_i s have length greater than $1/\epsilon$, the B_i have length greater than 1. Thus the B_i can be uniquely identified with the natural numbers by associating each B_i with the floor of its left endpoint. Since the natural numbers are countable, B_i is at most countable, and there exists a homeomorphism between the A_i and B_i , the A_i are countable. \square

Theorem 7 If A is an open subset of \mathbb{R} , then it consists of disjoint open intervals $A = \cup A_i$ (unbounded open intervals are acceptable).

Proof: If A is bounded, then by Lemma 5, A can be expressed as the union of open intervals, and by Corollary 4, A can be expressed as the disjoint union of bounded open intervals.

If A is unbounded and $A = \mathbb{R}$, the proof is trivial. Suppose A is unbounded in the positive direction. For simplicity, suppose that A is bounded in the negative direction. Since A does not equal \mathbb{R} , A^C is nonempty, and by A being unbounded in the positive direction, A^C is bounded above. Thus A^C has a least upper bound.

Denote the l.u.b. of A^C as a . By definition, $(a, \infty) \subset A$. Because A is open, A^C is closed, and since a is a limit point of A^C , $a \in A^C$.

Let $C = A - (a, \infty)$. For all $c \in C$, $c < a$. If $c = a$, then $C \subset A$ implies $c \in A$, which contradicts that $a \in A^C$. If $c > a$, then $c \in C$ and $c \in (a, \infty)$, which is a contradiction because by construction C and (a, ∞) are disjoint. Thus C is bounded above. C is bounded below because $C \subset A$ and A is bounded below. Thus C is bounded.

C is open. $C^C = A^C \cup (a, \infty) = A^C \cup [a, \infty)$ because $a \in A^C$. Thus C^C is the finite union of closed sets, and is closed.

Thus C is a bounded open subset of \mathbb{R} , and can be expressed as the disjoint union of bounded open intervals C_i . Adding back in (a, ∞) gives A as the disjoint union of open intervals A_i . The cases where A is bounded above and unbounded below, and case when A is unbounded in both directions, follow similarly. \square

Lemma 8 Let A_i be a collection of disjoint intervals in \mathbb{R} . Then there is at most one unbounded interval in the positive direction, and at most one (possibly the same) unbounded interval in the negative direction.

Proof: Let A_m and A_n be distinct intervals in A that are unbounded above. Because they are intervals, they are non-empty, so there exist $m \in A_m$ and $n \in A_n$. By the trichotomy property on \mathbb{R} , either $m = n$, $m > n$, or $n > m$. In any of the cases, one of m or n is in both sets, violating the assumption that A_i is disjoint. \square

Theorem 9 If A_i is a collection of disjoint open intervals in \mathbb{R} , then A_i is countable.

Proof: Create subcollections B_i , where $i = 0, 1, 2 \dots \mathbb{N}$. Let B_0 consist of the unbounded intervals of A_i , of which there are at most 2, B_1 the intervals of A_i with finite length greater than 1, and B_i the intervals of A_i with length greater than $1/i$ and less than or equal to $1/(i-1)$. By Lemma 6, B_i contains at most countable many intervals. Continuing this process for all \mathbb{N} , we see that the B_i s contains all the intervals of A_i . Because B_i contains a countable number of sets, each with a countable number of elements, B_i , and thus A_i is countable. \square

Corollary 10 If U is an open subset of \mathbb{R} then it consists of countably many disjoint intervals $U = \sqcup U_i$. (Unbounded intervals $(-\infty, b)$, (a, ∞) , and $(-\infty, \infty)$ are allowed).

Proof: Follows immediately from Theorems 7 and 9. \square

Part b

Prove that the intervals U_i are uniquely determined by U . In other words, there is only one way to express U as a disjoint union of open intervals.

I will assume that the U_i have to be open, otherwise the statement is false. For example, $(0, 1)$ can also be written as $(0, .5] \cup (.5, 1)$.

Lemma 11 *Let A_i be a disjoint union of open intervals and $A = \sqcup A_i$. Let A_k be a particular interval with left and right endpoints b and c , respectively. Then $b, c \in A^C$.*

Proof: Suppose $b \in A$. Since $b \in A_k^C$, b must be in a different open interval. Denote the open interval A_j . Since A_j is open, there exists $r > 0$ such that $(b - r, b + r) \subset A_j$. Consider $b - r/2$. $b - r/2 < b$, so $b - r/2 \in A_k$. $b - r/2 \in (b - r, b + r) \subset A_j$, so $b - r/2 \in A_j$. Thus A_k and A_j intersect, which contradicts the assumption that the A_i are disjoint. Thus $b \in A^C$. The exact same argument holds for c . \square

Now to prove the theorem. Let A be an open subset and B_i and C_j be disjoint collections of open intervals such that $A = \sqcup B_i = \sqcup C_j$. If A is unbounded in both directions, the proof is trivial.

Consider an arbitrary point $a \in A$. Because the B_i 's are disjoint and the C_j 's are disjoint, there exist unique B_a and C_a such that $a \in B_a$ and $a \in C_a$.

Letting x be the right endpoint of B_a and y the right endpoint of C_a , $x = y$. Suppose it's not. Without loss of generality, let $x < y$. Then $x \in A$ because $x \in C_a$, but $x \in A^C$ because x is the right endpoint of $B_a \subset A$, and so $x \in B_a$. Repeating this for $x > y$ shows that $x = y$.

A similar argument holds for the left endpoint. Repeat this argument for all points in A to show that for all $a \in A$, the endpoints of B_a equal the endpoints of C_a . Thus, $B_a = C_a$ and the two collections B and C are equal.

Part c

If $U, V \subset \mathbb{R}$ are both open, so $U = \sqcup U_i$ and $V = \sqcup V_j$ where U_i and V_j are open intervals, show that U and V are homeomorphic if and only if there are equally many U_i and V_j .

Proof: Let m be the number of element in U_i , and n be the number of elements in V_j . Note that m, n may be infinity.

For the forward, suppose there are not equally many U_i and V_j . Without loss of generality, let $m < n$. Let f be a homeomorphism between U_i and V_j . For arbitrary $k \in 1, 2 \dots m$, consider the restriction of f on U_k . Because U_k is an interval, it is connected, and so its image is connected. Repeating this for all U_i shows that V is the union of at most m connected sets. However, V consists of n connected intervals, so V can not be the image of U . Thus the homeomorphism f does not exist. A similar argument in reverse if $m > n$.

For the reverse, construct a homeomorphism as follows. By Part a above, U_i and V_j are countable. Label the sets. Let the homeomorphism be f such that each U_k is shifted and dilated so that $f(U_k) = V_k$. f is trivially a bijection. Since on each segment, f is a non-singular linear transformation, f and f^{-1} are continuous. \square

Problem 33

Part a

Find a metric space in which the boundary of $M_r p$ is not equal to the sphere of radius r at p , $\partial(M_r p) \neq \{x \in M : d(x, p) = r\}$.

Consider the discrete metric on a space with at least two elements. Note that since all sets under the discrete metric are clopen, the boundary of all sets is the null set. However, the sphere of radius 0 centered at p is p itself, and the sphere of radius 1 centered at p is the entire metric space, neither of which are the null set.

Part b

Need the boundary be contained in the sphere?

Yes. Suppose not. For an arbitrary set A in a metric space M , $\partial A \subset \bar{A}$. Thus the boundary not being contained in the sphere implies that $CB_r p = \{x \in M : d(x, p) \leq r\}$ is a proper subset of $\overline{M_r p}$, so that the boundary can potentially be outside of $CB_r p$. Because $\overline{M_r p}$ is the smallest closed set that contains $M_r p$ and $M_r p \subset CB_r p$, $CB_r p$ being a proper subset of $\overline{M_r p}$ implies that $CB_r p$ is not closed. This implies that $CB_r p^C = \{x \in M : d(x, p) > r\}$ is not open. If $CB_r p^C$ is the null set, this is a contradiction.

Otherwise, for all $x \in CB_r p^C$, let $d(x, p) = y > r$. Then the open ball $M_{\frac{y-r}{2}} x \subset CB_r p^C$ by the triangle inequality, and $CB_r p^C$ is open, implying $CB_r p$ is closed. Thus there is a closed set containing $M_r p$ that is strictly smaller than the closure of $M_r p$, a contradiction.

Problem 34

Assume that N is a metric subspace of M and is also a closed subset of M . Show that $L \subset N$ is closed in N if and only if it is closed in M . Similarly, if N is a metric subspace of M and also is an open subset of M then $U \subset N$ is open in N if and only if it is open in M .

Proof: For the closed version, we start with the forward. Because L is closed in N , by the Inheritance Principle, $L = N \cap P$, where P is a closed set in M . Viewing N and P as subsets of M , since N is closed by assumption, $L = N \cap P$ is closed in M . Conversely, let L be closed in M . Then by the Inheritance Principle, $L \cap N$ is a closed set in N . Since $L \subset N$, $L \cap N = L$ is closed in N . The open version is essentially the same, with the word 'open' replacing the word 'closed.' \square

Problem 35

Prove that S clusters at p if and only if for each $r > 0$ there is a point $q \in M_r p \cap S$ such that $q \neq p$.

This should be a direct consequence of Theorem 52, that p being a cluster point of S is characterized by each neighborhood of p containing at least one point of S other than p .

Problem 36

Construct a set with exactly three cluster points.

In \mathbb{R}^2 , let $A_i = \{(x, y) \in \mathbb{R}^2 | (x, y) = (i, 1/n) \text{ for some } n \in \mathbb{N}\}$. Let $A = A_1 \cup A_2 \cup A_3$. The three cluster points are $\{(0, 0), (1, 0), (2, 0)\}$.

Problem 37

Construct a function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is continuous only at points of \mathbb{Z} .

First we consider the function which is x on the rationals and $-x$ on the irrationals. It is continuous only at $x = 0$. Take miniature copies of this function, centered at the integers, all only on the interval $(n - 1/2, n + 1/2]$. For all integers z , $f(z) = 0$. The function is continuous at the integers, and nowhere else.

Problem 38

Let X, Y be metric spaces with metrics d_X, d_Y , and let $M = X \times Y$ be their Cartesian product. Prove that the three natural metrics d_E, d_{max} , and d_{sum} on M are actually metrics.

Symmetry is obvious. The positive definiteness of d_{max} follows from the positive definiteness of the d_X and d_Y . The positive definiteness of d_{sum} is similar, and the positive definiteness of d_E follows because the square root function only has a root at 0.

For the triangle inequality on d_{sum} ,

$$\begin{aligned} d_{sum}(p, p^*) + d_{sum}(p^*, p') &= d_X(x, x^*) + d_X(x^*, x') + d_Y(y, y^*) + d_Y(y^*, y') \\ &> d_X(x, x') + d_Y(y, y') = d_{sum}(p, p') \end{aligned}$$

by the triangle inequality on d_X and d_Y . For d_{max} , $d_{max}(p, p') = \max\{d_X(x, x'), d_Y(y, y')\}$. Without loss of generality, suppose that $d_X(x, x') \geq d_Y(y, y')$. Then

$$\begin{aligned} d_{max}(p, p^*) + d_{max}(p^*, p') &= \max\{d_X(x, x^*), d_Y(y, y^*)\} + \max\{d_X(x^*, x'), d_Y(y, y')\} \\ &\geq d_X(x, x^*) + d_X(x^*, x') \geq d_X(x, x') = d_{max}(p, p') \end{aligned}$$

The argument if $d_X(x, x') < d_Y(y, y')$ is similar. For the triangle inequality on d_E ,

Problem 40

Let M be a metric space with metric d . Prove that the following are equivalent.

1. M is homeomorphic to M equipped with the discrete metric.
2. Every function $f : M \rightarrow M$ is continuous.
3. Every bijection $g : M \rightarrow M$ is a homeomorphism.
4. M has no cluster points.
5. Every subset of M is clopen.
6. Every compact subset of M is finite.

If written with no other qualifiers, M is equipped with its metric d .

For $1 \rightarrow 2$, for any arbitrary set equipped with the discrete metric, if p_n is a sequence such that $p_n \rightarrow p$, then the tail of $p_n = p$. Because $M_{discrete}$ is homeomorphic to M_d , this is true on M_d . Thus every convergent sequence has a constant tail, which is trivially preserved under an arbitrary function. Thus arbitrary functions are continuous.

Thus for any function $f : M_1 \rightarrow M_2$, sequential limits are trivially preserved, so f is continuous.

For $2 \rightarrow 3$, this follows because g^{-1} exists due to g being a bijection, and $g : M \rightarrow M$ and $g^{-1} : M \rightarrow M$ are both continuous.

For $3 \rightarrow 4$, let $p \in M$ be a cluster point of M . Because p is a cluster point, there exists a sequence of distinct points $p_n \in M$ such that $p_n \rightarrow p$. Let $g : M \rightarrow M$ be a function that swaps points p and p_1 , and is the identity function everywhere else. g is trivially a bijection, and thus it is a homeomorphism, and thus g is continuous and preserves sequential limits. However, $g(p_n) \rightarrow p$, but $g(p) = p_1 \neq p$, contradicting sequential limit preservation. Thus M has no cluster points.

For $4 \rightarrow 5$, because M has no cluster points, for all $p \in M$, there exists some $r > 0$ such that the neighborhood of M about r contains only p . Thus the open ball with radius r is contained in p , so all singleton sets are open. The arbitrary union of open sets is open, so all subsets of M are open. By taking compliments, all subsets of M are closed, and thus clopen.

For $5 \rightarrow 6$, let A be an infinite, compact subset of M , and A_n be an open covering of singletons. Then by compactness, there exists a finite subcover A_{n_k} of A . But it's clear that a finite union of singletons can not cover an infinite set. Thus A can not be infinite.

For $6 \rightarrow 1$, if d is the discrete metric, the proof is trivial. Otherwise, let d be the metric on M . I will show that M has no cluster points under metric d . From above, that means that all singletons are clopen in M with d , thus meaning that all subsets are clopen in M with d . Then the identity function between M with d and M with the discrete metric is trivially a homeomorphism.

Let p be a cluster point of M . Then for every open neighborhood of p , there exists a point in M that is not p . Consider the open neighborhoods $C_n = B_{r=1/n}(p)$ for all $n \in \mathbb{Z}^+$. Then for all n , there exists x_n such that $x_n \in C_n$.

Consider the set $C = (\cup_{n=1}^{\infty} x_n) \cup p$, and let A be an open cover of C . Because A is an open cover, there exists $A_p \in A$ such that $p \in A_p$. Since A_p is open, there exists a neighborhood with $r > 0$ centered at p such that $B_r(p) \subset A_p$. By the construction of C , $B_r(p)$ covers all of the x_n with $n > 1/r$. Thus the finite subcover $A_1 \cup A_2 \dots A_p$ covers A , making A compact. But this contradicts the assumption that all compact subsets of M are finite. Thus M has no cluster points.

Problem 41

Let $\|\cdot\|$ be a norm on \mathbb{R}^m , and let $B = \{x \in \mathbb{R}^m : \|x\| \leq 1\}$. Prove that B is compact.

Let $\|\cdot\|_E$ be the Euclidean norm on \mathbb{R}^m . If $\|\cdot\|$ equals the Euclidean norm, this is trivial.

We first want to show that the identity transformation from $(\mathbb{R}^m, \|\cdot\|_E)$ to $(\mathbb{R}^m, \|\cdot\|)$ is continuous. Due to the translation invariance of norms, if we show that the identity transformation is continuous at the origin, we have shown it for all \mathbb{R}^m . We begin with some lemmas. Let $\epsilon > 0$, and $C = \{x \in \mathbb{R}^m : \|x\| = \epsilon\}$.

Lemma 12 *The identity function from $(\mathbb{R}^m, \|\cdot\|_E)$ to $(\mathbb{R}^m, \|\cdot\|)$ is continuous.*

Proof: *All norms on a finite-dimensional vector space are equivalent, which means that open sets under $\|\cdot\|$ are open sets under $\|\cdot\|_E$, and vice versa. Thus the preimage of open sets under the identity transformation is trivially open. \square*

Corollary 13 *If $A \subset \mathbb{R}^m$ is compact with respect to the Euclidean norm, then it is compact with respect to $\|\cdot\|$.*

Proof: *The identity function is continuous, so $A \subset (\mathbb{R}^m, \|\cdot\|)$ is the image of a compact with respect to the Euclidean norm. By the theorems in the book, this implies that A is compact with respect to the $\|\cdot\|$ norm. \square*

Thus, if we can prove that B is compact with respect to the Euclidean norm, it is compact with respect to the norm $\|\cdot\|$. By the Heine-Borel theorem, this is equivalent to B being closed and bounded under the Euclidean norm.

For being closed, norms are continuous functions from \mathbb{R}^m to \mathbb{R} . It's clear that B is the preimage of the closed set $[0, 1] \subset \mathbb{R}$, implying that B is closed with respect to $\|\cdot\|$. Since the identity transformation from the Euclidean norm to the $\|\cdot\|$ norm is continuous, this implies that B is closed with respect to the Euclidean norm.

For boundedness, let $A = \{x \in \mathbb{R}^m : \|x\|_E = 1\}$ be the surface of the unit ball in \mathbb{R}^m under the Euclidean norm. A is clearly closed and bounded, so because it is a subset of \mathbb{R}^m , it is a compact. Since all norms are continuous, $\|\cdot\| : \mathbb{R}^m \rightarrow \mathbb{R}$ is a continuous function, and so the image of A under $\|\cdot\|$ obtains maximum and minimum values. Let $y = \min \|A\|$ be the minimum

value. Because norms are positive definite, the zero vector is not an element of A , so $y > 0$.

Define $B' = \{x \in \mathbb{R}^m : \|x\| \leq y\}$, and A' be the unit ball in \mathbb{R}^m . By construction, $B' \subset A'$ and is thus bounded under the Euclidean metric. Dilate B' and A' by a factor of $1/y$. By the norm property that $\|\alpha x\| = |\alpha|\|x\|$, after dilation, B' maps to B , and A' maps to a ball with radius $1/x$ centered at the origin. It's clear that B is a subset of this ball, thus making B bounded under the Euclidean metric.

Problem 43

Assume that the Cartesian product of two nonempty sets $A \subset M$ and $B \subset N$ is compact in $M \times N$. Prove that A and B are compact.

Proof: Let $(a_n) \in M$ and $(b_n) \in N$ be arbitrary sequences. Consider their Cartesian product $(a, b)_n \in M \times N$. By compactness of $M \times N$, there exists a convergent subsequence $(a, b)_{n_k} \rightarrow (a, b)$. Since all metrics on a Cartesian product are equivalent, let's use the sum metric, d_{sum} . Then for all $\epsilon > 0$ and large n , $d_{sum}((a, b)_{n_k}, (a, b)) = d_M(a_{n_k}, a) + d_N(b_{n_k}, b) < \epsilon$. Since distances are nonnegative, this implies that for large n $d_M(a_{n_k}, a) + d_N(b_{n_k}, b) < \epsilon$ and $d_N(b_{n_k}, b) < \epsilon$. Thus (a_n) and (b_n) have convergent subsequences, indicating that M and N are compact. \square

Problem 44

Consider a function $f : M \rightarrow \mathbb{R}$. Its graph is the set

$$G(f) = \{(p, y) \in M \times \mathbb{R} : y = f(p)\}$$

(Note the notation is my own invention; I have no idea if it's standard or not).

Part a

Prove that if f is continuous then its graph is closed as a subset of $M \times \mathbb{R}$.

Proof: By the theorems in the book, the metrics on a Cartesian product are equivalent. I will use d_{sum} for simplicity. Let d_M be the metric on M , and d_E be the Euclidean metric on \mathbb{R} .

Let $((a, f(a))_n) \in G(f)$ be a convergent sequence such that $(a, f(a))_n \rightarrow (a, b) \in M \times \mathbb{R}$. Because of convergence, this implies that for all $\epsilon > 0$, for large n , $d_{sum}((a, f(a))_n, (a, b)) = d_M(a_n, a) + d_E(f(a_n), b) < \epsilon$. Note that this implies that $a_n \rightarrow a$ and $f(a_n) \rightarrow b$. By continuity, $a_n \rightarrow a$ implies that $f(a_n) \rightarrow f(a)$. Since limits are unique, this implies $f(a) = b$, and so $(a, b) = (a, f(a)) \in M \times \mathbb{R}$. Thus $M \times \mathbb{R}$ is closed. \square

Part b

Prove that if f is continuous and M is compact then its graph is compact.

Proof: Because f is continuous and M is compact, $f(M) \subset \mathbb{R}$ is a compact. Therefore $G(f) = M \times f(M)$ is the Cartesian product of two compacts, and thus compact. \square

Part c

Prove that if the graph of f is compact then f is continuous.

If M is empty, then f is trivially continuous. If M is nonempty but p is an isolated point of M , then f is trivially continuous at p .

Otherwise, let p be a cluster point of M and suppose f is discontinuous at p . Then there exists $\epsilon > 0$ such that for all $\delta > 0$, there exists a $x \neq p \in M$: $d_M(x, p) < \delta$ such that $|f(p) - f(x)| > \epsilon$.

Therefore for all $n \in \mathbb{N}$, there exists a point $a_n \neq p \in M$ such that $d_M(a_n, p) < \frac{1}{n}$ and $|f(a_n) - f(p)| > \epsilon$.

Consider the sequence $(a_n) \in M$, and its associated sequence $((a_n, f(a_n))) \in M \times \mathbb{R}$.

Problem 45

Draw a Cantor set C on the circle and consider the set A of all chords between points of C .

Without loss of generality, I will assume the circle has circumference 1. I will arbitrarily choose a point p to be 0 and count off distances going counter-clockwise.

Part a

Prove that A is compact.

For an interval I_a , denote the left and right endpoints of I_a as A_L and A_R , respectively.

Lemma 14 *Let I_A and I_B be intervals in C_n , the n th step of the Cantor set. Without loss of generality, let $A_R < B_L$. Then the set of all chords between the intervals is the shape bounded by the curve $A_L A_R$, the line segment $\overline{A_R B_L}$, the curve $B_L B_R$, and the line segment $\overline{A_R B_L}$. Denote this shape S_{AB} .*

Proof: All chords must have one of their endpoints on I_A and the other on I_B . Since the intervals are disjoint, mental playing around with the picture will show that the curve draws the desired shape. \square

Lemma 15 *Let I_A and I_B be intervals as defined in 14, and let I_C, I_D be subintervals such that $I_C, I_D \subset I_A \cup I_B$. Then $S_{CD} \subset S_{AB}$.*

Proof: For $p \in S_{CD}$, p is on a chord with one endpoint on I_C and the other on I_D . By subintervals, the endpoints are in $I_A \cup I_B$, so the chord that connects those endpoints is contained in S_{AB} . Thus $p \in S_{AB}$. \square

Corollary 16 S_{AB} as defined in Lemma 14 is compact.

Proof: The shape is closed and bounded in \mathbb{R}^2 . \square

Lemma 17 Denote A_n the set of all chords that are drawn between intervals in C_n . Then A_n is compact.

Proof: $A_n = \cup_{i,j=1}^n S_{ij}$. In other words, A_n is the union of all S_{ij} , where i, j are two intervals in C_n . A_n is bounded, and because it is the union of finite closed sets, A_n is closed. Since $A_n \subset \mathbb{R}^2$, A_n is compact. \square

Lemma 18 A_{n+1} is a subset of A_n .

Proof: Let $p \in A_{n+1}$. Then p is in an S between two intervals of C_{n+1} , which are subintervals of an interval or intervals in C_n . By Lemma 15, p is in the corresponding S in C_n , so $p \in A_n$. \square

Lemma 19 A_n is nonempty for all n .

Proof: $1/3, 2/3$ are always points in A_n , and their chord is nonempty. \square

Lemma 20 A is compact.

Proof: $A = \cap_{i=1}^{\infty} A_i$. By Lemmas 17, 18, and 19, A is the intersection of a nested series of nonempty compacts, and is thus compact. \square

Problem 47

Suppose $A, B \subset \mathbb{R}^2$.

Part a

If A and B are homeomorphic, are their complements homeomorphic? No. Let $A = (0, 1)$ and B be the x -axis. $(0, 1)$ is homeomorphic to \mathbb{R} , so A is homeomorphic to B . However, A^C is connected while B^C is not.

Part b

What if A and B are compact?