

## 6.6 INTEGRATION AND DIFFERENTIATION

**Problem 6.6:1.** On  $[a, b]$ , let  $\alpha$  be a strictly increasing function and  $f$  a continuous function, and for  $x \in [a, b]$  define  $F(x) = \int_a^x f(t)d\alpha(t)$ . Show that for all  $x \in [a, b]$ ,  $\frac{dF(x)}{d\alpha(x)} = f(x)$ , where the left-hand side is defined as  $\lim_{t \rightarrow x} \frac{F(x) - F(t)}{\alpha(x) - \alpha(t)}$ , and the equality includes the assertion that the limit exists.

*Proof.* First, we note that because  $f \in \mathfrak{R}(\alpha)$  on  $[a, b]$

$$F(x) - F(t) = \int_a^x f(s)d\alpha(s) - \int_a^t f(s)d\alpha(s) = \int_t^x f(s)d\alpha(s)$$

For all partitions  $P$ ,

$$\int_t^x f(s)d\alpha(s) \leq \overline{\int_t^x f(s)d\alpha(s)} = \sum_{x_i \in P} M_i \Delta\alpha_i \leq \left( \sup_{s \in [x, t]} f(s) \right) \sum_{x_i \in P} \Delta\alpha_i = \left( \sup_{s \in [x, t]} f(s) \right) (\alpha(x) - \alpha(t))$$

Since  $\alpha(x)$  is strictly increasing,  $\alpha(x) - \alpha(t) \neq 0$  when  $x \neq t$ , so

$$\frac{F(x) - F(t)}{\alpha(x) - \alpha(t)} \leq \sup_{s \in [x, t]} f(s)$$

Taking the limit as  $x$  approaches  $t$  on both sides and using the continuity of  $f$  gives

$$\lim_{t \rightarrow x} \frac{F(x) - F(t)}{\alpha(x) - \alpha(t)} \leq f(x)$$

The analogous result for the lower integral and the Squeeze Theorem complete the proof. □

**Problem 6.6:2.**

(a). Show that if  $f$  is continuous, then

$$\int_{t=1}^b \left( \int_{s=a}^t f(s)ds \right) dt = \int_{t=a}^b (b-t)f(t)dt$$

*Proof.* Let  $x \in [a, b]$ . Define  $P(x) = \int_{t=a}^x \left( \int_{s=a}^t f(s)ds \right) dt$  and  $Q(x) = \int_{t=a}^x (x-t)f(t)dt$ .

$f(t)$  being continuous on  $[a, b]$  implies that it is Riemann-integrable. This implies that  $f^*(t) = \int_{s=a}^t f(s)ds$  is continuous, and that  $P(x)$  is continuous and differentiable. Similarly,  $(b-t)f(t)$  is continuous on  $[a, b]$ , so  $Q(x)$  is continuous and differentiable.

By the Fundamental Theorem of Calculus,

$$P'(x) = \int_{s=a}^x f(s)ds$$

For  $Q(x)$ , since  $t$  and  $tf(t)$  are Riemann-integrable,

$$Q(x) = x \int_{t=a}^x f(t)dt - \int_{t=a}^x tf(t)dt$$

$x$  is trivially differentiable. Since  $t$  and  $tf(t)$  are continuous,

$$Q'(x) = \int_{t=a}^x f(t)dt + xf(x) - xf(x) = \int_{t=a}^x f(t)dt$$

Thus,  $P'(x) = Q'(x)$ . Integrating both sides from  $a$  to  $c$ , then setting  $c = b$ , produces the desired result.  $\square$

(c). Show that the result of Part (a) continues to hold if  $f$  is merely assumed Riemann-integrable, but not necessarily continuous.

*Proof.*  $P(x)$  has the same derivative as in Part (a), as the derivation only assumed that  $P(x)$  is Riemann-integrable. Similarly, for  $x_0 \in [a, b]$  where  $f(x_0)$  is continuous, the above derivations hold for  $Q(x)$ .

Let  $x_0$  be a point where  $f(x_0)$  is discontinuous. First, we will prove two lemmas.

**Lemma 0.0.1.** *If  $f(x)$  is bounded, then  $(x - x_0)f(x)$  is continuous at  $x_0$ .*

*Proof.* Let  $M = \sup |f(x)|$ . Then  $(x - x_0)f(x) \leq |(x - x_0)f(x)| \leq |x - x_0|M$ , which can be made arbitrarily small.  $\square$

**Lemma 0.0.2.** *If  $f(x)$  is continuous, then  $(x - x_0)f(x)$  is differentiable at  $x_0$  with derivative  $f(x_0)$ .*

*Proof.* By the definition of differentiability,

$$\lim_{x \rightarrow x_0} \frac{(x - x_0)f(x) - (x_0 - x_0)f(x_0)}{x - x_0} = \lim_{x \rightarrow x_0} f(x) = f(x_0)$$

by continuity.  $\square$

We can rewrite  $Q(x)$  as

$$Q(x) = \int_{t=a}^x ((x - x_0) + (x_0 - t))f(t)dt = (x - x_0) \int_{t=a}^x f(t)dt + \int_{t=a}^x (x_0 - t)f(t)dt$$

because the sub-functions are trivially Riemann-integrable.  $\int_{t=a}^x f(t)$  is a continuous function, so by Lemma 0.0.2  $(x - x_0) \int_{t=a}^x f(t)dt$  is differentiable at  $x = x_0$  with derivative  $\int_{t=a}^{x_0} f(t)dt$ . Similarly,  $f(t)$  is bounded because it is Riemann-integrable, so by Lemma 0.0.1  $(x_0 - t)f(t)$  is continuous at  $t = x_0$ . Therefore  $\int_{t=a}^x (x_0 - t)f(t)dt$  is differentiable at  $x = x_0$ , with derivative 0.

Therefore,  $Q(x)$  is differentiable at  $x = x_0$ , and  $Q'(x_0) = \int_{t=a}^{x_0} f(t)dt$ . The proof then follows using the same logic as in Part (a).  $\square$