

Dependency Discovery via Multiscale Generalized Correlation

Cencheng Shen

Johns Hopkins University

*Collaborators: Joshua T. Vogelstein, Carey E. Priebe, Shangsi Wang, Youjin Lee,
Mauro Maggioni, Qing Wang, Alex Badea.*

Acknowledgment: NSF DMS, DARPA SIMPLEX.

1/23/2018

Set-up

Set-up

Given paired data $(\mathcal{X}_n, \mathcal{Y}_n) = \{(x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}^q, \text{ for } i = 1, \dots, n\}$,

Set-up

Given paired data $(\mathcal{X}_n, \mathcal{Y}_n) = \{(x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}^q, \text{ for } i = 1, \dots, n\}$,

- Are they related?

Set-up

Given paired data $(\mathcal{X}_n, \mathcal{Y}_n) = \{(x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}^q, \text{ for } i = 1, \dots, n\}$,

- Are they related?
- How are they related?

Set-up

Given paired data $(\mathcal{X}_n, \mathcal{Y}_n) = \{(x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}^q, \text{ for } i = 1, \dots, n\}$,

- Are they related?
- How are they related?

Set-up

Given paired data $(\mathcal{X}_n, \mathcal{Y}_n) = \{(x_i, y_i) \in \mathbb{R}^p \times \mathbb{R}^q, \text{ for } i = 1, \dots, n\}$,

- Are they related?
- How are they related?

X	Y
brain connectivity	creativity / personality
brain shape	health
gene / protein	cancer
social networks	attributes
anything	anything else

Formal Definition of Independence Testing

Formal Definition of Independence Testing

$$(x_i, y_i) \stackrel{i.i.d.}{\sim} F_{XY}, \quad i = 1, \dots, n$$

$$H_0 : F_{XY} = F_X F_Y,$$

$$H_A : F_{XY} \neq F_X F_Y.$$

Formal Definition of Independence Testing

$$(x_i, y_i) \stackrel{i.i.d.}{\sim} F_{XY}, \quad i = 1, \dots, n$$

$$H_0 : F_{XY} = F_X F_Y,$$

$$H_A : F_{XY} \neq F_X F_Y.$$

A test is universally consistent if its power converges to 1 as $n \rightarrow \infty$ against any dependent F_{XY} .

Formal Definition of Independence Testing

$$(x_i, y_i) \stackrel{i.i.d.}{\sim} F_{XY}, \quad i = 1, \dots, n$$


$$H_0 : F_{XY} = F_X F_Y,$$

$$H_A : F_{XY} \neq F_X F_Y.$$

A test is universally consistent if its power converges to 1 as $n \rightarrow \infty$ against any dependent F_{XY} .


Without loss of generality, we shall assume F_{XY} has finite second moments.

Benchmarks




Correlation
Measures


Benchmarks



Benchmarks



Benchmarks



Motivations

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;
- is easy to understand and efficient to implement;

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;
- is easy to understand and efficient to implement;
- *provides insights into the dependency structure.

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;
- is easy to understand and efficient to implement;
- *provides insights into the dependency structure.

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;
- is easy to understand and efficient to implement;
- *provides insights into the dependency structure.

Existing method has pros and cons with respect to each point.

Motivations

Modern data sets may be **high-dimensional, nonlinear, noisy, of limited sample size, structured, from disparate spaces**. Thus we desire a test that

- is consistent against all dependencies;
- has good finite-sample testing performance;
- is easy to understand and efficient to implement;
- *provides insights into the dependency structure.

Existing method has pros and cons with respect to each point.

To that end, we propose the **multiscale generalized correlation** in [*Shen et al.(2017a)*][1].

Overview


1. Illustration
2. Experiments
3. Theory
4. Summary

Illustration

Introducing MGC

Introducing MGC


$(\mathcal{X}_n, \mathcal{Y}_n)$




Introducing MGC

$(\mathcal{X}_n, \mathcal{Y}_n) \rightarrow$ Computing
Distances
& Centering:
 $A, B \in \mathbb{R}^{n \times n}$

Introducing MGC




Introducing MGC




$$D_{cov} = \sum_{i \neq j}^n A_{ij} B_{ij}$$


Introducing MGC




Introducing MGC



Introducing MGC



Introducing MGC



Direct Maximum

Direct Maximum

Directly taking the maximum local correlation

$$\max_{(k,l) \in [n]^2} \{Dcorr^{k,l}(\mathcal{X}_n, \mathcal{Y}_n)\}$$

will yield a biased statistic under independence, i.e., the maximum is always larger than 0 in expectation even under independent relationship!

Smoothed Maximum

Smoothed Maximum

Instead, we take a smoothed maximum:

Smoothed Maximum

Instead, we take a smoothed maximum:

Pick a threshold $\tau \geq 0$, compute the set

$$\{(k, l) \text{ such that } Dcorr^{k,l}(\mathcal{X}_n, \mathcal{Y}_n) > \max\{\tau, Dcorr(\mathcal{X}_n, \mathcal{Y}_n)\}\},$$

Smoothed Maximum

Instead, we take a smoothed maximum:

Pick a threshold $\tau \geq 0$, compute the set

$$\{(k, l) \text{ such that } D_{corr}^{k,l}(\mathcal{X}_n, \mathcal{Y}_n) > \max\{\tau, D_{corr}(\mathcal{X}_n, \mathcal{Y}_n)\}\},$$

and calculate the largest connected component R of the set.

Smoothed Maximum

Instead, we take a smoothed maximum:

Pick a threshold $\tau \geq 0$, compute the set

$$\{(k, l) \text{ such that } Dcorr^{k,l}(\mathcal{X}_n, \mathcal{Y}_n) > \max\{\tau, Dcorr(\mathcal{X}_n, \mathcal{Y}_n)\}\},$$

and calculate the largest connected component R of the set. If there are sufficiently many elements in R ($> 2n$), take the maximum correlation within R as MGC.

Smoothed Maximum

Instead, we take a smoothed maximum:

Pick a threshold $\tau \geq 0$, compute the set

$$\{(k, l) \text{ such that } Dcorr^{k,l}(\mathcal{X}_n, \mathcal{Y}_n) > \max\{\tau, Dcorr(\mathcal{X}_n, \mathcal{Y}_n)\}\},$$

and calculate the largest connected component R of the set. If there are sufficiently many elements in R ($> 2n$), take the maximum correlation within R as MGC.

τ is chosen based on an approximate null distribution of $Dcorr$ (symmetric Beta), which converges to 0 as $n \rightarrow \infty$.

Smoothed Maximum

Instead, we take a smoothed maximum:

Pick a threshold $\tau \geq 0$, compute the set

$$\{(k, l) \text{ such that } Dcorr^{k,l}(\mathcal{X}_n, \mathcal{Y}_n) > \max\{\tau, Dcorr(\mathcal{X}_n, \mathcal{Y}_n)\}\},$$


and calculate the largest connected component R of the set. If there are sufficiently many elements in R ($> 2n$), take the maximum correlation within R as MGC.

τ is chosen based on an approximate null distribution of $Dcorr$ (symmetric Beta), which converges to 0 as $n \rightarrow \infty$.


It is a critical step for both the finite-sample performance and certain theoretical properties of MGC.

Examples



Examples



Examples





Examples




$$Dcorr(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$
$$MGC(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$

Examples





$$Dcorr(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$

$$MGC(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$





Examples





$$Dcorr(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$

$$MGC(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$





Examples



$$Dcorr(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$

$$MGC(\mathcal{X}_n, \mathcal{Y}_n) = 0.15$$



$$Dcorr(\mathcal{X}_n, \mathcal{Y}_n) = 0.01$$


$$MGC(\mathcal{X}_n, \mathcal{Y}_n) = 0.13$$

Experiments

Visualizations of 20 Simulation Settings

Visualizations of 20 Simulation Settings

MGC, Distance Correlation, and Pearson's Correlation for 20 Dependencies



Evaluation Criterion


- Power is the probability of rejecting the null when the alternative is true.

Evaluation Criterion

- Power is the probability of rejecting the null when the alternative is true.
- Required sample size $N_{\alpha,\beta}(c)$ to achieve a power of β at type 1 error level α using a statistic c .

Testing Power: Linear vs Nonlinear

Testing Power: Linear vs Nonlinear



$$\begin{aligned}n &= 30, p = q = 1, \\ X &\sim \text{Uniform}(-1, 1), \\ \epsilon &\sim \text{Normal}(0, \text{noise}), \\ Y &= X + \epsilon \text{ and } Y = X^2 + \epsilon.\end{aligned}$$

Required Sample Size

Required Sample Size

When $noise = 1$, $p = q = 1$, the required sample size $N_{\alpha=0.05, \beta=0.85}(c)$:

Required Sample Size

When $noise = 1$, $p = q = 1$, the required sample size $N_{\alpha=0.05, \beta=0.85}(c)$:
in linear relationship, 40 for all three methods;
in quadratic relationship, 80 for MGC, 180 for Dcorr, and > 1000 for Pearson.

Required Sample Size

When $noise = 1$, $p = q = 1$, the required sample size $N_{\alpha=0.05, \beta=0.85}(c)$:
in linear relationship, 40 for all three methods;
in quadratic relationship, 80 for MGC, 180 for Dcorr, and > 1000 for Pearson.

Next we compute the size for each simulation, and summarize by the median over close-to-linear (type 1-5) and strongly non-linear relationships (type 6-19).

Required Sample Size

When $noise = 1$, $p = q = 1$, the required sample size $N_{\alpha=0.05, \beta=0.85}(c)$:
in linear relationship, 40 for all three methods;
in quadratic relationship, 80 for MGC, 180 for Dcorr, and > 1000 for Pearson.

Next we compute the size for each simulation, and summarize by the median over close-to-linear (type 1-5) and strongly non-linear relationships (type 6-19).

We consider univariate (1D) and multivariate (10D) cases.

Median Size Table

Testing Methods	1D Lin	1D Non-Lin	10D Lin	10D Non-Lin
MGC	50	90	60	165
Dcorr	50	250	60	515
Pearson / RV / CCA	50	>1000	50	>1000
HHG	70	90	100	315
HSIC	70	95	100	400
MIC	120	180	n/a	n/a

A Proteomics Experiment: Identify Biomarker

A Proteomics Experiment: Identify Biomarker

The blood samples of 95 individuals are collected. Pancreatic (10), ovarian (24), colorectal cancer (28), and healthy controls (33).

A Proteomics Experiment: Identify Biomarker

The blood samples of 95 individuals are collected. Pancreatic (10), ovarian (24), colorectal cancer (28), and healthy controls (33).

The protein and peptide abundance levels are measured for each person for 318 peptides / proteins.

A Proteomics Experiment: Identify Biomarker

The blood samples of 95 individuals are collected. Pancreatic (10), ovarian (24), colorectal cancer (28), and healthy controls (33).

The protein and peptide abundance levels are measured for each person for 318 peptides / proteins. The target is to find potential biomarkers that best distinguish between pancreatic and non-pancreatic.

A Proteomics Experiment: Identify Biomarker

The blood samples of 95 individuals are collected. Pancreatic (10), ovarian (24), colorectal cancer (28), and healthy controls (33).

The protein and peptide abundance levels are measured for each person for 318 peptides / proteins. The target is to find potential biomarkers that best distinguish between pancreatic and non-pancreatic.

For each feature, we compute MGC and the p-values between the abundance level and the label vector ($n = 95, p = q = 1$).

A Proteomics Experiment: Identify Biomarker

The blood samples of 95 individuals are collected. Pancreatic (10), ovarian (24), colorectal cancer (28), and healthy controls (33).

The protein and peptide abundance levels are measured for each person for 318 peptides / proteins. The target is to find potential biomarkers that best distinguish between pancreatic and non-pancreatic.

For each feature, we compute MGC and the p-values between the abundance level and the label vector ($n = 95, p = q = 1$).

Adjusted for multiple testing, MGC uniquely revealed one particular protein, neurogranin, which is exclusively expressed in brain tissue among normal tissues and has not been linked with any other cancer type.

Comparison and Verification

Comparison and Verification

HSIC identifies two peptides, HHG identifies three peptides, and other methods do not identify any peptide as significant. However, there exists strong evidence that the other two peptides are upregulated in other cancers.

Comparison and Verification

HSIC identifies two peptides, HHG identifies three peptides, and other methods do not identify any peptide as significant. However, there exists strong evidence that the other two peptides are upregulated in other cancers.

If we compare kNN ($K=3$) leave-one-subject-error of these peptides:

Peptides	False Positives	True Positives
Neurogranin (MGC)	2	5
HSIC Features	4	5
HHG Features	2	1

Comparison and Verification

HSIC identifies two peptides, HHG identifies three peptides, and other methods do not identify any peptide as significant. However, there exists strong evidence that the other two peptides are upregulated in other cancers.

If we compare kNN ($K=3$) leave-one-subject-error of these peptides:

Peptides	False Positives	True Positives
Neurogranin (MGC)	2	5
HSIC Features	4	5
HHG Features	2	1

Theory

Basic Properties of Sample MGC

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.
3. Invariant: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\{\phi(x_i)\}, \{\delta(y_i)\})$ for any linear transformation ϕ, δ (i.e., rotation, scaling, translation, reflection).

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.
3. Invariant: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\{\phi(x_i)\}, \{\delta(y_i)\})$ for any linear transformation ϕ, δ (i.e., rotation, scaling, translation, reflection).
4. 1-Linear: $c(\mathcal{X}_n, \mathcal{Y}_n) = 1$ under linear relationships.

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.
3. Invariant: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\{\phi(x_i)\}, \{\delta(y_i)\})$ for any linear transformation ϕ, δ (i.e., rotation, scaling, translation, reflection).
4. 1-Linear: $c(\mathcal{X}_n, \mathcal{Y}_n) = 1$ under linear relationships.
5. 0-Indep: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} 0$ if and only if independence.

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.
3. Invariant: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\{\phi(x_i)\}, \{\delta(y_i)\})$ for any linear transformation ϕ, δ (i.e., rotation, scaling, translation, reflection).
4. 1-Linear: $c(\mathcal{X}_n, \mathcal{Y}_n) = 1$ under linear relationships.
5. 0-Indep: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} 0$ if and only if independence.
6. Consistency: At any type 1 error level α , testing power

$$\beta(c(\mathcal{X}_n, \mathcal{Y}_n)) \xrightarrow{n \rightarrow \infty} 1$$

against any dependent F_{XY} .

Basic Properties of Sample MGC

1. Boundedness: $c(\mathcal{X}_n, \mathcal{Y}_n) \in [-1, 1]$.
2. Symmetric: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\mathcal{Y}_n, \mathcal{X}_n)$.
3. Invariant: $c(\mathcal{X}_n, \mathcal{Y}_n) = c(\{\phi(x_i)\}, \{\delta(y_i)\})$ for any linear transformation ϕ, δ (i.e., rotation, scaling, translation, reflection).
4. 1-Linear: $c(\mathcal{X}_n, \mathcal{Y}_n) = 1$ under linear relationships.
5. 0-Indep: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} 0$ if and only if independence.
6. Consistency: At any type 1 error level α , testing power

$$\beta(c(\mathcal{X}_n, \mathcal{Y}_n)) \xrightarrow{n \rightarrow \infty} 1$$

against any dependent F_{XY} .

Population MGC

Suppose $(X, Y), (X', Y'), (X'', Y''), (X''', Y''')$ are *iid* as F_{XY} .

Population MGC

Suppose $(X, Y), (X', Y'), (X'', Y''), (X''', Y''')$ are *iid* as F_{XY} . Let $I(\cdot)$ be the indicator function, define two random variables

$$I_{X, X'}^{\rho_k} = I(\text{Prob}\{B(X, \|X' - X\|)\} \leq \rho_k)$$

$$I_{Y', Y}^{\rho_l} = I(\text{Prob}\{B(Y', \|Y - Y'\|)\} \leq \rho_l)$$

for $\rho_k, \rho_l \in [0, 1]$.

Population MGC

Suppose $(X, Y), (X', Y'), (X'', Y''), (X''', Y''')$ are *iid* as F_{XY} . Let $I(\cdot)$ be the indicator function, define two random variables

$$I_{X, X'}^{\rho_k} = I(\text{Prob}\{B(X, \|X' - X\|)\} \leq \rho_k)$$

$$I_{Y', Y}^{\rho_l} = I(\text{Prob}\{B(Y', \|Y - Y'\|)\} \leq \rho_l)$$

for $\rho_k, \rho_l \in [0, 1]$. Further define

$$d_X^{\rho_k} = (\|X - X'\| - \|X - X''\|) I_{X, X'}^{\rho_k}$$

$$d_{Y'}^{\rho_l} = (\|Y' - Y\| - \|Y' - Y'''\|) I_{Y', Y}^{\rho_l}$$

Population MGC

Suppose $(X, Y), (X', Y'), (X'', Y''), (X''', Y''')$ are *iid* as F_{XY} . Let $I(\cdot)$ be the indicator function, define two random variables

$$I_{X, X'}^{\rho_k} = I(\text{Prob}\{B(X, \|X' - X\|)\} \leq \rho_k)$$

$$I_{Y', Y}^{\rho_l} = I(\text{Prob}\{B(Y', \|Y - Y'\|)\} \leq \rho_l)$$

for $\rho_k, \rho_l \in [0, 1]$. Further define

$$d_X^{\rho_k} = (\|X - X'\| - \|X - X''\|) I_{X, X'}^{\rho_k}$$

$$d_{Y'}^{\rho_l} = (\|Y' - Y\| - \|Y' - Y'''\|) I_{Y', Y}^{\rho_l}$$

The population local covariance can be defined as

$$Dcov^{\rho_k, \rho_l}(X, Y) = E(d_X^{\rho_k} d_{Y'}^{\rho_l}) - E(d_X^{\rho_k}) E(d_{Y'}^{\rho_l}).$$

Population MGC

Suppose $(X, Y), (X', Y'), (X'', Y''), (X''', Y''')$ are *iid* as F_{XY} . Let $I(\cdot)$ be the indicator function, define two random variables

$$I_{X, X'}^{\rho_k} = I(\text{Prob}\{B(X, \|X' - X\|)\} \leq \rho_k)$$

$$I_{Y', Y}^{\rho_l} = I(\text{Prob}\{B(Y', \|Y - Y'\|)\} \leq \rho_l)$$

for $\rho_k, \rho_l \in [0, 1]$. Further define

$$d_X^{\rho_k} = (\|X - X'\| - \|X - X''\|) I_{X, X'}^{\rho_k}$$

$$d_{Y'}^{\rho_l} = (\|Y' - Y\| - \|Y' - Y'''\|) I_{Y', Y}^{\rho_l}$$

The population local covariance can be defined as

$$Dcov^{\rho_k, \rho_l}(X, Y) = E(d_X^{\rho_k} d_{Y'}^{\rho_l}) - E(d_X^{\rho_k}) E(d_{Y'}^{\rho_l}).$$

Normalizing and taking a smoothed maximum yield population MGC.

Sample to Population

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see *[Shen et al.(2017b)]*[2].

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see *[Shen et al.(2017b)]*[2].

1. 0-Indep: $c(X, Y) = 0$ if and only if independence.

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see *[Shen et al.(2017b)][2]*.

1. 0-Indep: $c(X, Y) = 0$ if and only if independence.
2. Convergence: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} c(X, Y)$.

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see [Shen et al.(2017b)][2].

1. 0-Indep: $c(X, Y) = 0$ if and only if independence.
2. Convergence: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} c(X, Y)$.
3. **Almost Unbiased**: $E(c(\mathcal{X}_n, \mathcal{Y}_n)) = c(X, Y) + \mathcal{O}(1/n)$.

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see [Shen et al.(2017b)][2].

1. 0-Indep: $c(X, Y) = 0$ if and only if independence.
2. Convergence: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} c(X, Y)$.
3. **Almost Unbiased**: $E(c(\mathcal{X}_n, \mathcal{Y}_n)) = c(X, Y) + \mathcal{O}(1/n)$.
4. **Diminishing Variance**: $Var(c(\mathcal{X}_n, \mathcal{Y}_n)) = \mathcal{O}(1/n)$.

Sample to Population

The population version can be equivalently defined via characteristic functions of F_{XY} , e.g.,

$$D_{cov}^{\rho_k=1, \rho_l=1}(X, Y) = \int_{t,s} |g_{XY}(t, s) - g_X(t)g_Y(s)|^2 dw(t, s),$$

see [Shen et al.(2017b)][2].

1. 0-Indep: $c(X, Y) = 0$ if and only if independence.
2. Convergence: $c(\mathcal{X}_n, \mathcal{Y}_n) \xrightarrow{n \rightarrow \infty} c(X, Y)$.
3. **Almost Unbiased**: $E(c(\mathcal{X}_n, \mathcal{Y}_n)) = c(X, Y) + \mathcal{O}(1/n)$.
4. **Diminishing Variance**: $Var(c(\mathcal{X}_n, \mathcal{Y}_n)) = \mathcal{O}(1/n)$.

The last three properties also hold for any local correlation by $(\rho_k, \rho_l) = (\frac{k-1}{n-1}, \frac{l-1}{n-1})$.

Advantage over Mantel and Dcorr

Advantage over Mantel and Dcorr

1. There exists dependent F_{XY} such that $Mantel = 0$, thus not universally consistent.

Advantage over Mantel and Dcorr

1. There exists dependent F_{XY} such that $Mantel = 0$, thus not universally consistent.
2. For any dependent F_{XY} , $MGC(X, Y) \geq Dcorr(X, Y)$.

Advantage over Mantel and Dcorr

1. There exists dependent F_{XY} such that $Mantel = 0$, thus not universally consistent.
2. For any dependent F_{XY} , $MGC(X, Y) \geq Dcorr(X, Y)$.
3. There exists dependent F_{XY} such that $MGC(X, Y) > Dcorr(X, Y)$.

Advantage over Mantel and Dcorr

1. There exists dependent F_{XY} such that $Mantel = 0$, thus not universally consistent.
2. For any dependent F_{XY} , $MGC(X, Y) \geq Dcorr(X, Y)$.
3. There exists dependent F_{XY} such that $MGC(X, Y) > Dcorr(X, Y)$.
- 4*. If the relationship is linear (or with independent noise), the global scale is always optimal and $MGC(X, Y) = Dcorr(X, Y)$.

Advantage over Mantel and Dcorr

1. There exists dependent F_{XY} such that $Mantel = 0$, thus not universally consistent.
2. For any dependent F_{XY} , $MGC(X, Y) \geq Dcorr(X, Y)$.
3. There exists dependent F_{XY} such that $MGC(X, Y) > Dcorr(X, Y)$.
- 4*. If the relationship is linear (or with independent noise), the global scale is always optimal and $MGC(X, Y) = Dcorr(X, Y)$.

Conversely, the optimal scale being local (i.e., $MGC(X, Y) > Dcorr(X, Y)$) implies a non-linear relationship.

Summary

Summary

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

- Proper distance transformation ensures the universal consistency.

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

- Proper distance transformation ensures the universal consistency.
- Compute all local correlations iteratively.

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

- Proper distance transformation ensures the universal consistency.
- Compute all local correlations iteratively.
- Identify the optimal local correlation without inflating the sample bias.

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

- Proper distance transformation ensures the universal consistency.
- Compute all local correlations iteratively.
- Identify the optimal local correlation without inflating the sample bias.

Summary

MGC builds on distance correlation, the locality principle, and taking a smoothed maximum.

- Proper distance transformation ensures the universal consistency.
- Compute all local correlations iteratively.
- Identify the optimal local correlation without inflating the sample bias.

They made MGC advantageous in theory and practice.

Advantages of MGC

Advantages of MGC

1. Performant under any joint distribution:

Advantages of MGC

1. Performant under any joint distribution:
 - Equals 0 asymptotically if and only if independence.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

2. It works for:

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

2. It works for:

- Low- and high-dimensional data.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

2. It works for:

- Low- and high-dimensional data.
- Euclidean and structured data (e.g., images, networks, shapes).

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

2. It works for:

- Low- and high-dimensional data.
- Euclidean and structured data (e.g., images, networks, shapes).
- Any dissimilarity / similarity / kernel matrix.

Advantages of MGC

1. Performant under any joint distribution:

- Equals 0 asymptotically if and only if independence.
- Amplify the dependency signal while mostly avoiding the sample bias.
- Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.

2. It works for:

- Low- and high-dimensional data.
- Euclidean and structured data (e.g., images, networks, shapes).
- Any dissimilarity / similarity / kernel matrix.

Advantages of MGC

1. Performant under any joint distribution:
 - Equals 0 asymptotically if and only if independence.
 - Amplify the dependency signal while mostly avoiding the sample bias.
 - Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.
2. It works for:
 - Low- and high-dimensional data.
 - Euclidean and structured data (e.g., images, networks, shapes).
 - Any dissimilarity / similarity / kernel matrix.
3. Intuitive to understand and efficient to implement in $\mathcal{O}(n^2 \log n)$.

Advantages of MGC

1. Performant under any joint distribution:
 - Equals 0 asymptotically if and only if independence.
 - Amplify the dependency signal while mostly avoiding the sample bias.
 - Superior finite-sample performance over all benchmarks, against linear / nonlinear / noisy / high-dimensional relationships.
2. It works for:
 - Low- and high-dimensional data.
 - Euclidean and structured data (e.g., images, networks, shapes).
 - Any dissimilarity / similarity / kernel matrix.
3. Intuitive to understand and efficient to implement in $\mathcal{O}(n^2 \log n)$.

MGC shares the same intrinsic idea as in nonlinear embedding, random forest, multiple kernel learning, deep learning.

Current Works

Current Works

- The sample method, algorithmic details, simulation advantages and real applications are demonstrated in [*Shen et al.(2017a)*][1].

Current Works

- The sample method, algorithmic details, simulation advantages and real applications are demonstrated in [*Shen et al.(2017a)*][1].
- Population MGC and most mathematical properties appear in [*Shen et al.(2017b)*][2].

Current Works

- The sample method, algorithmic details, simulation advantages and real applications are demonstrated in [*Shen et al.(2017a)*][1].
- Population MGC and most mathematical properties appear in [*Shen et al.(2017b)*][2].
- MGC is infused with diffusion maps for testing between graph vertices and attributes [*Lee et al.(2017)*][3].

Current Works

- The sample method, algorithmic details, simulation advantages and real applications are demonstrated in [*Shen et al.(2017a)*][1].
- Population MGC and most mathematical properties appear in [*Shen et al.(2017b)*][2].
- MGC is infused with diffusion maps for testing between graph vertices and attributes [*Lee et al.(2017)*][3].
- MGC is utilized for iterative signal subgraph extraction in [*Wang et al.(2018)*][4].

Future Directions

Future Directions

- Feature selection / dimension reduction.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.
- Better classification / regression in multiple graph setting.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.
- Better classification / regression in multiple graph setting.
- Rank or Kernel MGC; interpretation of the optimal local scale under RKHS framework.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.
- Better classification / regression in multiple graph setting.
- Rank or Kernel MGC; interpretation of the optimal local scale under RKHS framework.
- Relative efficiency among MGC, Dcorr, Mantel, and HSIC.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.
- Better classification / regression in multiple graph setting.
- Rank or Kernel MGC; interpretation of the optimal local scale under RKHS framework.
- Relative efficiency among MGC, Dcorr, Mantel, and HSIC.
- Faster MGC testing.

Future Directions

- Feature selection / dimension reduction.
- Magnitude of MGC versus prediction and classification error.
- Better classification / regression in multiple graph setting.
- Rank or Kernel MGC; interpretation of the optimal local scale under RKHS framework.
- Relative efficiency among MGC, Dcorr, Mantel, and HSIC.
- Faster MGC testing.
- Real data applications.

References

1. C. Shen, C. E. Priebe, M. Maggioni, Q. Wang, and J. T. Vogelstein, "Discovering relationships and their structures across disparate data modalities," <https://arxiv.org/abs/1609.05148>, 2017.
2. C. Shen, C. E. Priebe, and J. T. Vogelstein, "From distance correlation to the multiscale generalized correlation," <https://arxiv.org/abs/1710.09768>, 2017.
3. Y. Lee, C. Shen, and J. T. Vogelstein, "Network dependence testing via diffusion maps and distance-based correlations," <https://arxiv.org/abs/1703.10136>, 2017.
4. S. Wang, C. Shen, A. Badea, C. E. Priebe, and J. T. Vogelstein, "Signal subgraph estimation via vertex screening," *In prep*, 2018.