

Working master

COSMIC TIME

Working master

Updated: Sep 8, 2026

CC0 • Anonymous

Start here

This is the current entry point for Cosmic Time Architecture, planetary recurrence, and the weave comparator. It incorporates the September 7 recovery and September 8 input-verification records. The “CTA v0.7 source” tab preserves this document’s previous body, including the executed signed rematch.

Timekeeper audits clocks: name the clock, identify the timestamp or parameter’s source, specify the witness, and state the observable consequence. Geometry, physical mechanism, history, and symbolism remain separate evidence lanes.

Current source map

CTA III baseline, definitions and §§24–26: [COSMIC TIME ARCHITECTURE III](#)

PRP source-recovery record: [# CTA III §26 — PRP-0](#)

PRP input-provenance verification: [# CTA III §26 — PRP-0](#)

Modular Planetary Recurrence Geometry: [Modular Planetary Recurrence Geometry](#)

Time toroid and earlier post-freeze record: [Time toroid](#)

Timekeeper research-lane audit: [Timekeeper Ledger - Research-Lane Appendix](#)

Geometry + GQG working master: [# Geometry Maximization v2](#)

1. Current status

Planetary recurrence: saved results report module recurrence, episode structure, and differences between directed and orientation-forgetting witnesses. GLOBAL EXCESS SYNCHRONIZATION: NOT SHOWN.

PRP source recovery: the saved numerical dictionary and named upstream script are RECOVERED / VERIFIED by the cited September records. Historical execution-time identity, runtime-target equality, and exact replay remain UNRESOLVED.

Weave comparator: the signed rematch is reported PASS as deterministic model verification under the declared symmetric initialization. It is not a planetary scan, physical-clock calibration, or blind human prediction trial.

Physical mechanism and historical interpretation: no promotion follows from these geometric or recovery results. Physical Q2 remains NOT_RUN. Older Q2 sampler screens belong to separate pilot records and do not alter that physical-test status.

This consolidation reviews and organizes the records; it does not rerun their simulations, regenerate an ephemeris, or independently repeat their byte checks.

2. The geometry and its witnesses

A three-phase model $\theta \in T^3$ has two independent relative phases:

$$\varphi = (\theta_2 - \theta_1, \theta_3 - \theta_1) \in T^2.$$

The quotient removes common rotation. Keep absolute phase whenever the selected witness needs it. The collective phase carries a clock hand; the relative state diagnoses lock, slip, or drift.

A directed witness preserves signed geometry; a shell witness forgets orientation. A recurrence means return under the specified witness and tolerance or score, not identity of the full sky. Historical and symbolic views are downstream records.

Lift continuous relative paths after choosing an initial lift. Weave winding $m \in \mathbb{Z}^2$ records that path history. It is distinct from gauge winding $W \in \mathbb{Z}^3$ and parity $q = W \bmod 2$; shared terminology supplies no map between them.

Recurrence, crossing, slip, lock, and tick are different event classes. A tick requires a declared hand and oriented crossing rule. Sampling must resolve the chosen event; wrapped samples alone can hide turns or direction.

3. What the weave actually establishes

The exploratory coupled comparator is

$$\theta_i = \omega_i + (K/3) \sum_j \sin(\theta_j - \theta_i).$$

On the invariant branch $\theta_1 = \theta_2$, the relative phase $a = \theta_3 - \theta_1$ obeys $\dot{a} = \delta - K \sin a$. It is a model law with declared parameters, not an ephemeris-derived planetary interaction.

The collective overlap squared is $C(a) = (5 + 4 \cos a)/9$. States $a = \pm\pi/2$ both give $C = 5/9$, but their derivatives differ by $-8\delta/9$. At $\delta = 0.003$ and $K = 0.0015$, the rates are $-1/1500$ and $+1/500$. Thus the same overlap can be falling or rising under the same law. For sufficiently small positive sampling intervals, overlap is not an autonomous predictive state on the full symmetric circle.

Keeping $(\cos a, \sin a)$ restores the branch phase record.

The CTA3-SIGNED-REMATCH-001 record reports 18 trajectories: $\delta = +0.003$, -0.003 , and 0 at $K/d = 0, 0.5, 0.9, 1, 1.1$, and 2, with $d = 0.003$, $\theta(0) = (0, 0, 0)$, and fixed $u = 0.003t$ through $u = 600$.

Half-step runs and the analytic scalar flow provided checks.

Signed pairs had equal overlap and opposite lifted phase. Drifting pairs completed ± 95 , ± 82 , and ± 41 turns. At $K/d = 0.5$, principal-cut slip counts were ± 83 , distinct from complete turns. The maximum reported primary phase error was 1.616×10^{-8} radians against a frozen 5×10^{-7} tolerance. The symmetry was predicted analytically before verification.

4. PRP: recovered specification and saved findings

The recovered freeze declares Swiss Ephemeris 2.10.03, Moshier, apparent geocentric ecliptic coordinates of date, on daily samples from proleptic Gregorian 0001-01-01 through 2026-08-27 inclusive at 12:00 UT. UT is the source wording; a historical UTC realization is not independently established.

The eight primary bodies are Mercury, Venus, Mars, Jupiter, Saturn, Uranus, Neptune, and Pluto. The saved dictionary contains 72 entries: 61 primary across 16 source families and 11 extended.

The implementation compares all unordered index pairs within each ordered body tuple to numerical source templates. Ideal motif names do not replace those targets. Empirical rank probabilities, component Bonferroni adjustment, and strict contiguous K01≥2 episodes govern acceptance; 2° motif assignment is not a universal episode tolerance.

The null uses 500 family-shift controls per witness, moving family modules and exclusions together. The implementation resets the same seed, 437113, for each witness: separate evaluation does not mean independently seeded schedules. Upper-tail scalar p-values use $(1+\text{count})/501$.

Saved results, as checked in the recovery record:

Directed witness — 724,920 eligible days; 2,455 strict episodes; 338.6581 episodes per 100,000 days; episode-rate $p=1.0$; reported maximum-peak $p\approx 0.47106$.

Shell witness — 724,920 eligible days; 3,336 strict episodes; 460.1887 episodes per 100,000 days; episode-rate $p=1.0$; reported maximum-peak $p\approx 0.32136$.

These support the retained status: GLOBAL EXCESS SYNCHRONIZATION: NOT SHOWN.

5. What recovery settled—and what remains

The September 8 verification reports byte equality for the 110,892-byte dictionary and 30,928-byte upstream script against separately retrieved copies of the original archive. All 72 static definitions and 650 stored pair targets passed the stated consistency checks. Those copies remain evidence of the same historical artifacts, not independently generated runs.

The episode runner is manifest-bound, but its imported upstream dependency is not. It calls `build_modules()` to regenerate targets; it does not load the saved dictionary as its runtime input. Consequently, saved-target consistency does not establish historical runtime-target equality.

The historical manifest matches 15 of 16 entries. The full `run.log` differs, while its first 425 bytes match the recorded hash; an appended output suffix is consistent with the runner's ordering.

Preserve the exception.

The source prose and code also differ in directed operand order and seam convention. Keep that discrepancy visible; its effect on the historical scores has not been established.

Remaining recovery targets are the 61 original directed probability arrays, the original daily position-cache bytes and provenance, an execution-linked upstream digest, runtime targets, pinned numerical environment and launch record, and actual shifts or complete lexicographic controls. A regenerated replacement would be a new reconstruction run with new identities.

6. Rules for the next claim

Freeze bodies, interval, frame, source-to-phase constructor, witness, module dictionary, scoring, tolerances, orientation, exclusions, null, multiplicity handling, and code/run identities before inspecting the evaluation. Freeze geometric episodes before joining event labels.

Keep raw timestamps, coordinates, frame, and source immutable; each derived view gets its own identity. Label parameter custody: source-given, exact-derived, legacy, exploratory, fitted, or arbitrary control.

The legacy ≈ 43 -day and ≈ 21.5 -day arithmetic is not a detected cosmic periodicity. Days and degrees require an explicit conversion model. Timing adjacency alone does not establish historical causation or common control.

Use this first tab for current status. Keep source details and older editions behind it. The next research step is recovery of the remaining execution-linked inputs, or an explicitly new frozen reconstruction—not another summary claiming the old replay is complete.

7. Temporal presentation and sampling

The temporal exchange-rate increment is an application to interaction timing. In a Cosmic Time tool, distinguish the source timestamp, model parameter, sampling instant, computation completion, and user-visible release. A change in interface latency is not a change in astronomical or model time.

Preserve the raw sample times and orientation/lift information before buffering, resampling, or interpolation. Regular-looking output cannot recover an unobserved turn or crossing. The sampling requirements in source §29 and the existing source-to-phase constructor remain necessary.

For a declared interface task, test mean latency and one named irregularity measure against a primary user outcome. A benefit from added waiting and a stronger effect of irregularity at low latency are separate claims. The temporal note supplies exact examples and a test design, not a new ephemeris run, planetary mechanism, historical association, or physical-clock calibration.

Temporal exchange rate v0.3 — interface timing application:  # Temporal exchange rate

[Project sources and earlier versions](#)

[All active projects](#)

CTA v0.7 source

Cosmic Time Architecture III · v0.7 executed additions

****Predictive witnesses, signed controls, sampling, and the completed rematch****
6 September 2026

Sections 27–32 extend CTA III v0.6. The baseline and pre-run proposal are preserved in this packet; §32 records the new execution and updates the proposal's run status.

27. Predictive fidelity — does the witness retain the next step?

Recurrence fidelity and predictive fidelity are separate questions. A witness can correctly detect a repeated pattern while omitting information needed to predict its continuation.

For a declared state domain D , fixed dynamics $\Phi_{\Delta t}: D \rightarrow D$, sampling interval Δt , and witness V , an autonomous observed update exists precisely when

$$\begin{aligned} & \\ & V(x) = V(y) \\ & \quad \Longleftrightarrow \\ & V(\Phi_{\Delta t} x) = V(\Phi_{\Delta t} y) \\ & \quad \quad (x, y \in D). \end{aligned}$$

The proof is direct: define the observed successor using any full-state representative. It is well-defined exactly when the representative does not change the answer. Inputs, parameters, and the sampling interval must be the same for both states, or explicitly included in the predictive record.

For stochastic models, replace equality of successors by equality of their entire next-observation laws. A fitted transition table or matching average is insufficient to establish that condition.

An exact counterexample within the W_2 model

On W_2 's invariant symmetric branch $\theta_1 = \theta_2$, let $a = \theta_3 - \theta_1$. The existing reduction is

$$\begin{aligned} & \\ & \dot{a} = \delta - K \sin a. \end{aligned}$$

The collective overlap, written $R_{\{\mathrm{coll}\}}$ here to distinguish its scalar meaning, satisfies

$\$$

$$R_{\{\mathrm{coll}\}}=\frac{|2+e^{ia}|^3}{3},\quad C(a):=R_{\{\mathrm{coll}\}}^2=\frac{5+4\cos a}{9}.$$

The two states $a_+=\pi/2$ and $a_-=-\pi/2$ have the same observed value $C=5/9$. Nevertheless,

$$\dot{C}=-\frac{4}{9}\sin a, \quad (\delta-K\sin a),$$

so

$$\dot{C}(a_+)=\frac{4}{9}(K-\delta), \quad \dot{C}(a_-)=\frac{4}{9}(K+\delta),$$

and their difference is $-8\delta/9$. For nonzero detuning, the derivatives differ. Their observed successors therefore differ for all sufficiently small positive sampling intervals.

At the existing comparator parameter values $\delta=0.003$ and $K=0.0015$:

Initial relative phase	Initial overlap squared	Initial rate of overlap squared
$+\pi/2$	$5/9$	$-1/1500$ per model-time unit
$-\pi/2$	$5/9$	$+1/500$ per model-time unit

****The same overlap value is decreasing in one state and increasing in the other.**** These are newly constructed starting states under the same law, not additional observations from the original zero-initialized W_2 run.

This disproves predictive closure of collective overlap on the full symmetric relative-phase circle for sufficiently fine sampling. It does not claim failure on every restricted domain or at every possible sampling interval. In particular, an injective restriction of the witness can retain more predictive information than the unrestricted witness.

A sufficient continuous phase record on this branch is $(c,s)=(\cos a,\sin a)$:

$$\dot{c}=-s(\delta-Ks), \quad \dot{s}=c(\delta-Ks), \quad c^2+s^2=1.$$

The first two clocks coincide only on the symmetric branch. A general weave requires both relative phases. Cumulative winding additionally requires the chosen lift and its history.

The Geometry Maximization and Twin Timelines work supplies a complementary exact benchmark: two phases share 39 slip observations and differ at observation 40. That construction demonstrates finite-history ambiguity in its own phase model. It provides a test pattern for witness design, without identifying its constants or clock with the astronomical system.

28. Signed detuning — the mirror control

W_2 executed positive detuning. A useful extension is a paired $\pm d$ comparison with the same $K \geq 0$ and initial relative phase zero.

Uniqueness of the smooth scalar flow and oddness of sine give the exact relation

$$a(t;-d) = -a(t;+d).$$

Consequently the entire overlap trace is identical:

$$R_{\{\mathrm{coll}\}}(t;-d) = R_{\{\mathrm{coll}\}}(t;+d),$$

while the lifted phase changes sign. For $K < d$, the complete oriented turns have opposite signs and equal period

$$T_{\{\mathrm{turn}\}} = \frac{2\pi}{\sqrt{d^2 - K^2}}.$$

This comparison changes the parameter sign. It tests whether a witness preserves orientation; it is separate from §27's counterexample under one fixed law.

The positive-detuning descriptions in §24 remain valid for that run. A sign-general extension must use the direction of δ : at $K = d$, the zero-initialized relative phase approaches $\operatorname{sgn}(\delta)\pi/2$; for $K > d$, it approaches $\arcsin(\delta/K)$ on the stable branch.

Cut conventions are part of the signed record

With principal phase in $[-\pi, \pi)$, the deck label is

$$m(a)=\left\lfloor\frac{a+\pi}{2\pi}\right\rfloor.$$

Away from the cut, $m(-a)=-m(a)$. At the cut $a=(2k+1)\pi$, the half-open convention instead gives $m(-a)=1-m(a)$. The continuous lift still reverses exactly. Signed crossing events and instantaneous deck labels must therefore be checked with their stated endpoint conventions.

The existing planetary directed witness uses $(-180^\circ,180^\circ]$, while W_2 uses $[-\pi,\pi)$. At the antipodal point these display $+180^\circ$ and $-\pi$ respectively. They represent the same circle point, but their signed representatives differ. Unit conversion must transport the cut convention as well as multiply by a scale.

The paired numerical extension was executed on 6 September 2026 and passed all six coupling settings. Section 32 records its result. The flow symmetry and observation equality above remain exact mathematical consequences.

29. Sampling fidelity — a still picture can hide a full turn

Consider three synthetic lifted phases in cycles:

$$p_0(t)=0,\quad p_+(t)=t,\quad p_-(t)=-t.$$

At integer sampling times n , every wrapped observation equals zero. Yet the lifted displacements per interval are 0, +1, and -1 turns. All three corresponding overlap samples equal one.

This is exact sampling aliasing. Wrapped samples alone cannot determine the intervening winding when the intervening dynamics are unspecified. An independently known motion model or speed bound supplies additional information; a smoother visualization does not.

For the reduced weave equation,

$$|\dot{a}|\leq |\delta|+K.$$

Under exact phase readouts, a sufficient condition for a unique nearest-increment reconstruction between adjacent samples is

$$|$$

$(|\delta|+K)\Delta t < \pi$.

\$\$

This bounds the actual phase displacement strictly below half a turn. It is sufficient, not necessary; equality is ambiguous. It certifies the net lift increment under its assumptions. In a more general trajectory, detecting every intermediate crossing or recrossing additionally requires an event-resolving path record or justified event brackets. Observation uncertainty must be carried into the reconstruction; ambiguous intervals remain unresolved.

Sampling contract

Attach the following record to each numerical or astronomical extension:

Field	Required distinction
---	---
Source clock	Timestamp, time scale, origin, and units
Model clock	Physical time, declared model time, or normalized time; W_2 uses $\tau= \delta t$
Sampling rule	Fixed cadence, actual irregular times, or event-triggered output
Phase representation	Radians, degrees, or cycles; relative versus absolute phase
Lift and cut	Initial lift, principal interval, seam ties, signed crossing rule
Numerical record	Internal solver steps distinguished from saved samples and interpolated display points
Completeness	Missing intervals, uncertainty, and which crossings are certified

A constant rescaling of time and sampling only at selected events are different operations. Event records should not be interpreted as uniformly timed occupation samples without an appropriate argument or weighting.

This contract supplements the existing T_{dyn} , T_{addr} , and T_{bind} distinction. It supplies no conversion from model time or Monte Carlo updates to physical time.

30. Separate return, prediction, and exploration

The latest project work supplies three different tests:

Test	What it establishes	Current scope
---	---	---
W_2 detuning comparator	Detuning changes the declared relative dynamics and return geometry	Original execution reported in CTA III §24; fresh signed verification recorded in §32
Twin Timelines	Equal finite histories can have unequal next observations	Constructed exact phase example; no blind human trial result claimed
Toroidal Q2 pilot	Coarse sector diagnostics can pass while signed winding fails its screens	Completed separate lattice sampler; $L=3$ and $L=4$ remain unresolved

At L=4, for example, the Q2 pilot recorded signed-W effective sample size about 336 and R-hat about 1.03469, outside its declared screens. Physical Q2 remains unresolved. These are sampler results under a different state space and clock, not estimates of astronomical exploration or weave mixing.

Retain the existing distinction

\$\$

$$\mathbf{m}_{\mathrm{weave}} \in \mathbb{Z}^2,$$

$$\quad W_{\mathrm{gauge}} \in \mathbb{Z}^3,$$

$$\quad q_{\mathrm{gauge}} = W_{\mathrm{gauge}} \bmod 2.$$
 \$\$

No map between these registers is created by the shared word “winding.” Likewise recurrence does not establish mixing, and predictive closure does not establish stability.

The useful transferable practice is to check each retained observable against the claim being made. The [checks and evidence record](EVIDENCE_CROSSWALK.json) distinguishes the new exact examples from imported results.

31. Extension order and current status

The original symmetric weave question in §17 has a conditional mathematical answer in §24. Add a cross-reference there so the historical question is not mistaken for an experiment that still has not run.

The paired sign-control plan, witnesses, sampling schedule, and evaluation rules have now been executed, as recorded in §32. Overlap equality and signed-lift reversal were scored separately. The two-state witness collision and sampling alias also passed their checks. A future blind presentation may hide labels until a prediction is recorded; no blind human trial was performed here.

For a new planetary claim, the immediate priority remains §26's incomplete reproduction package: recover the scan's ephemeris realization, exact body set, complete module dictionary, numeric tolerances, code identity, run identity, and full null specification. W_1 , W_2 , Twin Timelines, and the Q2 pilot cannot fill those fields for the earlier aggregate scan.

****Status of this addendum:**** exact witness and sampling examples checked; paired numerical weave extension executed and verified within the declared comparator; no new ephemeris scan, physical mechanism test, or historical association claimed. The previously reported planetary result remains ****GLOBAL EXCESS SYNCHRONIZATION: NOT SHOWN****.

Keep the clock. Keep the direction. Keep the missing information visible. Then test whether the next step is actually determined.

32. Executed signed rematch — CTA3-SIGNED-REMATCH-001

****Status:** PASS / deterministic model verification. ****** The frozen plan compared $\delta=+0.003$, -0.003 , and 0 at $K/d=0, 0.5, 0.9, 1, 1.1$, and 2, from $\theta(0)=(0,0,0)$. All 18 full-system trajectories were repeated at half the time step and checked against the analytic scalar flow over $u=600$, with the same fixed clock $u=0.003t$ for every condition.

The paired overlap values matched at every saved time, while the signed relative phases reversed. The drifting pairs completed $\pm 95, \pm 82$, and ± 41 full turns. At $K/d=0.5$ the corresponding signed slip counts were ± 83 ; complete turns and principal-cut crossings remain separate. The critical case approached $\pm \pi/2$ without completing a turn, and the stronger couplings approached opposite locked offsets.

The maximum primary phase error against the analytic reference was $1.616e-08$ radians, below the frozen $5e-7$ tolerance. The same-law equal-overlap counterexample and exact sampling/cut checks passed. Settings and thresholds were not changed after execution began.

The trajectory symmetry was predicted analytically before this verification. No untouched discovery, blind human prediction, planetary mechanism, or physical-clock calibration is claimed. The symmetric initial condition remains part of the result's scope.

[Execution report](REPORT.md) · [Frozen plan](PLAN.json) · [Full results](RESULTS.json) · [Supporting cases](SUPPORTING_CASES.json) · [Analytic reference](ANALYTIC_REFERENCE.md) · [Saved-data verification](DATA_VERIFICATION.json)

****Reuse:**** CC0, following the source document.

****Baseline:**** [byte-preserved CTA III v0.6](baseline/CTA_III_v0.6.txt).

****Sources and verification:**** [source identities](FREEZE.json), [exact checks and imported evidence](EVIDENCE_CROSSWALK.json).