

## Large-vessel vasculitis in positron emission tomography and ultrasound fusion imaging

A 65-year-old female was referred with loss of weight, dyspnoea and night sweats. Laboratory tests revealed elevated inflammatory markers. Imaging with  $^{18}\text{F}$ -labelled fluorodeoxyglucose ( $^{18}\text{F}$ -FDG) positron emission tomography/computed tomography (PET/CT) showed increased  $^{18}\text{F}$ -FDG-uptake in the axillary arteries and the aorta (Fig. 1A, blue and green arrows). Large-vessel vasculitis was suspected, given the high specificity (89%) and sensitivity (80%) of PET/CT for this diagnosis [1]. The limited soft-tissue contrast of CT images impedes a detailed assessment of morphological vascular pathologies. These can be better examined using US, which depicts vascular wall thickening as a sign of inflammatory processes (Fig. 1B, blue arrow).

PET/US fusion was performed, combining the advantages of metabolic (PET) and morphological (US) examinations [2]. The Volume Navigation technology of GE Healthcare's LOGIQ<sup>TM</sup> E9 US device allows the merging of previously acquired PET datasets with real-time US. Figure 1C shows the accurate correlation of PET and US images in the left-sided axillary artery. Additional Doppler sonography was applied to assess the perfused vessel lumen (Fig. 1D). Sagittal images of PET, CT and US in Fig. 1E–G demonstrate the co-registration of PET and US images, using spine and liver as morphological landmarks. The increased  $^{18}\text{F}$ -FDG-uptake correlates with the pathologically thickened wall of the abdominal aorta (Fig. 1E–G, green arrows and videos of transversal and sagittal PET/US fusion imaging of the abdomen, available as supplementary data at *Rheumatology* Online).

**Funding:** No specific funding was received from any funding bodies in the public, commercial or not-for-profit sectors to carry out the work described in this manuscript.

**Disclosure statement:** The authors have declared no conflicts of interest.

## Supplementary data

Supplementary data are available at *Rheumatology* Online.

**Philipp Seifert<sup>1</sup>, Robert Drescher<sup>1</sup>, Alexander Pfeil<sup>2</sup> and Martin Freesmeyer<sup>1</sup>**

<sup>1</sup>Clinic of Nuclear Medicine and <sup>2</sup>Department of Rheumatology, Clinic of Internal Medicine, Jena University Hospital, Jena, Germany

Correspondence to: Martin Freesmeyer, Clinic of Nuclear Medicine, Jena University Hospital, Am Klinikum 1, Gebäude A4 Ebene U1, 07747 Jena Lobeda-Ost, Germany.  
E-mail: martin.freesmeyer@med.uni-jena.de

## References

- 1 Danve A, O'Dell J. The role of  $^{18}\text{F}$  fluorodeoxyglucose positron emission tomography scanning in the diagnosis and management of systemic vasculitis. *Int J Rheum Dis* 2015;18:714–24.
- 2 Drescher R, Müller A, Lesser T, Freesmeyer M. Differential diagnosis of thyroid nodules via real-time PET/ultrasound (US) fusion in a case of co-existing medullary thyroid cancer and adenoma. *J Clin Endocrinol Metab* 2013;98:4250–1.

**Fig. 1** PET/CT/US fusion images of axillary artery and abdominal aorta in large-vessel vasculitis

