

34 Jewett LR, Razykov I, Hudson M, Baron M, Thombs BD; Canadian Scleroderma Research Group. Prevalence of current, 12-month and lifetime major depressive disorder among patients with systemic sclerosis. *Rheumatology* 2014;53:1719.

35 Thombs BD, Jewett LR, Kwakkenbos L, Hudson M, Baron M; Canadian Scleroderma Research Group. Major depression diagnoses are often transient among patients with systemic sclerosis: baseline and 1-month follow-up. *Arthritis Care Res* 2015;67:411–6.

## Clinical vignette

*Rheumatology* 2017;56:1311

doi:10.1093/rheumatology/kex013

Advance Access publication 14 March 2017

### A case of vanishing skull: Gorham's disease

Gorham's disease (GD) is a rare form of lymphangioma-tosis associated with profound osteolysis. Outcome depends on the site affected. Small limb lesions can cause fractures, while spinal involvement or chylothorax associated with rib lesions can be life threatening. There is no consensus regarding treatment. Surgical resection and/or radiotherapy are most commonly used, but often unsuccessfully [1].

We report the case of an otherwise healthy female who presented at the age of 40 years with a small, painless skull indentation. Over the following 6 years the indentation evolved into an 11 × 11 cm cranial vault defect. There were no other abnormal findings on examination. Investigations were negative for malignancy, infection, biochemical and immunological abnormalities. CT demonstrated a defect in the occipital region (Fig. 1) but no abnormalities elsewhere in the skeletal system. A skull and dural biopsy confirmed GD.

Radiotherapy proved ineffective. Monthly 5 mg i.v. infusion of zoledronic acid was therefore commenced, with weekly s.c. pegylated IFN- $\alpha$ 2b 35  $\mu$ g. Clinical and radiological follow-up over 3 years demonstrated suppression of the osteolytic process with stabilization of the calvarial defect.

Lymphangiomatoses are non-malignant processes involving abnormal lymphatic proliferation. In GD, an increased number of osteoclasts are often present, which is thought to be promoted by elevated levels of IL-6 [2]. While IFN and bisphosphonate can provide effective therapy, cases with poor response may benefit from trials of anti-cytokine therapy or other biologic modulators in rheumatologic research settings.

**Funding:** No specific funding was received from any bodies in the public, commercial or not-for-profit sectors to carry out the work described in this article.

**Disclosure statement:** The authors have declared no conflicts of interest.

**Sizheng Zhao<sup>1,2</sup>, David Richardson<sup>3</sup>,  
Madhu Mahindrakar<sup>1</sup> and Jagdish R. Nair<sup>1</sup>**

<sup>1</sup>*Rheumatology Department, Aintree University Hospital,*

<sup>2</sup>*Institute of Ageing and Chronic Disease, University of*

**Fig. 1** CT head with 3D rendering of the skull defect



*Liverpool and <sup>3</sup>Department of Oral and Maxillofacial Surgery, Aintree University Hospital, Liverpool, UK*

Correspondence to: Jagdish Nair, Clinical Sciences Centre, University Hospital Aintree, Lower Lane, Liverpool, L9 7AL, UK. E-mail: jagdish.nair@aintree.nhs.uk

### References

- 1 Kuriyama DK, McElligott SC, Glaser DW, Thompson KS. Treatment of Gorham-Stout disease with zoledronic acid and interferon- $\alpha$ : a case report and literature review. *J Pediatr Hematol Oncol* 2010;32:579–84.
- 2 Dellinger MT, Garg N, Olsen BR. Viewpoints on vessels and vanishing bones in Gorham-Stout disease. *Bone* 2014;63:47–52.