

## Anti-NXP2 antibody-associated extensive subcutaneous calcinosis in adult-onset myositis

A 68-year-old woman was followed since 1988 for DM. A combined therapy of corticosteroids and MTX controlled muscle involvement. In 1993, multiple subcutaneous nodules of calcinosis appeared on the limbs, chest and abdomen. In 2013, extensive calcinosis severely impaired the patient's quality of life while myositis was under control (Fig 1). In 2016, a blood sample was collected in which circulating anti-NXP2 antibodies were detected.

Anti-NXP2 antibodies were first identified in 1997. Today, 25% of patients with juvenile DM demonstrate these autoantibodies in their blood samples, indicating a poor prognosis [1]. Anti-NXP2 antibodies are detected in only 1% of cases of adult patients with myositis, and are likely linked to the occurrence of neoplasms [2]. Therefore, although typical in paediatric patients, an extensive form of calcinosis in an adult case of anti-NXP2 antibody-associated myositis is extremely rare. In our case, the evolution of calcinosis is independent from muscle involvement. Furthermore, MTX, corticosteroids and IVIGs were inefficient in preventing the spreading of calcinosis. Currently, a topical sodium thiosulphate treatment is being trialled.

Thus, an early detection of anti-NXP2 antibodies could help identify a subset of adult patients at risk of developing calcinosis during the early stages of DM.

**Funding:** No specific funding was received from any bodies in the public, commercial or not-for-profit sectors to carry out the work described in this manuscript.

**Disclosure statement:** P.G. is a medical expert for Laboratoire Français du Biofractionnement (LFB) and has received fees from Abbvie, Boehringer Ingelheim France, Pfizer, and Roche in the last 5 years. The other authors have declared no conflicts of interest.

**Radjiv Goulabchand<sup>1,2</sup>, Philippe Guilpain<sup>1,2,3</sup>, Catherine Cyteval<sup>2,4,5</sup> and Alain Le Quellec<sup>1,2</sup>**


<sup>1</sup>Department of Internal Medicine and Multi-Organic Diseases, St Eloi Hospital, CHRU Montpellier, <sup>2</sup>Medical School, Montpellier University, <sup>3</sup>Inserm U1183, St Eloi Hospital, <sup>4</sup>Medical Imaging Department and <sup>5</sup>Biostatistics Department, Montpellier University Hospital, Montpellier, France

Correspondence to: Goulabchand Radjiv, Médecine Interne et Maladies Multi-Organiques (MIMMO, Pr Le Quellec), CHRU Montpellier, Hôpital Saint-Eloi, 80 Avenue Augustin Fliche, 34295 Montpellier Cedex 05.  
E-mail: r-goulabchand@chu-montpellier.fr

## References

- 1 Betteridge Z, McHugh N. Myositis-specific autoantibodies: an important tool to support diagnosis of myositis. *J Intern Med* 2016;280:8–23.
- 2 Fujimoto M, Watanabe R, Ishitsuka Y, Okiyama N. Recent advances in dermatomyositis-specific autoantibodies. *Curr Opin Rheumatol* 2016;28:636–44.

**Fig. 1** Extensive subcutaneous calcinosis: arm and back pictures



Total body CT imaging reconstruction (bone-window settings).