

Systemic Lupus Erythematosus/Series Editors: D. Isenberg and C. Gordon

Systemic lupus erythematosus—messages from experimental models

M. L. Stoll and J. Gavalchin

Division of Hematology/Oncology, Department of Medicine, SUNY Health Science Center at Syracuse, Syracuse, NY, USA

Systemic lupus erythematosus (SLE) is an autoimmune disease characterized by the production of autoreactive T cells and autoantibodies that can affect virtually every organ system. The most severe clinical manifestations include immune complex-mediated glomerulonephritis, arthritis, vasculitis, cerebritis, pericarditis, cytopenias and serositis [1]. The diversity of clinical manifestations and disease phenotype, together with limited access to patient tissues, has made the study of human lupus difficult. However, the availability of a number of animal models for SLE has allowed us to make significant progress towards understanding the pathogenetic mechanisms contributing to disease development, and developing therapeutic strategies that specifically target the critical immune cells that are involved. This review will summarize some of the findings in the experimental murine models for SLE that have led to our current understanding of the pathogenetic mechanisms involved in the development of the disease in humans.

Experimental models for systemic lupus nephritis

Spontaneous models

Initial efforts to identify pathogenetic mechanisms in SLE utilized murine models that spontaneously developed syndromes resembling human lupus. The serological and clinical characteristics of these models are summarized in Table 1.

One of the first autoimmune models, the New Zealand black (NZB) mouse, was found to develop a spontaneous autoimmune disease resembling haemolytic anaemia, and only rarely, glomerulonephritis [2]. However, this strain does share some immunological features in common with SLE patients, including abnormal B-cell tolerance [3], intrinsic B-cell hyperproliferation and immunoglobulin production [4–6], loss of suppressor cell function [7], and defective apoptosis of resting B cells [8]. Also, like human lupus, there is a genetic component contributing to NZB disease, and linkage

studies have identified several chromosomal loci associated with nephritis and autoantibody production [9].

When the NZB mice are crossed with either the New Zealand White (NZW) or the Swiss-Webster (SWR) strains, the female F₁ progeny, referred to as B/W or SNF₁, respectively, develop a lethal accelerated glomerulonephritis, due to the deposition of immune complexes, including anti-DNA Ig [4, 17–20]. Since B/W and SNF₁ disease resemble human lupus very closely, in both the immunological features of the disease and increased incidence of disease in females, studies in these models may identify the critical factors contributing to the development of human SLE. Some of the immunological features that are observed in the NZB mice, such as a loss of suppressor cell function and polyclonal B-cell activation, are similarly seen in their B/W and SNF₁ offspring [6, 7, 21, 22]. Genetic loci associated with susceptibility to renal disease have been identified. Carlsten and Tarkowski [23] mated B/W mice (H2^{d,z}) with the NZB parent (H2^d) and found that B/W mice with the H2^{d,d} haplotype, relative to their d,z siblings, had increased survival, decreased renal disease, and decreased levels of serum IgG + M and IgG against ss- and dsDNA, demonstrating a role for the major histocompatibility complex (MHC) in the development of the disease. However, the importance of class II MHC genes for lupus susceptibility was questioned recently by the results of studies in mice transgenic for E^z [24] or A^z [25] genes, which showed that production of autoantibodies was not linked with inheritance of the transgenes. Non-MHC loci from NZB or NZW mice have also been linked to various lupus traits, such as nephritis and autoantibody production. For instance, the *Nba1* locus on chromosome 4 has been associated with nephritis, as has the *Nba2* locus on chromosome 1. The same region of chromosome 1 also contains the *Sle1* and *Lbw-7* loci, and all three are linked to the production of autoantibodies, particularly IgG antibodies reactive with chromatin [9, 26–28]. Recently, this region in humans, specifically 1q41-q42, showed evidence of linkage with SLE, including the production of anti-chromatin antibodies [29]. Thus, important susceptibility genes for autoimmunity appear to be conserved between mice and humans [30], supporting approaches to look for candidate human SLE-associated genes based on mouse studies.

Accepted 20 August 1999.

Correspondence to: J. Gavalchin, Department of Medicine, SUNY HSC at Syracuse, 750 East Adams Street, Syracuse, NY 13210, USA.

TABLE 1. Murine models of SLE

	Mean survival (weeks)	Clinically relevant autoantibodies	Sensitivity to exogenous sex hormones	Clinical features
NZB	Males: 67 Females: 63	Anti-erythrocyte Anti-ssDNA Anti-dsDNA (low) NTA	None	Haemolytic anaemia Mild glomerulonephritis
(NZB × NZW) _{F1}	Males: 58 Females: 35	Anti-dsDNA NTA	Oestrogens accelerate; androgens protect	Severe glomerulonephritis
(SWR × NZB) _{F1}	Males: 64 Females: 29	Anti-dsDNA Anti-ssDNA	Oestrogens accelerate	Severe glomerulonephritis
(SNF ₁)	Males: 22 Females: 20	Anti-dsDNA Anti-Sm RF	Controversial; oestrogens may slightly accelerate disease	Severe glomerulonephritis, synovitis, polyarteritis lymphadenopathy, etc.
MRL/ <i>lpr</i>		Anti-Sm	Controversial	Late-onset glomerulonephritis
MRL/+	Males: 98 Females: 73	Anti-ssDNA Anti-dsDNA Anti-erythrocyte	None	Severe glomerulonephritis, haemolytic anaemia, lymphadenopathy, vasculitis
BXSB/MP	Males: 20 Females: 68			Severe glomerulonephritis, pneumonitis, alopecia
Motheaten	Males: 3.2 Females: 3.2	Anti-ssDNA Anti-dsDNA	Not tested	
Viable motheaten	Males: 10 Females: 10			

NTA, natural thymotoxic antibody; RF, rheumatoid factor.

Sources: [10–16]; M. L. Stoll, unpublished data.

Several other murine models that spontaneously develop a lupus-like syndrome are also available. However, in these models, genes have been identified that secondarily accelerate lupus. The MRL/Mp-*+/+* (MRL/+) and MRL/Lp-*lpr/lpr* (MRL/*lpr*) autoimmune models differ phenotypically by the development of early onset autoimmunity in the MRL/*lpr* mice, attributable to a single gene, *lpr* [11], which was later found to code for the Fas receptor for apoptosis [31]. This finding stimulated intensive interest in a possible role for the Fas protein in human lupus, but the results, as discussed below, have been ambiguous. When crossed on to different backgrounds, the *lpr* gene causes a variety of clinical and serological abnormalities, with most strains developing some degree of autoantibody production and lymphoproliferation, mainly consisting of CD3+ CD4- CD8- B220+ T cells [12, 32], but not the severe renal disease observed in the MRL/*lpr* strain [11]. This finding and others [33, 34], suggests that MRL background genes also play a role in the development of lupus. Interestingly, in MRL/*lpr* mice, there is less evidence for an MHC association with the disease, as selective matings have revealed that the disease occurs with other MHC haplotypes [24]. Recently, several additional chromosomal loci have been discovered that, independent of Fas expression, conferred increased susceptibility to autoantibody production and vasculitis [35].

A unique feature of the BXSB/Mp strain is that disease onset occurs at a much earlier age in males compared with females [11, 36]. This trait was mapped to one or more genes on the Y chromosome, and was not associated with male reproductive hormones, as castration of pre-pubertal males did not significantly

affect disease outcome [36]. This as yet unidentified gene has been designated *Yaa* (Y chromosome-linked autoimmune acceleration). As with the *lpr* gene [11, 12], the phenotype of male mice bearing the *Yaa* gene also depends on background genes; for instance, C57BL/6 males to whom the *Yaa* chromosome was transferred did not develop autoimmunity [37]. Using backcrosses of the lupus-prone (NZW × C57BL/6.Yaa)_{F1} males with the wild-type C57BL/6 strain, Santiago and coworkers [38] detected loci on three separate autosomal chromosomes that contributed to the disease, supporting the requirement of additional contributory genes besides *Yaa* in the development of autoimmunity in BXSB mice.

Mice that are homozygous for mutations at the 'motheaten' locus (*me*) and the allelic variant 'motheaten viable' (*me^v*) develop a severe combined syndrome with profound systemic autoimmunity characterized by abnormalities in natural killer cell function and differentiation, as well as macrophage/monocyte morphology and function. The mutation is located in the gene for protein tyrosine phosphatase 1C (PTP1-c), also known as haematopoietic cell phosphatase (HCP), an enzyme involved in multiple signalling pathways in B- and T-cell haematopoiesis [39]. However, recently it was reported that the presence of T and B cells was not required for the motheaten phenotype, except for autoantibody production, calling into question the use of this strain as a paradigm for human autoimmune disease [14].

Induced models

C3H [40] and Balb/c mice [41] have been reported to develop a lupus-like syndrome when injected at 2 months of age with an antibody bearing the 16/6 idiotype, which is frequently detected in humans with SLE [42, 43].

Antibodies reactive with ss- and dsDNA, as well as the 16/6 idiotype are produced, and the mice ultimately develop a lethal glomerulonephritis. T cells that induced production of the 16/6 idiotype in an MHC-restricted manner were found to be involved [44, 45]. In other studies, T cells which inhibited *in vivo* production of Ig bearing the 16/6 idiotype and inhibited disease development in mice immunized with the 16/6 idiotype were identified [46]. Thus, these studies appear to support a role for idiotypic network interactions in autoimmunity. However, it has been difficult for other laboratories to reproduce this model [47], with environmental factors in specific laboratories speculated to be responsible, as has been shown in the murine models for collagen arthritis [47], diabetes [48] and multiple sclerosis [49].

In selected parent \rightarrow F₁ combinations, graft-*vs*-host disease (GVHD) can be induced by injection of parental lymphocytes into adult non-irradiated F₁ hybrids, for example, (C57BL/10 \times DBA/2)F₁ mice injected with DBA/2 cells. The disease is characterized by a chronic autoimmune syndrome resembling human SLE, with hypergammaglobulinaemia, production of IgG antibodies to a variety of autoantigens, as well as antibody-mediated glomerulonephritis [50, 51]. T_H2-type cells from the donor appear to participate in the pathogenetic process, acting on host F₁ B cells, perhaps through the production of interleukin-4 (IL-4) and IL-5 [52, 53].

When some strains of mice are injected with pristane, a branched alkane routinely used to prime mice for hybridoma implantation, SLE-like changes in the glomerulus, with deposition of immunoglobulin and complement, are observed. Autoantibodies are produced that react with ssDNA and histones, but not dsDNA [54]. The genetic background, particularly at H-2, influences serology in this model; anti-DNA autoantibodies are produced in Balb/c mice (H-2^d) with pristane immunization, while SJL mice (H-2^s) produce antinuclear ribonucleoprotein antibodies [54].

Transgenic and knockout models

Transgenic and knockout models for specific genes that have been proposed to have a role in the pathogenesis of autoimmunity permit direct testing of their significance in disease. These genes include those for cytokines, T cell or Ig receptors, MHC, or co-stimulatory molecules. However, these new animal models for disease are limited by the assumptions that they make regarding the importance of a specific gene in disease pathogenesis. Thus, the results of studies using these mice must be validated by observations in spontaneous murine models and human disease. Examples of some of the 'knockout' mice currently under study include transforming growth factor-beta (TGF- β 1) $-/-$ mice, which develop serum antibodies that are reactive with dsDNA, ssDNA and the Sm ribonucleoprotein, resulting in Ig glomerular deposits [55]. Autoimmune disease also occurs in mice with deleted IL-2 [56] and IL-10 [57], CTLA-4 [58], and T cell receptor (TCR) genes [59]. In contrast, autoimmune mice deficient in CD4 [60] or terminal deoxynucleotidyl transferase (TdT) [61] show dimin-

ished autoimmunity. Polymorphisms in some human genes have been linked to SLE, and studies in mice deficient in these genes have provided insight into their role in disease pathogenesis. One example is the human IgG receptor (Fc γ R) genes, particularly Fc γ RIIa and Fc γ RIIIa [62–64]. Polymorphisms in these genes may be important risk factors for SLE nephritis, and differences in clinical manifestations could result from the genetically determined ability to bind and clear immune complexes. Moreover, studies in murine models suggest that Fc γ R receptors are necessary for nephritis, in addition to the formation of immune complexes. Clynes *et al.* [65] crossed B/W mice with a knockout strain lacking Fc γ RI and Fc γ RIII receptors (γ - $-/-$). These mice developed anti-DNA antibodies, and immune complexes were deposited in the glomeruli. However, inflammatory nephritis and renal failure did not occur, even though complement was deposited, indicating a requirement for Fc γ R in disease initiation. Studies in animal models have also clarified the mechanism by which C1q deficiency contributes to the development of autoimmunity. C1q deficiency has been associated with lupus in over 90% of the known occurrences of this deficiency, and defects in C1q are found in lupus patients [66]. These observations suggested that C1q plays a role in protecting against the development of SLE [67], and this hypothesis was tested in C1q-deficient mice (C1qa $-/-$). These mice developed significant titres of autoantibodies, and 25% of them had glomerulonephritis associated with immune complex deposition, and multiple apoptotic cell bodies, implying either that increased apoptosis and/or a defect in the clearance of apoptotic cells contributed to the development of autoimmunity. Cells undergoing apoptosis produce subcellular structures (blebs), containing nuclear and cytoplasmic constituents, and thus may be the source of antigen driving the production of autoantibodies in SLE [68, 69]. The data in C1q-deficient mice suggest that one way that C1q may protect from SLE is by binding to and promoting the clearance of apoptotic cells, thus removing the autoantigenic stimulus.

Transgenic models include mice transgenic for human *BCL2*, an oncogene that promotes cell survival; these mice develop a disease characterized by the production of autoantibodies and kidney disease, and lymphocytic infiltration of arteries and solid organs [70]. Transgenic mice expressing interferon- γ (IFN- γ) in the epidermis produced autoantibodies reactive with dsDNA and histones, and further, the levels of these antibodies were higher in females *vs* males, as is found in human SLE [71]. Mice transgenic for autoantibody heavy chain genes show that autoantibody-producing B cells and autoreactive T cells are strictly regulated in non-autoimmune mice [72, 73].

Immunopathogenesis of SLE

A number of immunological abnormalities have been noted in both lupus patients and murine models. An essential component of murine lupus is autoreactive

B cells, which are found at relatively higher numbers, and are substantially activated, compared with non-autoimmune mice [22, 74]. These B cells produce a variety of autoantibodies [75–79], including antibodies to dsDNA, which are virtually diagnostic of lupus, as they are both highly sensitive and specific for the disease [76], and in general correlate with nephritis in human lupus patients [76, 77, 80]. This has led to the hypothesis that DNA may be the eliciting antigen in SLE [81–83]. On the other hand, in both humans and mice, the correlation between nephritis and anti-DNA antibody levels is not universal [77], and the polyreactivity of many anti-DNA antibodies further undermines this relationship [84]. In addition, numerous investigators have observed that modulation of disease outcome in murine lupus need not have a corresponding effect on the levels of anti-DNA antibodies [13, 85–89]. Moreover, the existence of somatically mutated and class-switched anti-DNA immunoglobulins is highly indicative of T-cell help (see, for example [90]). Furthermore, DNA has been shown to be poorly immunogenic in laboratory animals, even those predisposed to autoimmunity [91, 92], reviewed in [93]. Still, the relevance of anti-DNA antibodies to the pathogenetic process in SLE is clearly evident in studies using LPJ-394, a synthetic immunomodulant that selectively targets B cells producing anti-dsDNA antibodies [94]. In BXSB mice, treatment with LPJ-394 led to a reduction in serum levels of anti-DNA antibodies, reduced proteinuria, and increased survival [94], and similar reductions in anti-dsDNA antibody levels have been noted in SLE patients receiving LPJ-394 therapy [95].

Although the exact mechanisms involved in the pathogenesis of lupus nephritis are as yet unclear, there has been general agreement that disease is mediated by glomerular deposition of autoantibodies as immune complexes formed *in situ* or by direct binding of the antibodies to an intrinsic renal antigen or a self-antigen deposited in the kidney [96]. This deposited immunoglobulin then induces renal injury primarily through complement fixation, and recruitment and activation of inflammatory mediators [97]. However, a recent study suggested that the production of autoantibodies was not required for renal disease; MRL/MpJ-Fas*lpr* (MRL/*lpr*) mice that expressed a mutant transgene that did not permit secretion of circulating Ig still developed renal disease, suggesting that Ig-expressing B cells might also participate in the disease process either as antigen-presenting cells or as part of the local inflammatory process [98].

Evidence from studies of newly diagnosed lupus patients who also have an acute Epstein–Barr virus (EBV) infection have implicated this virus in the pathogenesis of lupus [99–101]. Work in mice has corroborated these findings; a peptide from the Sm RNP (PPPGMRPP) was found to be similar to a peptide from the Epstein–Barr nuclear antigen (EBVNA) (PPPGRRP) with the latter peptide reacting with lupus autoantibodies, but not antibodies from EBV-infected individuals who did not have autoimmune disease [102].

Immunization of normal rabbits and certain inbred mouse strains with the EBVNA peptide by James *et al.* [103, 104], resulted in the production of antibodies that cross-reacted with the Sm peptide, and clinical disease. These studies suggested that autoimmune disease might develop as a result of B- and T-cell responses to a particular antigen, such as EBV, that diversify over time, so that eventually antibodies may also be directed against host proteins, such as Sm. This autoreactivity can potentially spread to other epitopes within the cross-reactive protein and even to other proteins, a concept known as epitope spreading (reviewed in [105–107]). However, this effect has not always been seen by other investigators; attempts to reproduce it independently have found limited epitope spreading but not autoimmune disease [108]. These data again illustrate the difficulties that may be encountered in reproducing and interpreting results in some animal models.

T cells, particularly $\alpha\beta$ CD4⁺ cells, are considered to be essential for the development of lupus, by inducing isotype switching and affinity maturation [109–112]. The derivation of $\alpha\beta$ CD4⁺ T-helper cell lines from mice that are capable of inducing lupus nephritis *in vivo* [13, 113, 114], as well as the beneficial effects of therapeutic strategies targeting T cells, like neonatal thymectomy [115], and the treatment of mice [116] and humans with SLE with anti-CD4 antibody [117] (reviewed in [118]) further support the role of $\alpha\beta$ CD4⁺ cells in disease pathogenesis. The importance of CD4⁺ T cells in disease was also clearly supported by the finding that anti-dsDNA antibody production was decreased in CD4-deficient MRL/*lpr* mice. However, they did eventually develop moderate nephritis [60]. One way that T cells mediate their effect may be via increased production of cytokines; in some models like GVHD, and in lupus patients, a T_H2 pattern of cytokine production is observed [119], while in others, for example, NZB/NZW and MRL/*lpr* mice, the pattern is more complex, with increases seen in IL-4, IL-6 and IL-10 [120]. Deficient production of IL-6, tumour necrosis factor- α (TNF- α), IL-1 and IL-12 has also been noted [121, 122]. Nakajima *et al.* [123] observed clinical improvement with the administration of IL-4, whereas others have seen improvement with TNF- α [124, 125]. In non-autoimmune mice, transgenic expression of IL-4 results in autoantibody production and immune complex glomerulonephritis [126]. However, disease is ameliorated in (NZB \times C57BL/6.Yaa)F₁ transgenic mice overexpressing IL-4 in B cells [127]. CD4⁺ T cells have also been proposed to cause renal damage directly, since they may be found in the interstitium and glomeruli of kidneys from SLE patients [128] and mice with nephritis (Silvin *et al.*, in preparation). Finally, $\gamma\delta$ T cells may play a role, through their ability to regulate both $\alpha\beta$ T-cell and B-cell autoreactivity [129–131].

Recent studies from the B/W model have attempted to reconcile the observations that autoantibody production is associated with polyclonal B- and T-cell activation, but the autoantibodies appear to result from specific antigen-driven responses [132, 133]. These studies tested

the hypothesis that self-peptides derived from the VH region of anti-DNA antibodies were ligands for autoreactive T cells. In support of this, it was noted that some of these peptides increased the production of IgG anti-DNA antibodies *in vitro* [134], immunization with these peptides accelerated autoantibody production and nephritis [135, 136], and adoptive transfer of T-cell clones reactive with these peptides accelerated nephritis [136]. Finally, immune tolerance induction to three of these peptides in pre-nephritic B/W mice delayed the development of IgG anti-DNA antibodies and significantly improved survival [137]. Multiple mechanisms might contribute to the ability of these peptides to propagate an autoimmune response, including reciprocal activation of T and B cells by autoantibodies leading to the continuous recruitment of neo-autoreactive T helper cells, and pathogenic autoimmunity in a genetically predisposed individual [134]. We have obtained similar results in the SNF₁ cross. Initially, we found that modulation of the production of IgG expressing a pathogenic idiotype associated with nephritis, Id^{LN}F₁ + , by anti-idiotypic antibody or Id-reactive T-cell clones, altered the time of disease onset [13, 87, 138, 139]. We identified a peptide derived from a pathogenic Id^{LN}F₁ + antibody that stimulated these T-cell clones, and found that it was structurally similar to one of the peptides relevant in B/W disease [135–137], as each contained a triple basic amino acid motif, KKKKK. Recently, a peptide with the same motif derived from a natural autoantibody was also shown to modulate B/W disease [140]. Since T cells from patients with SLE may also be activated by VH peptides [141], similar mechanisms are likely to be critical to the development of human lupus and therapeutic regimens using autoantibodies or peptide determinants derived from them as tolerogens might therefore be useful in the treatment of human SLE. In support of this, treatment of MRL/*lpr* mice with dsDNA–anti-dsDNA immune complexes increased their survival, perhaps due to anti-idiotypic regulation of pathogenic antibody production [142]. Furthermore, an idiotypic vaccine that was derived from an anti-dsDNA antibody from MRL/*lpr* mice, 3E10, showed some protective effects in murine lupus, and was found to stabilize disease activity in some lupus patients [143].

Another feature of the pathogenic autoantibodies of SNF₁ mice and other murine SLE models is that they are cationic, suggesting that immune complexes containing these antibodies may initiate lupus nephritis by binding to negatively charged proteoglycans in the glomerular basement membrane [110, 144–146]. B cells producing these antibodies most likely process and present DNA complexed with cationic protein, such as histones, to pathogenic T cells that are specific for nucleosomal antigen(s) presented by either I-A^d or I-A^g. These T helper clones can then provide help to B cells which produce autoantibodies to ssDNA, dsDNA and histones [147, 148], thus nucleosome-specific T helper cells play an important role in initiating and sustaining the production of pathogenic antinuclear antibody production in

murine [147, 149] and human lupus [150, 151]. In further support for the role of nucleosomes as the target of autoreactive T cells in the development of lupus nephritis, tolerogenic therapy with nucleosome-derived peptides was recently found not only to delay the onset of nephritis in pre-nephritic SNF₁ mice, but was also effective in halting the progression of established disease [152].

Studies in the SNF₁ model have also established the importance of the contact-dependent, cognate interaction between pathogenic T helper and B cells in lupus nephritis. Antibody to CD40L (gp39), which transduces a co-stimulatory signal required for B-cell growth and differentiation, prevented pathogenic T helper clones and primary cells from inducing the production of autoantibodies by syngeneic B cells [153]. Treatment of SNF₁ mice with antibody to CD40L prior to the expression of elevated levels of IgG autoantibodies delayed the development of lupus nephritis [153], and was also found to be beneficial for the treatment of established disease [154]. Since disease mechanisms in SNF₁ mice and lupus patients are similar, and increased expression of CD40L has been noted in SLE patient T cells [155, 156], it is highly likely that anti-CD40L therapy will be effective for human lupus, and recruitment of patients for clinical trials is now under way [157]. However, other data examining the role of CD40–CD40L interactions in CD40L-deficient *lpr/lpr* mice suggested that this interaction was not an absolute requirement for class switching and that the production of IgG autoantibodies may arise by both CD40L-dependent and -independent pathways, although renal disease appears to be CD40L-dependent [158]. Studies in murine models have also shown that blocking other co-stimulatory interactions such as CTLA-4/B7, may be a useful therapeutic approach for SLE. In B/W mice, administration of murine CTLA-4Ig blocked autoantibody production, suppressed nephritis, and increased survival [159], and clinical trials in SLE patients are planned [160].

As discussed above, the *lpr* gene of MRL/*lpr* mice was found to encode for an apoptosis signalling receptor [31], suggesting that abnormalities in lymphocyte apoptosis might play a role in the development of lupus (reviewed in [161]). Supporting data include: (1) increased levels of soluble Fas in the serum of human lupus patients [162, 163]; and (2) the association of a rare lupus-like autoimmune condition with the absence of Fas [164]. On the other hand, the initial study associating high soluble Fas (sFas) levels with SLE looked at a very small group of patients, and only six of 10 had high sFas levels [162]. Later studies confirmed that elevated levels of sFas were not consistently found in lupus patients and do not correlate with disease activity [165, 166]. Furthermore, some investigators have found normal or high levels of Fas expression on lymphocytes derived from lupus patients [167] and increased levels of *in vitro* apoptosis in patients with SLE [168]. However, while MRL/*lpr* disease is largely associated with deficient apoptosis *in vivo* [31], apoptosis among lymphoid cells *in vitro* is increased [169]. Thus,

the role of sFas in SLE requires further study, with the data suggesting that perhaps a subgroup of SLE patients may have structural Fas or Fas ligand (FasL) defects, that contribute to their disease.

Conclusion

Contributions to the development of SLE in humans include multiple genetic factors such as intrinsic immunological abnormalities, MHC genes, sex hormones, and environmental factors such as infectious agents, among others, with a complex network involved in its regulation. Although no single murine model for the disease encompasses all of these factors, we are fortunate that each of these models possesses many of the features of human SLE, enabling investigators to address specifically the relevance of contributory agents and identify specific immunoregulatory defects that may predispose to the disease. In addition, the use of murine models facilitates the development and testing of immunotherapies that may specifically downregulate the pathogenic immune processes without the use of immunosuppressive agents.

References

- Boumpas DT, Fessler BJ, Austin HA III, Balow JE, Klippel JH, Lockshin MD. Systemic lupus erythematosus: emerging concepts. Part 2: Dermatologic and joint disease, the antiphospholipid antibody syndrome, pregnancy and hormonal therapy, morbidity and mortality, and pathogenesis. *Ann Intern Med* 1995;123:42–53.
- Bielschowsky M. Spontaneous anemia in mice of the NZB/ml strain. *Proc Univ Otago Med Sch* 1959;37:9–11.
- Wither J, Vukusic B. Autoimmunity develops in lupus-prone NZB mice despite normal T cell tolerance. *J Immunol* 1998;161:4555–62.
- Manny N, Datta SK, Schwartz RS. Synthesis of IgM by cells of NZB and SWR mice and their crosses. *J Immunol* 1979;122:1220–7.
- Reininger L, Winkler TH, Kalberer CP, Jourdan M, Melchers F, Rolink AG. Intrinsic B cell defects in NZB and NZW mice contribute to systemic lupus erythematosus in (NZB × NZW)_{F1} mice. *J Exp Med* 1996;184:853–61.
- Jongstra-Bilen J, Vukusic B, Boras K, Wither JE. Resting B cells from autoimmune lupus-prone New Zealand Black and (New Zealand Black × New Zealand White)_{F1} mice are hyper-responsive to T cell-derived stimuli. *J Immunol* 1997;159:5810–20.
- Talal N, Steinberg AD. The pathogenesis of autoimmunity in New Zealand black mice. *Curr Top Microbiol Immunol* 1974;64:79–103.
- Kozono Y, Kotzin BL, Holers VM. Resting B cells from New Zealand Black mice demonstrate a defect in apoptosis induction following surface IgM ligation. *J Immunol* 1996;156:4498–503.
- Vyse TJ, Drake CG, Rozzo SJ, Roper E, Izui S, Kotzin BL. Genetic linkage of IgG autoantibody production in relation to lupus nephritis in New Zealand hybrid mice. *J Clin Invest* 1996;98:1762–72.
- Andrews BS, Eisenberg RA, Theofilopoulos AN *et al*. Spontaneous murine lupus-like syndromes. Clinical and immunopathological manifestations in several strains. *J Exp Med* 1978;148:1198–215.
- Theofilopoulos AN, Dixon FJ. Murine models of systemic lupus erythematosus. *Adv Immunol* 1985;37:269–390.
- Cohen PL, Eisenberg RA. Lpr and gld: single gene models of systemic autoimmunity and lymphoproliferative disease. *Annu Rev Immunol* 1991;9:243–69.
- Knapp CJ, Uner AH, Tatum AH, Kakanar JR, Gavalchin J. IdLNF1-specific T cell clones accelerate the production of IdLNF1+IgG and nephritis in SNF1 mice. *J Autoimmun* 1995;8:367–80.
- Yu CC, Tsui HW, Ngan BY, Shulman MJ, Wu GE, Tsui FW. B and T cells are not required for the viable motheaten phenotype. *J Exp Med* 1996;183:371–80.
- Westhoff CM, Whittier A, Kathol S *et al*. DNA-binding antibodies from viable motheaten mutant mice: implications for B cell tolerance. *J Immunol* 1997;159:3024–33.
- Shultz LD, Coman DR, Bailey CL, Beamer WG, Sidman CL. 'Viable motheaten', a new allele at the motheaten locus. I. Pathology. *Am J Pathol* 1984;116:179–92.
- Helyer BJ, Howie JB. Renal disease associated with positive lupus-erythematosus tests in a cross-bred strain of mice. *Nature* 1963;197:197.
- Lambert PH, Dixon FJ. Pathogenesis of the glomerulonephritis of NZB/W mice. *J Exp Med* 1968;127:507–22.
- Datta SK, Manny N, Andrzejewski C, Andre-Schwartz J, Schwartz RS. Genetic studies of autoimmunity and retrovirus expression in crosses of New Zealand black mice. I. Xenotropic virus. *J Exp Med* 1978;147:854–71.
- Gavalchin J, Nicklas JA, Eastcott JW *et al*. Lupus prone (SWR × NZB)_{F1} mice produce potentially nephritogenic autoantibodies inherited from the normal SWR parent. *J Immunol* 1985;134:885–94.
- Cohen PL, Ziff M. Abnormal polyclonal B cell activation in NZB/NZW _{F1} mice. *J Immunol* 1977;119:1534–7.
- Klinman DM, Steinberg AD. Systemic autoimmune disease arises from polyclonal B cell activation. *J Exp Med* 1987;165:1755–60.
- Carlsten H, Tarkowski A. Histocompatibility complex gene products and exposure to oestrogen: two independent disease accelerating factors in murine lupus. *Scand J Immunol* 1993;38:341–7.
- Vyse TJ, Kotzin BL. Genetic susceptibility to systemic lupus erythematosus. *Annu Rev Immunol* 1998;16:261–92.
- Rozzo SJ, Vyse TJ, David CS, Palmer E, Izui S, Kotzin BL. Analysis of MHC class II genes in the susceptibility to lupus in New Zealand mice. *J Immunol* 1999;162:2623–30.
- Kono DH, Burlingame RW, Owens DG *et al*. Lupus susceptibility loci in New Zealand mice. *Proc Natl Acad Sci USA* 1994;91:10168–72.
- Mohan C, Alas E, Morel L, Yang P, Wakeland EK. Genetic dissection of SLE pathogenesis. Sle1 on murine chromosome 1 leads to a selective loss of tolerance to H2A/H2B/DNA subnucleosomes. *J Clin Invest* 1998;101:1362–72.
- Morel L, Yu Y, Blenman KR, Caldwell RA, Wakeland EK. Production of congenic mouse strains carrying genomic intervals containing SLE-susceptibility genes derived from the SLE-prone NZM2410 strain. *Mamm Genome* 1996;7:335–9.
- Tsao BP, Cantor RM, Kalunian KC *et al*. Evidence for linkage of a candidate chromosome 1 region to human

- systemic lupus erythematosus. *J Clin Invest* 1997;99:725–31.
30. Becker KG, Simon RM, Bailey-Wilson JE *et al.* Clustering of non-major histocompatibility complex susceptibility candidate loci in human autoimmune diseases. *Proc Natl Acad Sci USA* 1998;95:9979–84.
 31. Nagata S, Golstein P. The Fas death factor. *Science* 1995;267:1449–56.
 32. Morse HC 3rd, Davidson WF, Yetter RA, Murphy ED, Roths JB, Coffman RL. Abnormalities induced by the mutant gene *lpr*: expansion of a unique lymphocyte subset. *J Immunol* 1982;129:2612–5.
 33. Kelley VE, Roths JB. Interaction of mutant *lpr* gene with background strain influences renal disease. *Clin Immunol Immunopathol* 1985;37:220–9.
 34. Pisetsky DS, McCarty GA, Peters DV. Mechanisms of autoantibody production in autoimmune MRL mice. *J Exp Med* 1980;152:1302–10.
 35. Gu L, Weinreb A, Wang XP *et al.* Genetic determinants of autoimmune disease and coronary vasculitis in the MRL-*lpr/lpr* mouse model of systemic lupus erythematosus. *J Immunol* 1998;161:6999–7006.
 36. Murphy ED, Roths JB. A Y chromosome associated factor in strain BXSB producing accelerated autoimmunity and lymphoproliferation. *Arthritis Rheum* 1979;22:1188–94.
 37. Izui S, Higaki M, Morrow D, Merino R. The Y chromosome from autoimmune BXSB/MpJ mice induces a lupus-like syndrome in (NZW \times C57BL/6) F_1 male mice, but not in C57BL/6 male mice. *Eur J Immunol* 1988;18:911–5.
 38. Santiago ML, Mary C, Parzy D *et al.* Linkage of a major quantitative trait locus to Yaa gene-induced lupus-like nephritis in (NZW \times C57BL/6) F_1 mice. *Eur J Immunol* 1998;28:4257–67.
 39. Tsui FW, Tsui HW. Molecular basis of the motheaten phenotype. *Immunol Rev* 1994;138:185–206.
 40. Mendlovic S, Brocke S, Shoenfeld Y *et al.* Induction of a systemic lupus erythematosus-like disease in mice by a common human anti-DNA idiotype. *Proc Natl Acad Sci USA* 1988;85:2260–4.
 41. Swissa M, Lerner A, Sasaki T, Sela E, Blank M, Shoenfeld Y. The diverse pathogenic potential of anti-DNA antibodies from various sources to induce experimental systemic lupus erythematosus. *Pathobiology* 1996;64:32–9.
 42. Villarreal GM, Drenkard C, Villa AR *et al.* Prevalence of 13 autoantibodies and of the 16/6 and related pathogenic idiotype in 465 patients with systemic lupus erythematosus and their relationship with disease activity. *Lupus* 1997;6:425–35.
 43. Isenberg DA, Shoenfeld Y, Madaio MP *et al.* Anti-DNA antibody idiotype in systemic lupus erythematosus. *Lancet* 1984;2:417–22.
 44. Fricke H, Mendlovic S, Blank M, Shoenfeld Y, Ben-Bassat M, Mozes E. Idiotype specific T-cell lines inducing experimental systemic lupus erythematosus in mice. *Immunology* 1991;73:421–7.
 45. Blank M, Mendlovic S, Mozes E, Coates AR, Shoenfeld Y. Induction of systemic lupus erythematosus in naive mice with T-cell lines specific for human anti-DNA antibody SA-1 (16/6 Id+) and for mouse tuberculosis antibody TB/68 (16/6 Id+). *Clin Immunol Immunopathol* 1991;60:471–83.
 46. Blank M, Ben-Bassat M, Shoenfeld Y. Modulation of SLE induction in naive mice by specific T cells with suppressor activity to pathogenic anti-DNA idiotype. *Cell Immunol* 1991;137:474–86.
 47. Isenberg DA, Katz D, Le Page S *et al.* Independent analysis of the 16/6 idiotype lupus model. A role for an environmental factor? *J Immunol* 1991;147:4172–7.
 48. Leiter EH, Serreze DV, Prochazka M. The genetics and epidemiology of diabetes in NOD mice [news]. *Immunol Today* 1990;11:147–9.
 49. Teuscher C, Hickey WF, Grafer CM, Tung KS. A common immunoregulatory locus controls susceptibility to actively induced experimental allergic encephalomyelitis and experimental allergic orchitis in BALB/c mice. *J Immunol* 1998;160:2751–6.
 50. Gleichmann H, Gleichmann E. GVHD, a model for Spanish toxic oil syndrome? [Letter]. *Lancet* 1984;1:1474.
 51. Rolink AG, Radaszkiewicz T, Melchers F. The autoantigen-binding B cell repertoires of normal and of chronically graft-versus-host-diseased mice. *J Exp Med* 1987;165:1675–87.
 52. Dobashi K, Ono S, Murakami S, Takahama Y, Katoh Y, Hamaoka T. Polyclonal B cell activation by a B cell differentiation factor, B151-TRF2. III. B151-TRF2 as a B cell differentiation factor closely associated with autoimmune disease. *J Immunol* 1987;138:780–7.
 53. Dautrelepont JM, Moser M, Leo O *et al.* Hyper IgE in stimulatory graft-versus-host disease: role of interleukin-4. *Clin Exp Immunol* 1991;83:133–6.
 54. Satoh M, Kumar A, Kanwar YS, Reeves WH. Anti-nuclear antibody production and immune-complex glomerulonephritis in BALB/c mice treated with pristane. *Proc Natl Acad Sci USA* 1995;92:10934–8.
 55. Dang H, Geiser AG, Letterio JJ *et al.* SLE-like autoantibodies and Sjögren's syndrome-like lymphoproliferation in TGF-beta knockout mice. *J Immunol* 1995;155:3205–12.
 56. Sadlack B, Merz H, Schorle H, Schimpl A, Feller AC, Horak I. Ulcerative colitis-like disease in mice with a disrupted interleukin-2 gene. [See comments] *Cell* 1993;75:253–61.
 57. Kuhn R, Lohler J, Rennick D, Rajewsky K, Muller W. Interleukin-10-deficient mice develop chronic enterocolitis. [See comments] *Cell* 1993;75:263–74.
 58. Marengere LE, Waterhouse P, Duncan GS, Mittrucker HW, Feng GS, Mak TW. Regulation of T cell receptor signaling by tyrosine phosphatase SYP association with CTLA-4. *Science* 1996;272:1170–3 (errata appeared in 274:1597 and 276:21).
 59. Mombaerts P, Mizoguchi E, Grusby MJ, Glimcher LH, Bhan AK, Tonegawa S. Spontaneous development of inflammatory bowel disease in T cell receptor mutant mice. [See comments] *Cell* 1993;75:274–82.
 60. Chesnutt MS, Finck BK, Killeen N, Connolly MK, Goodman H, Wofsy D. Enhanced lymphoproliferation and diminished autoimmunity in CD4-deficient MRL/*lpr* mice. *Clin Immunol Immunopathol* 1998;87:23–32.
 61. Conde C, Weller S, Gilfillan S, Marcellin L, Martin T, Pasquali JL. Terminal deoxynucleotidyl transferase deficiency reduces the incidence of autoimmune nephritis in (New Zealand Black \times New Zealand White) F_1 mice. *J Immunol* 1998;161:7023–30.
 62. Duits AJ, Bootsma H, Derksen RH *et al.* Skewed distribution of IgG Fc receptor IIa (CD32) polymorphism is associated with renal disease in systemic lupus erythematosus patients. *Arthritis Rheum* 1995;38:1832–6.
 63. Salmon JE, Millard S, Schachter LA *et al.* Fc gamma

- RIIA alleles are heritable risk factors for lupus nephritis in African Americans. *J Clin Invest* 1996;97:1348–54.
64. Wu J, Edberg JC, Redecha PB *et al.* A novel polymorphism of FcγRIIIa (CD16) alters receptor function and predisposes to autoimmune disease. *J Clin Invest* 1997;100:1059–70.
 65. Clynes R, Dumitru C, Ravetch JV. Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis. *Science* 1998;279:1052–4.
 66. Slingsby JH, Norsworthy P, Pearce G *et al.* Homozygous hereditary C1q deficiency and systemic lupus erythematosus. A new family and the molecular basis of C1q deficiency in three families. *Arthritis Rheum* 1996;39:663–70.
 67. Botto M, Dell'Agnola C, Bygrave AE *et al.* Homozygous C1q deficiency causes glomerulonephritis associated with multiple apoptotic bodies. [See comments] *Nature Genet* 1998;19:56–9.
 68. Casciola-Rosen LA, Anhalt G, Rosen A. Autoantigens targeted in systemic lupus erythematosus are clustered in two populations of surface structures on apoptotic keratinocytes. [See comments] *J Exp Med* 1994;179:1317–30.
 69. Rosen A, Casciola-Rosen L, Ahearn J. Novel packages of viral and self-antigens are generated during apoptosis. *J Exp Med* 1995;181:1557–61.
 70. Strasser A, Whittingham S, Vaux DL *et al.* Enforced BCL2 expression in B-lymphoid cells prolongs antibody responses and elicits autoimmune disease. *Proc Natl Acad Sci USA* 1991;88:8661–5.
 71. Seery JP, Carroll JM, Cattell V, Watt FM. Antinuclear autoantibodies and lupus nephritis in transgenic mice expressing interferon gamma in the epidermis. *J Exp Med* 1997;186:1451–9.
 72. Erikson J, Radic MZ, Camper SA, Hardy RR, Carmack C, Weigert M. Expression of anti-DNA immunoglobulin transgenes in non-autoimmune mice. *Nature* 1991;349:331–4.
 73. Shinde S, Gee R, Santulli-Marotto S, Bockenstedt LK, Clarke SH, Mamula MJ. T cell autoimmunity in Ig transgenic mice. *J Immunol* 1999;162:7519–24.
 74. Shlomchik MJ, Madaio MP, Ni D, Trounstein M, Huszar D. The role of B cells in lpr/lpr-induced autoimmunity. *J Exp Med* 1994;180:1295–306.
 75. Bell DA, Maddison PJ. Serologic subsets in systemic lupus erythematosus: an examination of autoantibodies in relationship to clinical features of disease and HLA antigens. *Arthritis Rheum* 1980;23:1268–73.
 76. Tomer Y, Buskila D, Shoenfeld Y. Pathogenic significance and diagnostic value of lupus autoantibodies. *Int Arch Allergy Immunol* 1993;100:293–306.
 77. Lefkowitz JB, Gilkeson GS. Nephritogenic autoantibodies in lupus: current concepts and continuing controversies. *Arthritis Rheum* 1996;39:894–903.
 78. Isshi K, Hirohata S. Association of anti-ribosomal P protein antibodies with neuropsychiatric systemic lupus erythematosus. *Arthritis Rheum* 1996;39:1483–90 (erratum appeared in 40:977).
 79. Cabral AR, Alarcon-Segovia D. Autoantibodies in systemic lupus erythematosus. *Curr Opin Rheumatol* 1997;9:387–92.
 80. Tan EM. Antinuclear antibodies: diagnostic markers for autoimmune diseases and probes for cell biology. *Adv Immunol* 1989;44:93–151.
 81. Karounos DG, Grudier JP, Pisetsky DS. Spontaneous expression of antibodies to DNA of various species origin in sera of normal subjects and patients with systemic lupus erythematosus. *J Immunol* 1988;140:451–5.
 82. Marion TN, Tillman DM, Jou NT. Interclonal and intracolon diversity among anti-DNA antibodies from an (NZB × NZW)F₁ mouse. *J Immunol* 1990;145:2322–32.
 83. Shlomchik M, Mascelli M, Shan H *et al.* Anti-DNA antibodies from autoimmune mice arise by clonal expansion and somatic mutation. *J Exp Med* 1990;171:265–92.
 84. Emlen W, Pisetsky DS, Taylor RP. Antibodies to DNA. A perspective. *Arthritis Rheum* 1986;29:1417–26.
 85. Oldstone MB, Dixon FJ. Inhibition of antibodies to nuclear antigen and to DNA in New Zealand mice infected with lactate dehydrogenase virus. *Science* 1972;175:784–6.
 86. Kubo C, Day NK, Good RA. Influence of early or late dietary restriction on life span and immunological parameters in MRL/Mp-lpr/lpr mice. *Proc Natl Acad Sci USA* 1984;81:5831–5.
 87. Uner AH, Knupp CJ, Tatum AH, Gavalchin J. Treatment with antibody reactive with the nephritogenic idiotype, IdLNF1, suppresses its production and leads to prolonged survival of (NZB × SWR)F₁ mice. *J Autoimmun* 1994;7:27–44.
 88. Gilkeson GS, Ruiz P, Pippen AM, Alexander AL, Lefkowitz JB, Pisetsky DS. Modulation of renal disease in autoimmune NZB/NZW mice by immunization with bacterial DNA. *J Exp Med* 1996;183:1389–97.
 89. Mittleman BB, Morse HC III, Payne SM, Shearer GM, Mozes E. Amelioration of experimental systemic lupus erythematosus (SLE) by retrovirus infection. *J Clin Immunol* 1996;16:230–6.
 90. Davidson A, Manheimer-Lory A, Aranow C, Peterson R, Hannigan N, Diamond B. Molecular characterization of a somatically mutated anti-DNA antibody bearing two systemic lupus erythematosus-related idiotypes. *J Clin Invest* 1990;85:1401–9.
 91. Madaio MP, Hodder S, Schwartz RS, Stollar BD. Responsiveness of autoimmune and normal mice to nucleic acid antigens. *J Immunol* 1984;132:872–6.
 92. Mor G, Singla M, Steinberg AD, Hoffman SL, Okuda K, Klinman DM. Do DNA vaccines induce autoimmune disease? *Hum Gene Ther* 1997;8:293–300.
 93. Pisetsky DS. Anti-DNA antibodies in systemic lupus erythematosus: a case of mistaken identity? [Editorial] *J Rheumatol* 1998;25:195–7.
 94. Weisman MH, Bluestein HG, Berner CM, de Haan HA. Reduction in circulating dsDNA antibody titer after administration of LJP 394. *J Rheumatol* 1997;24:314–8.
 95. Tumlin JA. Lupus nephritis: novel immunosuppressive modalities and future directions. *Semin Nephrol* 1999;19:67–76.
 96. Couser WG. Mediation of immune glomerular injury. *J Am Soc Nephrol* 1990;1:13–29.
 97. Vlahakos DV, Foster MH, Adams S *et al.* Anti-DNA antibodies form immune deposits at distinct glomerular and vascular sites. *Kidney Int* 1992;41:1690–700.
 98. Chan OT, Hannum LG, Haberman AM, Madaio MP, Shlomchik MJ. A novel mouse with B cells but lacking serum antibody reveals an antibody-independent role for B cells in murine lupus. *J Exp Med* 1999;189:1639–48.
 99. Bhimma R, Adhikari M, Coovadia HM. Epstein-Barr virus-induced systemic lupus erythematosus. [See comments] *S Afr Med J* 1995;85:899–900.
 100. Dror Y, Blachar Y, Cohen P, Livni N, Rosenmann E, Ashkenazi A. Systemic lupus erythematosus associated

- with acute Epstein-Barr virus infection. *Am J Kidney Dis* 1998;32:825-8.
101. James JA, Kaufman KM, Farris AD, Taylor-Albert E, Lehman TJ, Harley JB. An increased prevalence of Epstein-Barr virus infection in young patients suggests a possible etiology for systemic lupus erythematosus. *J Clin Invest* 1997;100:3019-26.
 102. James JA, Harley JB. Linear epitope mapping of an Sm B/B' polypeptide. *J Immunol* 1992;148:2074-9.
 103. James JA, Gross T, Scofield RH, Harley JB. Immunoglobulin epitope spreading and autoimmune disease after peptide immunization: Sm B/B'-derived PPPGMRPP and PPPGIRGP induce spliceosome autoimmunity. *J Exp Med* 1995;181:453-61.
 104. James JA, Harley JB. A model of peptide-induced lupus autoimmune B cell epitope spreading is strain specific and is not H-2 restricted in mice. *J Immunol* 1998;160:502-8.
 105. James JA, Harley JB. B-cell epitope spreading in autoimmunity. *Immunol Rev* 1998;164:185-200.
 106. McCluskey J, Farris AD, Keech CL *et al.* Determinant spreading: lessons from animal models and human disease. *Immunol Rev* 1998;164:209-29.
 107. Singh RR, Hahn BH. Reciprocal T-B determinant spreading develops spontaneously in murine lupus: implications for pathogenesis. *Immunol Rev* 1998;164:201-8.
 108. Mason LJ, Timothy LM, Isenberg DA, Kalsi JK. Immunization with a peptide of Sm B/B' results in limited epitope spreading but not autoimmune disease. *J Immunol* 1999;162:5099-105.
 109. Ando DG, Sercarz EE, Hahn BH. Mechanisms of T and B cell collaboration in the *in vitro* production of anti-DNA antibodies in the NZB/NZW F₁ murine SLE model. *J Immunol* 1987;138:3185-90.
 110. Datta SK, Patel H, Berry D. Induction of a cationic shift in IgG anti-DNA autoantibodies. Role of T helper cells with classical and novel phenotypes in three murine models of lupus nephritis. *J Exp Med* 1987;165:1252-68.
 111. Naiki M, Chiang BL, Cawley D *et al.* Generation and characterization of cloned T helper cell lines for anti-DNA responses in NZB.H-2bm12 mice. *J Immunol* 1992;149:4109-15.
 112. Sobel ES, Kakkanaiah VN, Kakkanaiah M, Cheek RL, Cohen PL, Eisenberg RA. T-B collaboration for autoantibody production in lpr mice is cognate and MHC-restricted. *J Immunol* 1994;152:6011-6.
 113. Adams S, Leblanc P, Datta SK. Junctional region sequences of T-cell receptor beta-chain genes expressed by pathogenic anti-DNA autoantibody-inducing helper T cells from lupus mice: possible selection by cationic autoantigens. *Proc Natl Acad Sci USA* 1991;88:11271-5.
 114. Mao C, Osman GE, Adams S, Datta SK. T cell receptor alpha-chain repertoire of pathogenic autoantibody-inducing T cells in lupus mice. *J Immunol* 1994;152:1462-70.
 115. Steinberg AD, Roths JB, Murphy ED, Steinberg RT, Raveche ES. Effects of thymectomy or androgen administration upon the autoimmune disease of MRL/Mp-lpr/lpr mice. *J Immunol* 1980;125:871-3.
 116. Wofsy D, Seaman WE. Reversal of advanced murine lupus in NZB/NZW F₁ mice by treatment with monoclonal antibody to L3T4. *J Immunol* 1987;138:3247-53.
 117. Santoro TJ, Portanova JP, Kotzin BL. The contribution of L3T4+ T cells to lymphoproliferation and autoantibody production in MRL-lpr/lpr mice. *J Exp Med* 1988;167:1713-8.
 118. Perosa F, Scudeletti M, Imro MA, Dammacco F, Luccarelli G, Indiveri F. Anti-CD4 monoclonal antibody (mAb) and anti-idiotypic mAb to anti-CD4 in the therapy of autoimmune diseases. *Clin Exp Rheumatol* 1997;15:201-10.
 119. Rus V, Svetic A, Nguyen P, Gause WC, Via CS. Kinetics of Th1 and Th2 cytokine production during the early course of acute and chronic murine graft-versus-host disease. Regulatory role of donor CD8+ T cells. *J Immunol* 1995;155:2396-406.
 120. Handwerker BS, Rus V, da Silva L, Via CS. The role of cytokines in the immunopathogenesis of lupus. *Springer Semin Immunopathol* 1994;16:153-80.
 121. Alleva DG, Kaser SB, Beller DI. Aberrant cytokine expression and autocrine regulation characterize macrophages from young MRL+/+ and NZB/W F₁ lupus-prone mice. *J Immunol* 1997;159:5610-9.
 122. Alleva DG, Kaser SB, Beller DI. Intrinsic defects in macrophage IL-12 production associated with immune dysfunction in the MRL+/+ and New Zealand Black/White F₁ lupus-prone mice and the Leishmania major-susceptible BALB/c strain. *J Immunol* 1998;161:6878-84.
 123. Nakajima A, Hirose S, Yagita H, Okumura K. Roles of IL-4 and IL-12 in the development of lupus in NZB/W F₁ mice. *J Immunol* 1997;158:1466-72.
 124. Jacob CO, McDevitt HO. Tumour necrosis factor-alpha in murine autoimmune 'lupus' nephritis. *Nature* 1988;331:356-8.
 125. Gordon C, Ranges GE, Greenspan JS, Wofsy D. Chronic therapy with recombinant tumor necrosis factor-alpha in autoimmune NZB/NZW F₁ mice. *Clin Immunol Immunopathol* 1989;52:421-34.
 126. Erb KJ, Ruger B, von Brevens M, Ryffel B, Schimpl A, Rivett K. Constitutive expression of interleukin (IL)-4 *in vivo* causes autoimmune-type disorders in mice. *J Exp Med* 1997;185:329-39.
 127. Santiago ML, Fossati L, Jacquet C, Muller W, Izui S, Reininger L. Interleukin-4 protects against a genetically linked lupus-like autoimmune syndrome. *J Exp Med* 1997;185:65-70.
 128. Caligaris-Cappio F, Bergui L, Tesio L, Ziano R, Camussi G. HLA-Dr+ T cells of the Leu 3 (helper) type infiltrate the kidneys of patients with systemic lupus erythematosus. *Clin Exp Immunol* 1985;59:185-9.
 129. Braun MY, Lowin B, French L, Acha-Orbea H, Tschopp J. Cytotoxic T cells deficient in both functional fas ligand and perforin show residual cytolytic activity yet lose their capacity to induce lethal acute graft-versus-host disease. *J Exp Med* 1996;183:657-61.
 130. Budd RC, Mixter PF. The origin of CD4-CD8-TCR alpha beta+ thymocytes: a model based on T-cell receptor avidity. *Immunol Today* 1995;16:428-31.
 131. Renno T, Hahne M, Tschopp J, MacDonald HR. Peripheral T cells undergoing superantigen-induced apoptosis *in vivo* express B220 and upregulate Fas and Fas ligand. *J Exp Med* 1996;183:431-7.
 132. Dziarski R. Autoimmunity: polyclonal activation or antigen induction? *Immunol Today* 1988;9:340-2.
 133. Radic MZ, Weigert M. Genetic and structural evidence for antigen selection of anti-DNA antibodies. *Annu Rev Immunol* 1994;12:487-520.
 134. Singh RR, Hahn BH, Tsao BP, Ebling FM. Evidence for multiple mechanisms of polyclonal T cell activation in murine lupus. *J Clin Invest* 1998;102:1841-9.
 135. Ebling FM, Tsao BP, Singh RR, Sercarz E, Hahn BH.

- A peptide derived from an autoantibody can stimulate T cells in the (NZB \times NZW) F_1 mouse model of systemic lupus erythematosus. *Arthritis Rheum* 1993;36:355–64.
136. Singh RR, Kumar V, Ebling FM *et al.* T cell determinants from autoantibodies to DNA can upregulate autoimmunity in murine systemic lupus erythematosus. *J Exp Med* 1995;181:2017–27.
 137. Singh RR, Ebling FM, Sercarz EE, Hahn BH. Immune tolerance to autoantibody-derived peptides delays development of autoimmunity in murine lupus. *J Clin Invest* 1995;96:2990–6.
 138. Gavalchin J, Datta SK. The NZB \times SWR model of lupus nephritis. II. Autoantibodies deposited in renal lesions show a distinctive and restricted idiotypic diversity. *J Immunol* 1987;138:138–48.
 139. Knupp CJ, Uner AH, Korthas C, Gavalchin J. Characterization of IdLNFI-specific T cell clones from the (NZB \times SWR) F_1 murine model for systemic lupus erythematosus. *Clin Immunol Immunopathol* 1993;68:273–82.
 140. Jouanne C, Avrameas S, Payelle-Brogard B. A peptide derived from a polyreactive monoclonal anti-DNA natural antibody can modulate lupus development in (NZB \times NZW) F_1 mice. *Immunology* 1999;96:333–9.
 141. Williams WM, Staines NA, Muller S, Isenberg DA. Human T cell responses to autoantibody variable region peptides. *Lupus* 1995;4:464–71.
 142. Burny W, Lebrun P, Cosyns JP, Saint-Remy JM. Treatment with dsDNA–anti-dsDNA antibody complexes extends survival, decreases anti-dsDNA antibody production and reduces severity of nephritis in MRL/lpr mice. *Lupus* 1997;6:4–17.
 143. Weisbart RH, Noritake DT, Wong AL, Chan G, Kacena A, Colburn KK. A conserved anti-DNA antibody idiotype associated with nephritis in murine and human systemic lupus erythematosus. *J Immunol* 1990;144:2653–8.
 144. Shivakumar S, Tsokos GC, Datta SK. T cell receptor alpha/beta expressing double-negative (CD4 $^-$ /CD8 $^-$) and CD4 $^+$ T helper cells in humans augment the production of pathogenic anti-DNA autoantibodies associated with lupus nephritis. *J Immunol* 1989;143:103–12.
 145. Ebling F, Hahn BH. Restricted subpopulations of DNA antibodies in kidneys of mice with systemic lupus. Comparison of antibodies in serum and renal eluates. *Arthritis Rheum* 1980;23:392–403.
 146. Gauthier VJ, Mannik M. A small proportion of cationic antibodies in immune complexes is sufficient to mediate their deposition in glomeruli. *J Immunol* 1990;145:3348–52.
 147. Mohan C, Adams S, Stanik V, Datta SK. Nucleosome: a major immunogen for pathogenic autoantibody-inducing T cells of lupus. *J Exp Med* 1993;177:1367–81.
 148. Datta SK, Kaliyaperumal A. Nucleosome-driven autoimmune response in lupus. Pathogenic T helper cell epitopes and costimulatory signals. *Ann NY Acad Sci* 1997;815:155–70.
 149. Kaliyaperumal A, Mohan C, Wu W, Datta SK. Nucleosomal peptide epitopes for nephritis-inducing T helper cells of murine lupus. *J Exp Med* 1996;183:2459–69.
 150. Desai-Mehta A, Mao C, Rajagopalan S, Robinson T, Datta SK. Structure and specificity of T cell receptors expressed by potentially pathogenic anti-DNA autoantibody-inducing T cells in human lupus. *J Clin Invest* 1995;95:531–41.
 151. Voll RE, Roth EA, Girkontaite I *et al.* Histone-specific Th0 and Th1 clones derived from systemic lupus erythematosus patients induce double-stranded DNA antibody production. *Arthritis Rheum* 1997;40:2162–71.
 152. Kaliyaperumal A, Michaels MA, Datta SK. Antigen-specific therapy of murine lupus nephritis using nucleosomal peptides: tolerance spreading impairs pathogenic function of autoimmune T and B cells. *J Immunol* 1999;162:5775–83.
 153. Mohan C, Datta SK. Lupus: key pathogenic mechanisms and contributing factors. *Clin Immunol Immunopathol* 1995;77:209–20.
 154. Kalled SL, Cutler AH, Datta SK, Thomas DW. Anti-CD40 ligand antibody treatment of SNF1 mice with established nephritis: preservation of kidney function. *J Immunol* 1998;160:2158–65.
 155. Koshy M, Berger D, Crow MK. Increased expression of CD40 ligand on systemic lupus erythematosus lymphocytes. *J Clin Invest* 1996;98:826–37.
 156. Desai-Mehta A, Lu L, Ramsey-Goldman R, Datta SK. Hyperexpression of CD40 ligand by B and T cells in human lupus and its role in pathogenic autoantibody production. *J Clin Invest* 1996;97:2063–73.
 157. Illei GG, Klippel JH. Novel approaches in the treatment of lupus nephritis. *Lupus* 1998;7:644–8.
 158. Ma J, Xu J, Madaio MP *et al.* Autoimmune lpr/lpr mice deficient in CD40 ligand: spontaneous Ig class switching with dichotomy of autoantibody responses. *J Immunol* 1996;157:417–26.
 159. Finck BK, Linsley PS, Wofsy D. Treatment of murine lupus with CTLA4Ig. *Science* 1994;265:1225–7.
 160. Strand V. Approaches to the management of systemic lupus erythematosus. *Curr Opin Rheumatol* 1997;9:410–20.
 161. Rose LM, Latchman DS, Isenberg DA. Apoptosis in peripheral lymphocytes in systemic lupus erythematosus: a review. *Br J Rheumatol* 1997;36:158–63.
 162. Cheng J, Zhou T, Liu C *et al.* Protection from Fas-mediated apoptosis by a soluble form of the Fas molecule. *Science* 1994;263:1759–62.
 163. Tokano Y, Miyake S, Kayagaki N *et al.* Soluble Fas molecule in the serum of patients with systemic lupus erythematosus. *J Clin Immunol* 1996;16:261–5.
 164. Elkon KB, Marshak-Rothstein A. B cells in systemic autoimmune disease: recent insights from Fas-deficient mice and men. *Curr Opin Immunol* 1996;8:852–9.
 165. Goel N, Ulrich DT, St. Clair EW, Fleming JA, Lynch DH, Seldin MF. Lack of correlation between serum soluble Fas/APO-1 levels and autoimmune disease. [See comments]. *Arthritis Rheum* 1995;38:1738–43.
 166. Onel KB, Tucek-Szabo CL, Ashany D, Lacy E, Nikolic-Zugic J, Elkon KB. Expression and function of the murine CD95/FasR/APO-1 receptor in relation to B cell ontogeny. *Eur J Immunol* 1995;25:2940–7.
 167. Ohsako S, Hara M, Harigai M, Fukasawa C, Kashiwazaki S. Expression and function of Fas antigen and bcl-2 in human systemic lupus erythematosus lymphocytes. *Clin Immunol Immunopathol* 1994;73:109–14.
 168. Emlen W, Niebur J, Kadera R. Accelerated *in vitro* apoptosis of lymphocytes from patients with systemic lupus erythematosus. *J Immunol* 1994;152:3685–92.
 169. Van Houten N, Budd RC. Accelerated programmed cell death of MRL-lpr/lpr T lymphocytes. *J Immunol* 1992;149:2513–7.