OSCE Exam Case 6

Door note:

A doctor is waiting inside for your recommendation, **document your response.**

Ref: Asthma in infants and children

EXAM CASE 26

- Name: Peter Jones Age: 9 Years
- Medical condition: Asthma
- Allergies: No Known Allergies

- Medications:
- Ventolin MDI as needed– since 6 months
- Advair MDI twice daily – since 6 months

New Rx: Montelukast 10 mg PO at bedtime daily

Outcome

- Confirmed any updates on allergies,
- Any updates on current medications, any non-prescription medications.
- Child overall health!
- Correct the dose of Montelukast 5 mg chewable tablet at bedtime.

  Montelukast: 6–14 y: 5 mg (chewable tablet) QHS PO

- Ask the patient to report any symptoms. Boxed warning issued regarding neuropsychiatric effects (e.g., depression, agitation/aggression, hallucinations, suicidal ideation); monitor for symptoms.

Pharmacy PREP OSCE Check List. Case. Constipation

PharmacyPREP.COM

Communication skills

Patient Centred. Opening Statement (respond to patient clue, good listener, did not interrupt patient, engaging). Non-verbal Skills (eye contact, gesture, posture), Identified patient? Understands the sequence of problem

Information gathering process. (Asked M-A-M-A) allergies, medicine history, medical conditions. Builds trust and confidence with patient, Specific lifestyle such as alcohol and smoking habits prn; and/or other considerations relevant to the purpose of the case.

Discuss Follow up/Monitoring relevant to case

Time to check references, and use of proper references and Overall time management

Overall Communications	Acceptable 4/4	Marginally Acceptable 3/4	Marginally unacceptable 2/4	Unacceptable 1-0/4

Problem Solving

key elements of the counselling task

Check for red flags, impaction/ bowel obstruction

Patient education: "normal" BM frequency varies, non pharms, develop bowel routine, monitoring

Nonpharms: create routine for toileting e.g. attempt 5-15 min before every meal until 1 BM; dietary fibre (age plus 5g/day; or 10 g/day); sorbitol/barley malt/corn syrup (stool softener); prune; diet (fruits, vegetables, whole grains); exercise and hydration

Pharms:

Best: lactulose 5 ml bid [or sorbitol 1-3ml/kg/d in divided doses] (osmotic agents) → onset 24-48h for both

Others (not best choice):

Second line: senna, stool softeners (e.g. docusate)

see MD if >2 weeks chronic constipation or no BM > 7 days despite laxatives;

Overall Problem Solving	Fully solved 4/4	Marginally solved 3/4	Marginally unsolved 2/4	Unsolved 1-0/4
Note.	Misinformation?		Put patients in to risk?	O