Exam Case # 7

CANDIDATE'S INSTRUCTIONS
A women approaches you for a concern. Kindly counsel her as you would in practice. Also, assist her with any concerns she has regarding her child condition

.

Time Frame: You have 7 minutes to complete this station

References: PSC and other references

PharmacyPREP.COM


Dentinox®
INFANT
Colic Drops
Oral Suspension
Activated Dimeticone
Effective pain relief
from colic from
birth onwards
SYRINGE FOR
EASY APPLICATION
Dentinox

PSP
Baby's Choice
GRIPE WATER
EAU CONTRE LA COLIQUE
250 mL
life
Gripe
Water
For Quick Relief of Baby's
Minor Stomach Upsets
250 mL
equate
Alcohol Free
Gripe Water
250 mL

Standardized patient

SP’s opening statement, my child is crying lot and cranky. I donʼt know what to do?
I am feeling terrible?.
If candidate says did you talk to your doctor? Say no
SPs says My daughter started crying a lot from last night?
If a candidate asks?
Does your child have a fever? No
Does your child cry continuously for more than 3 h? No she cries frequently for five to 10 minutes and is cranky and screaming

Child information.
Child age: 4 mo, Child weight 15 lbs, NOT giving any medication, otherwise healthy. No vomiting.
The last immunization was 3 mo ago.
Weight growth seems normal

Problem solved

- Infant colic:
- - NICE guidelines (US routine postnatal care for women and their babies): http://www.nice.org.uk/nicemedia/live/10988/30144/30144.pdf pages 32-33 has a short section on colic
- - check that s/s are consistent with colic: "duration of crying 3h per day on several days of the week..." Patient Self Care p322 Figure 1
- - check for red flags: "fever, lack of weight gain, excessive vomiting, changes in stooling or urination, dehydration, crying 3h nonstop." from Patient Self Care p322 Figure 1
- - key points: **reassurance** and **empathy for parents**, colic is selflimiting (3-4 months), offer and focus on non-pharms (PSC Ch 36), pharmacological treatment is not required.
- - specifically don't recommend dicyclomine (can cause breathing problems and coma) and gripe water --> reference: NICE guidelines and Patient Self Care Ch 36 p325

Pharmacy PREP OSCE Check List. Case Infant Colic

Communication skills

Patient Centred. Opening Statement (respond to patient clue, good listener, did not interrupt patient, engaging). Non-verbal Skills (eye contact, gesture, posture), Identified patient? Understands the sequence of problem . Empathy.

Information gathering process. (Asked M-A-M-A) allergies, medicine history, medical conditions. Builds trust and confidence with patient, Specific lifestyle such as alcohol and smoking habits prn; and/or other considerations relevant to the purpose of the case.

Discuss Follow up/Monitoring relevant to case

Time to check references, and use of proper references and Overall time management

Overall Communications	Acceptable 4/4	Marginally Acceptable 3/4	Marginally unacceptable 2/4	Unacceptable 1 to 0/4

Problem Solving

key elements of the counseling task. NICE guidelines
•check for red flags: "fever, lack of weight gain, excessive vomiting, changes in stooling or urination, dehydration, crying 3h nonstop." from CTMA Chapter Infant Colic Figure 1
•- key points: reassurance and empathy for parents, colic is self limiting (3-4 months), offer and focus on non-pharms (CTMA), pharmacological treatment is not required.
•Specifically don't recommend dicyclomine (can cause breathing problems and coma) and gripe water
y elements of the counseling task

Overall Problem Solving	Fully solved 4/4	Marginally solved 3/4	Marginally unsolved 2/4	Unsolved 1-0/4
Note.	Misinformation?		Put patients in to risk?	O