Pharmacy prep OSCE case 9

Door Note

A woman coming for OTC recommendation. Respond as you would in regular practice.

Time: you have 7 minutes to complete the station.

Reference: Minor ailment N&V

EXAM CASE 46

SP instructions:

- A woman complains that she is sick to the stomach and throwing up. Want to buy some medication for her condition.
- On average in a day for how long do you feel nauseated or sick to the stomach? 2 to 3 hour
- On average in a day how much do you vomit or through up. 1-2
- On average in a day how many times do you retouch or dry heaves without bringing anything up? 1-2

Patient profile

Name: Sara Azam Age: 28 yr ,
Medical condition: Pregnant (1st trimester)
Allergies: No known allergies
Medications: Pregvit → started taking 15 days ago

On the Table: Ginger tablets, Gravol, Diphenhydramine, Imodium, Tylenol

Reference: CTMA

Solution

Rule out red flags: fever, dehydration, blood in vomit.

Assess the severity of N&V in pregnancy.

On average in a day for how long do you feel nauseated or sick to the stomach? 2 to 3 hour

On average in a day how much do you vomit or through up. 1-2

On average in a day how many times do you retouch or dry heaves without bringing anything up? 1-2

Based on her presentation. A pharmacist recommends : Ginger tablets 250 mg q6h

Counsel about the sedation side effects, and avoid driving after taking medications for few hours.

If your conditions is worsening and more frequent and prolonged period of nausea and vomiting, consult the doctor.

Nonpharms: Eat before getting out of bed in the morning to help prevent early morning nausea

Eat small, frequent meals, eat bland food, and avoid spicy and fatty food. Take frequent naps.

Follow-up: will call in 2 days to see if any improvement.


When pregnancy confirmed, assess presence and severity of nausea and vomiting
Mild NVP
Moderate-to-severe NVP or hyperemesis gravidarum
Nonpharmacologic therapies:
• Eat small, frequent meals
• Eat bland foods
• Avoid spicy, fatty or strong-smelling foods
• Take frequent naps/rests
• Shorten work day, if possible
• Practice P6 acupressure
Pharmacologic therapies:
• Ginger tablet 250 mg Q6H PO
• Pyridoxine 25 mg Q8H PO
Start/continue any or all interventions as for mild NVP and add a first-line agent:
• Dimenhydrinate 50 mg Q4–6H PRN PO/PR/IV
• Diphenhydramine 25 mg Q4–6H or 50 mg Q6–8H PRN PO
• Promethazine 12.5–25 mg Q4–6H PRN PO
Persistent NVP
Persistent NVP or hyperemesis without dehydration
Persistent NVP or hyperemesis with dehydration