



## Cimduo

lamivudine/tenofovir DF (3TC/TDF)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN COMBINED WITH TIVICAY, ISENTRESS HD, OR ISENTRESS

One tablet (300 mg lamivudine/ 300 mg tenofovir DF), once daily.  
Take with or without food.



## Descovy

emtricitabine/tenofovir alafenamide (FTC/TAF)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

One tablet (200 mg emtricitabine/ 25 mg tenofovir alafenamide), once daily.  
Take with or without food.



## Emtriva

emtricitabine (FTC)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

One 200 mg capsule, once daily.  
Take with or without food.



## Epivir

lamivudine (3TC)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND ABACAVIR

One 300 mg tablet, once daily;  
or one 150 mg tablet, twice daily.  
Take with or without food.  
**GENERIC IS AVAILABLE.**



## Epzicom

abacavir/lamivudine (ABC/3TC)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE IF HLA-B\*5701-NEGATIVE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR (AS TRIUMEQ)

One tablet (600 mg abacavir/ 300 mg lamivudine), once daily.  
Take with or without food.  
**GENERIC IS AVAILABLE.**



## Truvada

emtricitabine/tenofovir DF (FTC/TDF)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

One tablet (200 mg emtricitabine/ 300 mg tenofovir DF), once daily.  
Take with or without food.  
**APPROVED AS GENERIC, WHICH IS NOT YET COMMERCIALY AVAILABLE.**



## Viread

tenofovir disoproxil fumarate (TDF)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

One 300 mg tablet, once daily.  
Take with or without food.  
**GENERIC IS AVAILABLE.**



## Ziagen

abacavir sulfate (ABC)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE IF HLA-B\*5701-NEGATIVE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND LAMIVUDINE (AS TRIUMEQ)

Two 300 mg tablets, once daily;  
or one 300 mg tablet, twice daily.  
Take with or without food.  
**GENERIC IS AVAILABLE.**



## Temixys

lamivudine/tenofovir DF (3TC/TDF)

★ RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN COMBINED WITH TIVICAY, ISENTRESS HD, OR ISENTRESS

One tablet (300 mg lamivudine/ 300 mg tenofovir DF), once daily.  
Take with or without food.