

Envelope Equations

USPAS - Beam Physics with Intense Space Charge

Austin Hoover Steven Lund Daniel Winklehner John Barnard

July 2026

Paraxial ray equation

Envelope equations for axially symmetric beams

Envelope equations for elliptically symmetric beams

Envelope equations: reduced-order characterization of beam dynamics in terms of second-order moments

[Image]

Road map

- Start from single particle equation of motion: $\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$.
- Make use of:
 - Paraxial (near-axis) approximation
 - Conservation of angular momentum
 - Axisymmetry of external forces
- Arrive at single-particle paraxial ray equation.
- Take statistical averages to obtain moment/envelope equations.

Paraxial ray equation

Equation of motion in cylindrical coordinates

Equation of motion for particle of mass m and charge q in electric field $\mathbf{E}$ and magnetic field $\mathbf{B}$:

$$\frac{d\mathbf{p}}{dt} = \frac{d(\gamma m \dot{\mathbf{x}})}{dt} = q(\mathbf{E} + \dot{\mathbf{x}} \times \mathbf{B}), \quad (1)$$

Switch to cylindrical coordinates (r, θ, z) :

$$\begin{aligned} \mathbf{p} &= p_x \hat{x} + p_y \hat{y} + p_z \hat{z} \\ &= p_r \hat{r} + p_\theta^* \hat{\theta} + p_z \hat{z}. \end{aligned} \quad (2)$$

The components p_r and p_θ^* are given by

$$\begin{aligned} p_r &= \gamma m \dot{r}, \\ p_\theta^* &= \gamma m r \dot{\theta}. \end{aligned} \quad (3)$$

The symbol p_θ^* is used to reserve p_θ for the *canonical* angular momentum, defined later.

In cylindrical coordinates, the unit vectors depend on the particle position and thus change with time:

$$\begin{aligned}\dot{\hat{r}} &\equiv d\hat{r}/dt = \dot{\theta}\hat{\theta}, \\ \dot{\hat{\theta}} &\equiv d\hat{\theta}/dt = -\dot{\theta}\hat{r}.\end{aligned}\tag{4}$$

Thus, the time derivative of the momentum vector in Eq. (2) is:

$$\begin{aligned}\frac{d\mathbf{p}}{dt} &= \frac{d}{dt} (p_r\hat{r} + p_\theta^*\hat{\theta} + p_z\hat{z}) \\ &= \dot{p}_r\hat{r} + p_r\dot{\hat{r}} + \dot{p}_\theta\hat{\theta} + p_\theta^*\dot{\hat{\theta}} + \dot{p}_z\hat{z} \\ &= (\dot{p}_r - p_\theta^*\dot{\theta})\hat{r} + (\dot{p}_\theta + p_r\dot{\theta})\hat{\theta} + \dot{p}_z\hat{z} \\ &= \left[\frac{d(\gamma m \dot{r})}{dt} - \gamma m r \dot{\theta}^2 \right] \hat{r} + \left[\frac{d(\gamma m r \dot{\theta})}{dt} + \gamma m \dot{r} \dot{\theta} \right] \hat{\theta} + \left[\frac{d(\gamma m \dot{z})}{dt} \right] \hat{z}. \\ &= \left[\frac{d(\gamma m \dot{r})}{dt} - \gamma m r \dot{\theta}^2 \right] \hat{r} + \left[\frac{1}{r} \frac{d}{dt} (\gamma m r^2 \dot{\theta}) \right] \hat{\theta} + \left[\frac{d(\gamma m \dot{z})}{dt} \right] \hat{z}.\end{aligned}\tag{5}$$

This takes care of the left side of the Lorentz force equation (Eq. (1)).

The right side of Eq. (1) requires the cylindrical field components:

$$\begin{aligned}\mathbf{E} &= E_r \hat{r} + E_\theta \hat{\theta} + E_z \hat{z}, \\ \mathbf{B} &= B_r \hat{r} + B_\theta \hat{\theta} + B_z \hat{z}.\end{aligned}\tag{6}$$

Evaluating the cross product $\dot{\mathbf{x}} \times \mathbf{B}$,

$$\dot{\mathbf{x}} \times \mathbf{B} = \begin{vmatrix} \hat{r} & \hat{\theta} & \hat{z} \\ \dot{r} & r\dot{\theta} & \dot{z} \\ B_r & B_\theta & B_z \end{vmatrix},\tag{7}$$

leads to

$$\begin{aligned}\frac{d(\gamma m \dot{r})}{dt} - \gamma m r \dot{\theta}^2 &= q (E_r + r \dot{\theta} B_z - \dot{z} B_\theta), \\ \frac{1}{r} \frac{d(\gamma m r^2 \dot{\theta})}{dt} &= q (E_\theta + \dot{z} B_r - \dot{r} B_z), \\ \frac{d(\gamma m \dot{z})}{dt} &= q (E_z + \dot{r} B_\theta - r \dot{\theta} B_r)\end{aligned}\tag{8}$$

Conservation of canonical angular momentum

Write the fields in terms of the scalar potential $\phi(\mathbf{x})$ and vector potential $\mathbf{A}(\mathbf{x})$:

$$\begin{aligned}\mathbf{E} &= -\nabla\phi - \frac{\partial\mathbf{A}}{\partial t}, \\ \mathbf{B} &= \nabla \times \mathbf{A}.\end{aligned}\tag{9}$$

The gradient $\nabla\phi$ in cylindrical coordinates is:

$$\nabla\phi = \left[\frac{\partial\phi}{\partial r}\right]\hat{r} + \left[\frac{1}{r}\frac{\partial\phi}{\partial\theta}\right]\hat{\theta} + \left[\frac{\partial\phi}{\partial z}\right]\hat{z}.\tag{10}$$

And the curl $\nabla \times \mathbf{A}$ is:

$$\nabla \times \mathbf{A} = \left[\frac{1}{r}\frac{\partial A_z}{\partial A_\theta} - \frac{\partial A_\theta}{\partial z}\right]\hat{r} + \left[\frac{\partial A_r}{\partial z} - \frac{\partial A_z}{\partial r}\right]\hat{\theta} + \left[\frac{1}{r}\left(\frac{\partial(rA_\theta)}{\partial r} - \frac{\partial A_r}{\partial\theta}\right)\right]\hat{z}.\tag{11}$$

Axial symmetry means all partial derivatives with respect to θ are zero in Eq. (10) and Eq. (??). This gives the following expressions for the fields in terms of the potentials in cylindrical coordinates:

Series expansion of axially symmetric fields

Consider external fields $\mathbf{E}$ and $\mathbf{B}$ with azimuthal symmetry ($\partial/\partial\theta = 0$). In the absence of charge and current densities,

$$\begin{aligned}\nabla \cdot \mathbf{E} &= 0, \\ \nabla \cdot \mathbf{B} &= 0, \\ \nabla \times \mathbf{E} &= 0, \\ \nabla \times \mathbf{B} &= 0.\end{aligned}\tag{19}$$

The vanishing curl means we can express the fields as

$$\begin{aligned}\mathbf{E} &= -\nabla\phi, \\ \mathbf{B} &= -\nabla\phi_m,\end{aligned}\tag{20}$$

for scalar potentials ϕ and ϕ_m . Additionally, the vanishing divergence leads to the Laplace equation for the scalar potential:

$$\nabla^2\phi = 0.\tag{21}$$

In cylindrical coordinates, the Laplace equation reads

Derivation of paraxial ray equation

We will now take the *paraxial* limit of small transverse velocity relative to longitudinal velocity: $\dot{r} \ll \dot{z}$, $r\dot{\theta} \ll \dot{z}$. If $v = (v_r^2 + v_\theta^2 + v_z^2)^{1/2}$ is the total velocity, where $v_r = \dot{r}$, $v_\theta = r\dot{\theta}$, and $v_z = \dot{z}$, the paraxial limit gives $\dot{z} = (v^2 - \dot{r}^2 - r^2\dot{\theta}^2)^{1/2} \approx v$. We will therefore only keep terms linear in r in the equations of motion.

We will also add the self-generated fields to the equation of motion, writing

$$\begin{aligned}\mathbf{E} &= \mathbf{E}^{(ext)} + \mathbf{E}^{(self)}, \\ \mathbf{B} &= \mathbf{B}^{(ext)} + \mathbf{B}^{(self)}.\end{aligned}\tag{35}$$

[Is there a way to derive the ray equation first, then add the self-fields later?]
To first order in r , the external fields are:

$$\begin{aligned}E_r^{(ext)} &\approx \frac{r}{2} \frac{\partial^2 V(z)}{\partial z^2}, \\ E_z^{(ext)} &\approx -\frac{\partial V(z)}{\partial z}, \\ B_r^{(ext)} &\approx -\frac{r}{2} \frac{\partial B(z)}{\partial z},\end{aligned}\tag{36}$$

Envelope equations for axially symmetric beams

Envelope equations for axially symmetric beams

The strategy in this section will be to compute the second moments of the transverse distribution function $f(x, x', y, y', s)$. For axially symmetric beams, the Vlasov equation will generate a set of differential equations for the evolution of the second moments such as $\langle rr \rangle$, $\langle rr' \rangle$, $\langle rr'' \rangle$, etc. Combining with the paraxial ray equation will generate a set of ordinary differential equations (ODEs) for the evolution of the beam *envelope* radius $r_b = \sqrt{2\langle rr \rangle}$.

Start with the Vlasov equation:

$$\frac{\partial f}{\partial s} + \mathbf{x}' \cdot \frac{\partial f}{\partial \mathbf{x}} + \mathbf{x}'' \cdot \frac{\partial f}{\partial \mathbf{x}'} = 0, \quad (53)$$

where $\mathbf{x} = (x, y)$. Multiply the Vlasov equation by $g(\mathbf{x}, \mathbf{x}')$, an arbitrary function of the phase space coordinates, and integrate over all phase space:

$$\iint \left[g \frac{\partial f}{\partial s} + g \mathbf{x}' \cdot \frac{\partial f}{\partial \mathbf{x}} + g \mathbf{x}'' \cdot \frac{\partial f}{\partial \mathbf{x}'} \right] d\mathbf{x} d\mathbf{x}' = 0. \quad (54)$$

Introduce bracket notation to indicate the expected value under the distribution; e.g.,

Envelope equations for elliptically symmetric beams
