

CHANGE OF VARIABLES IN INTEGRATION

AUSTIN DAVID BROWN

1. CHANGE OF VARIABLES IN INTEGRATION

The main idea behind change of variables is changing the integration from the domain to the codomain of some function.

COV 1

See Linear Algebra Done Right for some insight.

Let $f : \mathbb{R}^n \rightarrow [0, +\infty]$ be measurable. Let $T \in GL_n(\mathbb{R})$. Since T is linear, $vol(T(\mathbb{R}^n)) = |\det(T)|vol(\mathbb{R}^n)$. Hence,

$$\int_{T(\mathbb{R}^n)} f d\mu = |\det(T)| \int_{\mathbb{R}^n} f \circ T d\mu.$$

COV 1 for differentiation

Let $f : \mathbb{R}^n \rightarrow [0, +\infty]$ be measurable.

Let $\varphi : U \rightarrow \mathbb{R}^n$ be a diffeomorphism.

Since the differential $\varphi(x) + D\varphi(x) : U \rightarrow \mathbb{R}^n$ is a linear approximation to φ near x , shifting by $\varphi(x)$ does not change volume, and since $D\varphi(x)$ is linear,

$$\int_{\varphi(U)} f(y) dy = \int_U f \circ \varphi(x) |\det(D\varphi(x))| dx.$$

COV 2

Lebesgue integral version:

This is the pushforward change of variable. The idea is that the way that this pushforward measure is defined, there is no change in volume.

Let $f : X \rightarrow [0, +\infty]$ be a measurable map. Let $g : X \rightarrow Y$ be a measurable map.

If μ is a measure on X , then μg^{-1} is a pushforward measure or image measure on the sigma algebra of Y . Then

$$\int_{g(X)} f(y) d\mu(g^{-1}(y)) = \int_X f(g(x)) d\mu(g g^{-1}(x)) = \int_X f(g(x)) d\mu(x).$$

Example

The main change of variables for random variables is this. If $f = id$, and $X : \Omega \rightarrow \mathbb{R}$ is a *RV*, then

$$E(X) = \int_{\Omega} X(\omega) dP(\omega) = \int_{X(\Omega)} x dPX^{-1}(x).$$

By definition,

$$\int_{X(\Omega)} x dPX^{-1}(x) = \int_{X(\Omega)} x dP_X(x) = \int_{X(\Omega)} x dF_X(x).$$

COV 3

Riemann-Stieltjes integral version:

Let $f : [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Let $g : [a, b] \rightarrow \mathbb{R}$ be continuous and g' is Riemann integrable. Then

$$\int_{[a,b]} f(x) d(g(x)) = \int_{[a,b]} f(x) g'(x) dx$$

Lebesgue integral version:

Let g be absolutely continuous and it will hold.

Examples:

This one is useful when $g = F_X$ is the distribution function of a random variable.