INTRODUCTION TO MEDICAL STATISTICS: FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

Mirjana Kujundžić Tiljak

FIRST STEP

→ summarize data to get a "feel" for the data

DIAGRAMS

TABLES

SUMMARY STATISTICS

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

ONE VARIABLE

EMPIRICAL FREQUENCY DISTIRBUTION of

variable relates each possible observation, class of observations (i.e. range of values) or category, as appropriate, to its observed frequency of occurrence

replacement of each frequency by a relative frequency (the percentage of the total frequency) → to compare frequency distributions in two or more groups of individuals

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

3


CATEGORICAL OR DISCRETE NUMERICAL DATA


BAR or COLUMN CHART

- separate horizontal or vertical bar is drawn for each category,
- bar's length being proportional to the frequency in that category
- bars are separated by small gaps to indicate that the data are categorical or discrete

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION






PIE CHART

- circular "pie" is split into sections
- · one section for each category
- area of each section is proportional to the frequency in that category

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION



CONTINUOUS NUMERICAL DATA

more difficult to display

 \rightarrow the data may need to bi summarized before being drawn

HISTOGRAM

- similar to a bar chart no gaps between the bars as the data are continuous
- width of each bar of the histogram relates to a range of values for the variable

26.06.2006


FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

9

- the area of the bar is proportional to the frequency in that range
- if one of the groups covers a wider range than the others – base will be wider and height shorter to compensate
- usually between five and 20 groups are chosen
- ranges should be so narrow to illustrate patterns in the data, but not so narrow that they are the row data
- histogram should be labeled carefully to make it clear where the boundaries lie

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION




DOT PLOT

- each observation is represented by on dot on a horizontal or vertical line
- · very simple to draw
- summary measure of the dana (i.e. mean or median) is shown on the diagram
- · may be used for discrate data

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION



STEM-AND-LEAF PLOT

- · mixture of a diagram and a table
- looks similar to histograms turned on its side
- Is effectively the data values written in increasing order of size
- vertical stem first few digits of the values, arranged in order
- Protruding from the stem = leaves (i.e. final digit of each of the ordered values, which are written horizontally in increasing numerical order

26.06.2006 FREQUENCY DISTRIBUTIONS AND

GRAPHIC PRESENTATION

STEM AND LEAF PLOT

database "STUDENT", variable "HEIGHT", female examinees

| Stem and Leaf Plot: HEIGHT, FEMALES one leaf=1 case | | |
|---|---------|-------------|
| | Class n | Percentiles |
| 15 3 | 2 | |
| 15 67889 | 8 | |
| 16 0000001222222333344444 | 44 | 25% |
| 16 555556667777777788888888889999 | 58 | median |
| 17 00000000001111122222222333334444 | 66 | 75% |
| 17 55555667889 | 21 | |
| 18 00134 | 7 | |
| 18 | 0 | |
| min = 153,0000 max = 184,0000 Total N: | 206 | |

26.06.2006 FREQUENCY DISTRIBUTIONS AND
GRAPHIC PRESENTATION

15

BOX PLOT or BOX-AND-WHISKER PLOT

- · vertical or horizontal rectangle
- the ends of the rectangle = upper and lower quartiles of the data values; standard deviation
- line drawn through the rectangle = median value; mean value
- whiskers, starting at the ends of the rectangle = minimum and maximum values; particular percentiles (e.g. 5th and 95th percentiles
- · outliers may be marked

26.06.2006 FREQUENCY DISTRIBUTIONS AND 16 GRAPHIC PRESENTATION

FREQUENCY DISTRIBUTION – "SHAPE"

- The choice of the most appropriate statistical method will often depend on the shape of distribution
- unimodal single "peak"
- bimodal two peaks
- uniform each value is equally likely, no peaks
- Unimodal distribution main aim = to see where the majority of the data values lie, relative to the maximum and minimum values

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

17

FREQUENCY DISTRIBUTION – "SHAPE"

- symmetrical centered around some mid-poing, with one side being a mirror-image of the other
- skewed to the right (positively skewed) long tail to the right with one of a few high values
- skewed to the left (negatively skewed)

 long tail to the left with one or a few low values

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

TWO VARIABLES

- one variable is categorical → separate diagrams showing the distribution of the second variable can be drawn for each of the categories
- · clustered or segmented bar or column charts

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

19

TWO VARIABLES

- both continuous or ordinal variable the relationship between the two
 - → SCATTER DIAGRAM
- plots one variable against the other in a two way diagram
- · X variable on horizontal axis
- · Y variable on vertical axis

26.06.2006

FREQUENCY DISTRIBUTIONS AND GRAPHIC PRESENTATION

