

Resolving Pisanski's Conjecture on Quadrilateral Embeddings

Alexander Metzger¹ and Austin Ulrigg²

¹University of California, Berkeley, alexmetzger@berkeley.edu

²University of Washington, austinul@uw.edu

Abstract

We prove Pisanski's conjecture on quadrilateral embeddings of regular bipartite graphs. For integers $m \geq d \geq 4$ with $4 \mid md$, we construct both a connected simple d -regular bipartite graph G with m vertices in each bipartition class and an orientable quadrangulation. This graph's genus is

$$\gamma(G) = 1 + \frac{m(d-4)}{4}.$$

These quadrangulations attain the lower bound on the genus from Euler's formula. We give dihedral constructions for even m and a cyclic construction for every $m \geq d$ with $4 \mid d$. In each case, two cyclic permutations of a connection set of size d determine all of the local rotations.

1 Introduction

In 1980, Tomaž Pisanski conjectured the following for positive integers m and d . Whenever $m \geq d \geq 4$ and $4 \mid md$, there exists a connected d -regular bipartite graph, with m vertices in each class, that has an orientable quadrangulation [8, Conjecture 1]. We prove this conjecture by constructing an orientable quadrangulation for every such pair (m, d) .

For a graph G , let $\gamma(G)$ denote its orientable genus. If a d -regular bipartite graph with m vertices in each class has an orientable quadrangulation, the surface has genus $1 + m(d-4)/4$, attaining the lower bound on the graph's genus from Euler's formula [8, p. 33]. Having d distinct neighbors in the opposite class also requires $m \geq d$. The surface's genus must be an integer, so $4 \mid md$ is necessary. Pisanski's conjecture asks whether these necessary conditions are sufficient when $d \geq 4$.

Many of the graph families whose orientable genus is known are highly symmetric. In 1965, Ringel determined the genus of the complete bipartite graphs [9]. In particular, $K_{m,m}$ has an orientable quadrangulation for every even m . Mohar, Parsons and Pisanski determined the orientable genus of nearly complete bipartite graphs [6]. For a perfect matching M in $K_{m,m}$, they proved

$$\gamma(K_{m,m} - M) = \left\lceil \frac{(m-1)(m-4)}{4} \right\rceil \quad (m \geq 3).$$

When $m \equiv 0, 1 \pmod{4}$, the expression inside the ceiling is an integer and the bound is attained by a quadrangulation. For $m \geq 5$, these give cases of Pisanski's conjecture with $d = m - 1$. Alpert and Gross [1] and Anderson [2] proved further partial cases. Anderson proved existence when m and d are both even, and when m is odd with $d \equiv 4 \pmod{8}$ and $d \geq 12$. More recently, Ulrigg and Metzger [11] determined the genus of an infinite family of bipartite Kneser graphs, giving explicit quadrangulations that realize the bound in Pisanski's conjecture.

The constructions of this paper exploit group-invariant rotation systems, where we choose the group and connection set together with the local rotations. These freedoms allow us to ensure that every facial walk closes after exactly 4 steps while ensuring that both permutations are cyclic.

Theorem 1.1 (Main theorem). *Let m and d be integers with $m \geq d \geq 4$. There exists a connected simple d -regular bipartite graph with m vertices in each bipartition class and a cellular orientable embedding whose face boundaries are simple 4-cycles if and only if $4 \mid md$. Whenever $4 \mid md$, the graph G constructed below has minimum orientable genus*

$$\gamma(G) = 1 + \frac{m(d-4)}{4}.$$

We construct Haar graphs [4] of groups of order m , using two cyclic orders on a connection set of size d to specify all local rotations. For even m and d , this recovers the graph family quadrangulated by Anderson [2, pp. 357–358]. The dihedral construction also covers every admissible pair with odd d . Our cyclic construction applies whenever $4 \mid d$, extending the degree range for the cyclic graph family in Anderson’s Corollary 2.2 to include $8 \mid d$. Together these constructions cover every pair (m, d) satisfying the conditions of Pisanski’s conjecture.

Sections 2–3 give the preliminary definitions and background material, and Sections 4–6 give the constructions and proof.

2 Preliminaries

We follow the definitions of *Graphs on Surfaces* by Mohar and Thomassen [7]. Throughout this paper, graphs are finite, simple, and undirected, and surfaces are connected and closed.

An embedding of a graph on a surface represents its vertices by distinct points and its edges by simple arcs joining their endpoints. These arcs meet only at common endpoints and do not pass through any other vertices. The connected components of the complement of the embedded graph are its *faces*. An embedding is *cellular*, or a *2-cell embedding*, if every face is homeomorphic to an open disk. The *orientable genus* $\gamma(G)$ of a graph G is the smallest nonnegative integer g for which G embeds in the orientable surface S_g , the connected sum of g tori, where S_0 is the sphere. It is known that every minimum-genus embedding of a connected graph in an orientable surface is cellular [7, 13].

An *orientable quadrangulation* is a cellular embedding in an orientable surface in which every facial walk has length four. If G is connected, simple, bipartite, and d -regular, with m vertices in each bipartition class and $d \geq 4$, Euler’s formula gives the lower bound on the orientable genus of G [8, p. 33].

$$\gamma(G) \geq \left\lceil 1 + \frac{m(d-4)}{4} \right\rceil. \quad (1)$$

If all faces of an orientable embedding of G are quadrilaterals, as in an orientable quadrangulation, the surface on which G embeds has genus $1 + m(d-4)/4$, attaining this bound and having $4 \mid m \times d$.

Cellular embeddings can be described combinatorially by listing the local rotations at each vertex. Given a cellular embedding of G in an orientable surface S , fix an orientation of S . Around each vertex v , we can list its incident edges in clockwise order. This defines a cyclic permutation π_v , where $\pi_v(e)$ is the edge that follows e in that order. Each permutation π_v is called the *local rotation* at v , and the set $\Pi = \{\pi_v \mid v \in V(G)\}$ is the *rotation system* of the embedding. More generally, a choice of one cyclic permutation of the incident edges at each vertex is called a rotation system of G . The Heffter–Edmonds–Ringel rotation principle gives a one-to-one correspondence between rotation systems and cellular orientable embeddings [7, pp. 87–90].

Theorem 2.1 (Heffter–Edmonds–Ringel). *Every rotation system of a connected graph with at least one edge determines a cellular embedding in an orientable surface. This embedding is unique up to a homeomorphism that induces the identity on the graph.*

We now describe the facial walks determined by a given rotation system. A *dart* of G is a pair (v, e) , where e is an edge incident with v . If e joins v and w , its two darts are (v, e) and (w, e) . Let R denote vertex rotation and τ denote dart reversal, so

$$R(v, e) = (v, \pi_v(e)), \quad \tau(v, e) = (w, e).$$

We compose permutations from right to left. The *facial permutation* is

$$F = R\tau. \tag{2}$$

Repeatedly applying F returns to the starting dart and traces a closed walk, called a *facial walk*. The orbits of F are the facial walks of the embedding, up to cyclic shift. Therefore, a rotation system of a connected graph whose facial walks all have length four determines an orientable quadrangulation.

3 Haar graphs and Quadrangulations

Representing the vertices of G by elements of a finite group, with one copy for each bipartition class, we choose one cyclic order for the connection set for all A -vertices and another for all B -vertices.

Definition 3.1 (Haar graph). Let Γ be a finite group and $S \subseteq \Gamma$. Define $H(\Gamma, S)$ with vertices

$$\{A_g : g \in \Gamma\} \cup \{B_g : g \in \Gamma\}$$

and edges $\{A_g B_{gs} : g \in \Gamma, s \in S\}$.

The two displayed vertex sets are disjoint, even when their group subscripts agree. In particular, the identity may belong to S . The graph $H(\Gamma, S)$ is simple, bipartite, and $|S|$ -regular, with $|\Gamma|$ vertices in each class. For nonempty S , it is connected if and only if $SS^{-1} = \{st^{-1} : s, t \in S\}$ generates Γ [4, Lemma 1(i)]. In particular, if $1_\Gamma \in S$ and S generates Γ , then $H(\Gamma, S)$ is connected. Let $|S| = d$, and let ρ_A and ρ_B be d -cycles on S . Write $(g, s)_A$ for the dart $A_g \rightarrow B_{gs}$ and $(g, s)_B$ for the dart $B_g \rightarrow A_{gs^{-1}}$. Define the rotations by

$$R(g, s)_A = (g, \rho_A(s))_A, \quad R(g, s)_B = (g, \rho_B(s))_B.$$

At every vertex, the label determines the neighboring vertex uniquely, so these are cyclic orders of all incident darts. Dart reversal is given by

$$\tau(g, s)_A = (gs, s)_B, \quad \tau(g, s)_B = (gs^{-1}, s)_A.$$

Thus, the face permutation satisfies

$$\begin{aligned} F(g, s)_A &= (gs, \rho_B(s))_B, \\ F(g, s)_B &= (gs^{-1}, \rho_A(s))_A, \end{aligned} \tag{3}$$

and hence

$$F^2(g, s)_A = (gs\rho_B(s)^{-1}, \rho_A\rho_B(s))_A. \tag{4}$$

In (4), the label changes by $\rho_A\rho_B$ and the group subscript is multiplied on the right by $s\rho_B(s)^{-1}$. We choose ρ_A and ρ_B so that applying this rule twice returns to the starting dart, but applying it once does not. Every face contains an A -dart, so this gives all facial orbits. In the dihedral constructions, we take $\rho_A = \rho_B^{-1}$ and choose the cyclic order ρ_B so that $s\rho_B(s)^{-1}$ has order two for every $s \in S$. By (4), two facial steps leave the label fixed and change the group subscript. Two further steps then return to the original dart.




Figure 1: A face of Lemma 3.2. Each arrow is labeled by its dart and its group multiplier [11].

Lemma 3.2 (Quadrangulations from involutions). *Let $s_0, \dots, s_{d-1}$ be distinct elements of a finite group Γ , and put $S = \{s_0, \dots, s_{d-1}\}$. Suppose that*

$$q_i = s_i s_{i+1}^{-1}$$

is a nonidentity involution for every i modulo d . For connected $H(\Gamma, S)$, the rotations

$$\rho_B = (s_0 \ s_1 \ \cdots \ s_{d-1}), \quad \rho_A = \rho_B^{-1}$$

give an orientable quadrangulation of $H(\Gamma, S)$. Every face boundary is a simple 4-cycle.

Proof. Both local label permutations are d -cycles. By (4),

$$F^2(g, s_i)_A = (gq_i, s_i)_A, \quad F^4(g, s_i)_A = (gq_i^2, s_i)_A = (g, s_i)_A.$$

The first equality does not return the original dart because $q_i \neq 1_\Gamma$. Since F interchanges the two types of darts, every facial orbit has length exactly four. Since $q_i^2 = 1_\Gamma$ and $q_i = s_i s_{i+1}^{-1}$, we have $q_i s_i = s_{i+1}$. The face starting with $(g, s_i)_A$ has vertices

$$A_g, \quad B_{gs_i}, \quad A_{gq_i}, \quad B_{gs_{i+1}}. \quad (5)$$

The two A -vertices are distinct because $q_i \neq 1_\Gamma$. The two B -vertices are distinct because $s_i \neq s_{i+1}$. Vertices from different classes are distinct by definition. $\square$

4 The dihedral construction

Suppose $m = 2n$ is even. We use the dihedral group of order $2n$, denoted by

$$D_n = \langle r, t \mid r^n = t^2 = 1, \ trt = r^{-1} \rangle.$$

Its elements are $1, r, \dots, r^{n-1}$ and $t, rt, \dots, r^{n-1}t$. We call these the rotations and reflections, respectively. The relation $tr^a = r^{-a}t$ shows that every reflection is an involution. Further, when one of x and y is a rotation and the other is a reflection, the element xy^{-1} is a reflection:

$$r^a(r^b t)^{-1} = r^{a+b}t, \quad (r^a t)(r^b)^{-1} = r^{a+b}t. \quad (6)$$

Thus, a cyclic list of elements of D_n that alternates between rotations and reflections satisfies the order-two condition in Lemma 3.2.

Lemma 4.1 (Alternating rotations and reflections). *Let $d \geq 2$ be even, and let $s_0, \dots, s_{d-1} \in D_n$ alternate between rotations and reflections. For every i modulo d , the element*

$$q_i = s_i s_{i+1}^{-1}$$

satisfies $q_i \neq 1$ and $q_i^2 = 1$.

Proof. Since d is even, every consecutive pair consists of one rotation and one reflection, including the last and first entries. By (6), q_i is therefore a reflection, say $q_i = r^a t$. A reflection is not the identity, and

$$q_i^2 = (r^a t)^2 = r^a (t r^a) t = r^a r^{-a} t^2 = 1.$$

□

4.1 Dihedral rotations for even m and d

Anderson gave an orientable quadrangulation for this graph family [2, pp. 357–358] using White’s composition theorem [12]. We give explicit dihedral rotations for the same graphs.

Proposition 4.2. *Let $m = 2n$, $d = 2k$ with $2 \leq k \leq n$, and $S = \{r^a, r^a t : 0 \leq a < k\}$. Then*

$$\rho_B = (1 \ t \ r \ rt \ r^2 \ r^2 t \ \dots \ r^{k-1} \ r^{k-1} t), \quad \rho_A = \rho_B^{-1} \quad (7)$$

give an orientable quadrangulation of the connected simple d -regular bipartite graph $H(D_n, S)$, with m vertices in each class and simple 4-cycles as face boundaries. For $k = 2$, the cyclic list in (7) stops at rt .

Proof. The entries of ρ_B are distinct and alternate cyclically between rotations and reflections. By Lemma 4.1, every product of a label and the inverse of its successor has order two. Since $1, r, t \in S$, the graph is connected. Lemma 3.2 therefore gives the required quadrangulation. □

When $d = m$, this is $K_{m,m}$ with genus $(m - 2)^2/4$, recovering Ringel’s formula for even m [9]. Figure 2 shows the construction for $m = 6$ and $d = 4$.




Figure 2: The even-degree dihedral group example $H(D_3, \{1, t, r, rt\})$.

4.2 Dihedral rotations for odd d and $4 \mid m$

Some of the odd-degree cases were established by Alpert and Gross [1], and the case $d = m - 1$ follows from the genus of nearly complete bipartite graphs [6]. The following construction applies to every odd $d \geq 5$ and every $m \geq d$ divisible by four.

Suppose $d = 2k + 1$. An odd-length cyclic order cannot alternate between rotations and reflections throughout. We use $k + 1$ rotations and k reflections, arranged so that exactly one consecutive pair consists of two rotations. Since d is odd, the condition $4 \mid md$ requires $4 \mid m$, so $n = m/2$ is even. The rotation $z = r^{n/2}$ has order two, since $z \neq 1$ and $z^2 = r^n = 1$. We take 1 and z as the consecutive rotations, giving $1z^{-1} = z$. Every other consecutive pair will contain one rotation and one reflection.

Proposition 4.3. *Let $m = 2n$ with $4 \mid m$ and $5 \leq d = 2k + 1 \leq m$. Choose distinct rotations $u_0, \dots, u_k$ such that*

$$u_0 = 1, \quad u_1 = z = r^{n/2}, \quad u_2 = r.$$

Put $v_j = r^j t$ for $0 \leq j < k$, and set

$$S = \{u_0, \dots, u_k\} \cup \{v_0, \dots, v_{k-1}\}.$$

Define

$$\rho_B = (u_0 \ u_1 \ v_0 \ u_2 \ v_1 \ \cdots \ u_k \ v_{k-1}), \quad \rho_A = \rho_B^{-1}. \quad (8)$$

These rotations give an orientable quadrangulation of the connected simple d -regular bipartite graph $H(D_n, S)$, with m vertices in each class and simple 4-cycles as face boundaries.

Proof. Since d is odd and $d \leq 2n$, we have $2 \leq k \leq n - 1$. Also $n \geq 4$, because $4 \mid m$ and $m \geq d \geq 5$. Thus $1, z, r$ are distinct, and $k + 1 \leq n$ leaves enough rotations to choose the remaining u_i . The k reflections are distinct as well, so S has exactly d elements. Write $\rho_B = (s_0 \ s_1 \ \cdots \ s_{d-1})$. The first consecutive product is

$$s_0 s_1^{-1} = u_0 u_1^{-1} = z,$$

which has order two. Every other consecutive pair contains one rotation and one reflection, including the last and first entries. Applying Lemma 4.1 to each of these two-element lists shows that $s_i s_{i+1}^{-1}$ has order two for every remaining index i . Thus, for every i modulo d ,

$$s_i s_{i+1}^{-1} \neq 1, \quad (s_i s_{i+1}^{-1})^2 = 1.$$

Since $1, r, t \in S$, the graph is connected. Lemma 3.2 therefore gives the required quadrangulation. $\square$

Figure 3 shows the construction for $m = 8$ and $d = 5$, with $z = r^2$.




Figure 3: The odd-degree dihedral group example $H(D_4, \{1, r^2, t, r, rt\})$.

5 Cyclic rotations for degrees divisible by four

The dihedral construction covers every admissible pair (m, d) with even m . For odd m , the condition $4 \mid md$ requires $4 \mid d$. We give cyclic rotations that work for every $m \geq d \geq 4$ with $4 \mid d$. For $m > d$, Anderson quadrangulated the same cyclic graph family when $d \equiv 4 \pmod{8}$ and $d \geq 12$ [2, Corollary 2.2]. His graph is isomorphic to $H(\mathbb{Z}_m, \{0, \dots, d-1\})$. The rotations below also apply when $8 \mid d$, completing the odd m case. For $d = 4$, the graph is the circulant $C_{2m}(1, 3)$, and our rotations recover its known toroidal quadrangulation [3, Proposition 5].

Write $d = 2h$, where h is even, and using additive notation in $\mathbb{Z}_m$, set

$$S = \{0, 1, \dots, 2h-1\}, \quad \rho_B = (0 \ 1 \ \dots \ h-1 \ 2h-1 \ 2h-2 \ \dots \ h), \quad \rho_A = \rho_B^{h-1}. \quad (9)$$

The labels in S are distinct modulo m . Since ρ_B is a $2h$ -cycle and h is even, $\gcd(2h, h-1) = \gcd(2, h-1) = 1$. For a k -cycle π , it is standard that π^N has exactly $\gcd(k, N)$ cycles each of length $k/\gcd(k, N)$. Thus our previous computation shows ρ_A is also a $2h$ -cycle. The first half of the ρ_B -cycle increases from 0 to $h-1$, and the second decreases from $2h-1$ to h . Thus labels h positions apart sum to $2h-1$. Put $\sigma(s) = 2h-1-s$. Then

$$\rho_B^h = \sigma, \quad \rho_A \rho_B = \sigma, \quad \rho_B \sigma = \sigma \rho_B.$$

The permutation σ has no fixed points, since $2s$ is an even integer it cannot equal $2h-1$.

Proposition 5.1. *Let $m \geq d \geq 4$ with $4 \mid d$. The rotations in (9) give an orientable quadrangulation of $H(\mathbb{Z}_m, S)$, with m vertices in each class, degree d , and simple 4-cycles as face boundaries.*

Proof. The graph is simple, bipartite, and d -regular. It is connected because S contains the identity 0 and the generator 1 of $\mathbb{Z}_m$. Both local rotations are d -cycles. In additive notation, (4) gives

$$F^2(x, s)_A = (x + s - \rho_B(s), \sigma(s))_A. \quad (10)$$

Applying this twice yields

$$F^4(x, s)_A = (x + s - \rho_B(s) + \sigma(s) - \rho_B(\sigma(s)), s)_A.$$

Since σ commutes with ρ_B ,

$$F^4(x, s)_A = (x + \underbrace{s + \sigma(s)}_{2h-1} - \underbrace{(\rho_B(s) + \sigma(\rho_B(s)))}_{2h-1}, s)_A.$$

Thus $F^4(x, s)_A = (x, s)_A$. Since F interchanges A -darts and B -darts and σ has no fixed points, every facial orbit has length exactly four. The face starting with $(x, s)_A$ has vertices

$$A_x, \quad B_{x+s}, \quad A_{x+s-\rho_B(s)}, \quad B_{x+(2h-1)-\rho_B(s)}. \quad (11)$$

The two A -vertices are distinct because s and $\rho_B(s)$ are distinct modulo m . If the two B -vertices were equal, then $\rho_B(s) \equiv \sigma(s) \pmod{m}$. Both labels belong to S , so this would imply $\rho_B(s) = \sigma(s) = \rho_B^h(s)$. Hence $\rho_B^{h-1}(s) = s$, which is impossible because ρ_B is a $2h$ -cycle and $0 < h-1 < 2h$. $\square$

For $m = 7$ and $d = 4$, the graph is shown in Figure 4.

Figure 4: The cyclic group example $H(\mathbb{Z}_7, \{0, 1, 2, 3\}) \cong C_{14}(1, 3)$.

For a nonorientable quadrangulation, Euler’s formula gives genus $k = 2 + m(d - 4)/2$, so $2 \mid md$ is necessary. However, it is also insufficient: for $(m, d) = (5, 4)$, the only graph is $K_{5,5}$ minus a perfect matching, which has no nonorientable quadrangulation [5, Theorem 3], although Proposition 5.1 gives an orientable one. The nonorientable genus of nearly complete bipartite graphs is known [5, 10]. An extension of Pisanski’s conjecture could ask which pairs $m \geq d \geq 4$ admit a regular bipartite nonorientable quadrangulation.

We note that the freedom to choose the connection set is useful in this problem. We can choose its elements together with the local rotations so that the facial walks have length four and both local permutations are cyclic. In the cyclic construction, the same label permutations work for every $m \geq d$. This motivates the question of whether similar choices give minimum-genus embeddings for other families of Haar graphs.

On the other hand, for any fixed Haar graph the connection set is fixed, and satisfying the conditions that make every facial walk return to its starting dart after exactly 4 steps, while keeping each local permutation as a single cycle, can be much more difficult. It would be useful to find structural conditions on a connection set that guarantee suitable cyclic permutations. One can also ask which changes to the label permutations preserve facial closure while combining their cycles. Such conditions or operations could extend these constructions to prescribed graph families whose genus is unknown.

References

- [1] S. R. Alpert and J. L. Gross, *Components of branched coverings of current graphs*, Journal of Combinatorial Theory, Series B **20** (1976), no. 3, 283–303.
- [2] I. Anderson, *Quadrilateral embeddings of bipartite graphs*, Journal of Graph Theory **5** (1981), no. 4, 355–361.
- [3] M. Conder and R. Grande, *On embeddings of circulant graphs*, Electronic Journal of Combinatorics **22** (2015), no. 2, Paper P2.28.
- [4] I. Estélyi and T. Pisanski, *Which Haar graphs are Cayley graphs?*, Electronic Journal of Combinatorics **23** (2016), no. 3, Paper P3.10.
- [5] B. Mohar, *Nonorientable genus of nearly complete bipartite graphs*, Discrete & Computational Geometry **3** (1988), 137–146.
- [6] B. Mohar, T. D. Parsons, and T. Pisanski, *The genus of nearly complete bipartite graphs*, Ars Combinatoria **20B** (1985), 173–183.
- [7] B. Mohar and C. Thomassen, *Graphs on Surfaces*, Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, Baltimore, 2001.
- [8] T. Pisanski, *Genus of Cartesian products of regular bipartite graphs*, Journal of Graph Theory **4** (1980), no. 1, 31–42.
- [9] G. Ringel, *Das Geschlecht des vollständigen paaren Graphen*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg **28** (1965), 139–150.
- [10] W. Singh and T. Sun, *Settling the nonorientable genus of the nearly complete bipartite graphs*, Graphs and Combinatorics **39** (2023), no. 5, Article 96.
- [11] A. Ulrigg and A. Metzger, *The genus of bipartite Kneser graphs*, arXiv:2610.03159 (2026).
- [12] A. T. White, *On the genus of the composition of two graphs*, Pacific Journal of Mathematics **41** (1972), no. 1, 275–279.
- [13] J. W. T. Youngs, *Minimal imbeddings and the genus of a graph*, Journal of Mathematics and Mechanics **12** (1963), no. 2, 303–315.