

## Solutions to Final Exam


1. (a) True. Periodic  $\xleftrightarrow{\mathcal{F}}$  Discrete.
- (b) True.
- (c) False.  $\because \omega_s < 2 * 3$ .
- (d) False.  $\because X(e^{j(\omega+2\pi)}) \neq X(e^{j\omega})$ .
- (e) False.  $x(t)$  should be real.
- (f) False. “real and odd”  $\xleftrightarrow{\mathcal{F}}$  “purely imaginary and odd”.
- (g) True.  $\because x[0] = 0$ .
- (h) True.  $\because \sum_{n=-\infty}^{\infty} x[n] = 0$ .
- (i) True.
- (j) False. “real and even”  $\xleftrightarrow{\mathcal{F}}$  “real and odd”.

2. Let  $Y(j\omega) = \text{sinc}\left(\frac{\omega}{2\pi}\right)$ . Since

$$X(j\omega) = \text{sinc}^2\left(\frac{\omega}{2\pi}\right) = Y(j\omega) \cdot Y(j\omega)$$

we have

$$x(t) = y(t) * y(t) = \text{rect}(t) * \text{rect}(t) = \begin{cases} 1+t, & -1 \leq t \leq 0 \\ 1-t, & 0 \leq t \leq 1 \\ 0, & \text{otherwise.} \end{cases}$$



3. Since

$$\text{sgn}(t) \xleftrightarrow{\mathcal{F}} \frac{2}{j\omega}$$

we have

$$\frac{2}{jt} \xleftrightarrow{\mathcal{F}} 2\pi \text{sgn}(-\omega) = -2\pi \text{sgn}(\omega)$$

according to the duality. Hence,

$$x(t) = \frac{1}{\pi t} \xleftrightarrow{\mathcal{F}} -j \text{sgn}(\omega).$$

4. (a) We know

$$\text{sgn}[n] \xleftrightarrow{\mathcal{F}} 2/(1 - e^{-j\omega}).$$

Since  $u[n] = (1 + \text{sgn}[n])/2$ , we have

$$U(e^{j\omega}) = \frac{1}{2} \left( \sum_{k=-\infty}^{\infty} 2\pi\delta(\omega - 2\pi k) + \frac{2}{1 - e^{-j\omega}} \right) = \frac{1}{1 - e^{-j\omega}} + \pi \sum_{k=-\infty}^{\infty} \delta(\omega - 2\pi k).$$

(b) Let

$$y[n] = \sum_{m=-\infty}^n x[m] = x[n] * u[n].$$

We have the Fourier transform of  $y[n]$  given by

$$\begin{aligned} X(e^{j\omega})U(e^{j\omega}) &= \frac{1}{1 - e^{-j\omega}}X(e^{j\omega}) + \pi \sum_{k=-\infty}^{\infty} X(e^{j\omega})\delta(\omega - 2\pi k) \\ &= \frac{1}{1 - e^{-j\omega}}X(e^{j\omega}) + \pi \sum_{k=-\infty}^{\infty} X(e^{j2\pi k})\delta(\omega - 2\pi k) \\ &= \frac{1}{1 - e^{-j\omega}}X(e^{j\omega}) + \pi X(e^{j0}) \sum_{k=-\infty}^{\infty} \delta(\omega - 2\pi k). \end{aligned}$$

5. From (iii),

$$\begin{aligned} x_o[n] \xleftrightarrow{\mathcal{F}} j\Im\{X(e^{j\omega})\} &= j\sin(\omega) - j\sin(2\omega) \\ &= \frac{1}{2}e^{j\omega} - \frac{1}{2}e^{-j\omega} - \frac{1}{2}e^{2j\omega} + \frac{1}{2}e^{-2j\omega}. \end{aligned}$$

Hence,

$$x_o[n] = \frac{1}{2}\delta[n+1] - \frac{1}{2}\delta[n-1] - \frac{1}{2}\delta[n+2] + \frac{1}{2}\delta[n-2].$$

Since  $x_o[n] = (x[n] + x[-n])/2$  and (i), we have, for  $n < 0$ ,

$$x[n] = 2x_o[n] = \delta[n+1] - \delta[n-1] - \delta[n+2] + \delta[n-2] = \delta[n+1] - \delta[n+2].$$

Furthermore, according to Parseval's relation and (iv), we have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega = \sum_{n=-\infty}^{\infty} |x[n]|^2 = |x[0]|^2 + 2 = 3$$

implying

$$x[0] = \pm 1.$$

From (ii), we have  $x[0] = 1$ . Therefore,

$$x[n] = \delta[n] + \delta[n+1] - \delta[n+2].$$


6. (a) The Fourier transform of the impulse train  $p[n]$  is

$$P(e^{j\omega}) = \frac{2\pi}{4} \sum_{k=-\infty}^{\infty} \delta\left(\omega - \frac{2k\pi}{4}\right) = \frac{\pi}{2} \sum_{k=-\infty}^{\infty} \delta\left(\omega - \frac{k\pi}{2}\right).$$

- (b) The Fourier transform of  $w[n]$  is

$$\begin{aligned} W(e^{j\omega}) &= \frac{1}{2\pi} \int_{2\pi} P(e^{j\theta}) X(e^{j(\omega-\theta)}) d\theta \\ &= \frac{1}{2\pi} \int_{-\frac{\pi}{4}}^{\frac{7\pi}{4}} \frac{\pi}{2} \sum_{k=-\infty}^{\infty} \delta\left(\theta - \frac{k\pi}{2}\right) X(e^{j(\omega-\theta)}) d\theta \\ &= \frac{1}{4} \int_{-\frac{\pi}{4}}^{\frac{7\pi}{4}} \sum_{k=0}^3 \delta\left(\theta - \frac{k\pi}{2}\right) X(e^{j(\omega-\theta)}) d\theta \\ &= \frac{1}{4} \sum_{k=0}^3 X\left(e^{j(\omega - \frac{k\pi}{2})}\right). \end{aligned}$$

The spectrum is depicted below.



7. Let

$$x_1(t) = A \cdot \text{rect}\left(\frac{t}{T}\right) \xleftrightarrow{\mathcal{F}} X_1(j\omega) = AT \text{sinc}\left(\frac{\omega T}{2\pi}\right)$$

and

$$x_2(t) = \cos(\omega_c t) \xleftrightarrow{\mathcal{F}} X_2(j\omega) = \pi\delta(\omega - \omega_c) + \pi\delta(\omega + \omega_c).$$

The Fourier transform of  $x(t)$  is

$$\begin{aligned} X(j\omega) &= \frac{1}{2\pi} X_1(j\omega) * X_2(j\omega) \\ &= \frac{AT}{2} \left[ \text{sinc}\left(\frac{\omega T}{2\pi}\right) * \delta(\omega - \omega_c) + \text{sinc}\left(\frac{\omega T}{2\pi}\right) * \delta(\omega + \omega_c) \right] \\ &= \frac{AT}{2} \text{sinc}\left(\frac{(\omega - \omega_c)T}{2\pi}\right) + \frac{AT}{2} \text{sinc}\left(\frac{(\omega + \omega_c)T}{2\pi}\right). \end{aligned}$$