

ON RANK-MONOTONE GRAPH OPERATIONS AND MINIMAL OBSTRUCTION GRAPHS FOR THE LOVÁSZ–SCHRIJVER SDP HIERARCHY

YU HIN (GARY) AU AND LEVENT TUNÇEL

ABSTRACT. We study the lift-and-project rank of the stable set polytopes of graphs with respect to the Lovász–Schrijver SDP operator LS_+ , with a particular focus on finding and characterizing the smallest graphs with a given LS_+ -rank (the needed number of iterations of the LS_+ operator on the fractional stable set polytope to compute the stable set polytope). We introduce a generalized vertex-stretching operation that appears to be promising in generating LS_+ -minimal graphs and study its properties. We also provide several new LS_+ -minimal graphs, most notably the first known instances of 12-vertex graphs with LS_+ -rank 4, which provides the first advance in this direction since Escalante, Montelar, and Nasini’s discovery of a 9-vertex graph with LS_+ -rank 3 in 2006.

1. INTRODUCTION

Given a simple, undirected graph $G = (V(G), E(G))$, we say that $S \subseteq V(G)$ is a *stable set* if no two vertices in S are joined by an edge. The (*maximum*) *stable set problem*, which aims to find a stable set of maximum cardinality in a given graph G , is one of the most well-studied problems in combinatorial optimization. While this problem is \mathcal{NP} -hard, a standard approach for tackling the problem is to associate stable sets of G with points in $\mathbb{R}^{V(G)}$, and model it as a convex optimization problem. Given a set $S \subseteq V(G)$, its *incidence vector* $\chi_S \in \{0, 1\}^{V(G)}$ is defined so that $[\chi_S]_i = 1$ if $i \in S$, and $[\chi_S]_i = 0$ otherwise. Then we define the *stable set polytope* of a given graph G to be the convex hull of the incidence vectors of stable sets of G :

$$\text{STAB}(G) := \text{conv}(\{\chi_S : S \subseteq V(G) \text{ is a stable set of } G\}).$$

Observe that if we let $\alpha(G)$ be the cardinality of a maximum stable set in G , then

$$(1) \quad \alpha(G) = \max \left\{ \sum_{i \in V(G)} x_i : x \in \text{STAB}(G) \right\}.$$

While (1) is a linear program, considering again that the underlying combinatorial problem is \mathcal{NP} -hard, it is a difficult task to find an explicit description (e.g., via listing its facets) of $\text{STAB}(G)$ for a general graph G . This naturally leads to the pursuit of “nice” convex relaxations of $\text{STAB}(G)$. Below we list several desirable characteristics of such a convex relaxation P :

Date: November 28, 2024.

Key words and phrases. stable set problem, lift and project, combinatorial optimization, semidefinite programming, integer programming.

Yu Hin (Gary) Au: Corresponding author. Department of Mathematics and Statistics, University of Saskatchewan, Saskatoon, Saskatchewan, S7N 5E6 Canada. E-mail: gary.au@usask.ca.

Levent Tunçel: Research of this author was supported in part by an NSERC Discovery Grant. Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, N2L 3G1 Canada. E-mail: levent.tuncel@uwaterloo.ca.

- (i) $P \cap \{0, 1\}^{V(G)} = \text{STAB}(G) \cap \{0, 1\}^{V(G)}$. That is, a 0-1 vector is in P if and only if it is the incidence vector of a stable set in G .
- (ii) P is tractable. That is, one can optimize a linear function over P with arbitrary precision in polynomial time.
- (iii) P is a “strong” relaxation. This is a relatively subjective measure, and can mean that important families of valid inequalities of $\text{STAB}(G)$ are also valid for P , and/or that $\max \left\{ \sum_{i \in V(G)} x_i : x \in P \right\}$ is “close” to $\alpha(G)$.

One of the simplest convex relaxations of $\text{STAB}(G)$ is the *fractional stable set polytope*

$$\text{FRAC}(G) := \left\{ x \in [0, 1]^{V(G)} : x_i + x_j \leq 1, \forall \{i, j\} \in E(G) \right\}.$$

While $\text{FRAC}(G)$ satisfies properties (i) and (ii), it is rather weak in general. For a stronger relaxation, we call $C \subseteq V(G)$ a *clique* if every pair of vertices in C is joined by an edge. Then notice that, for every clique C , the inequality

$$\sum_{i \in C} x_i \leq 1$$

is valid for $\text{STAB}(G)$. Thus, if we define the *clique polytope*

$$\text{CLIQ}(G) := \left\{ x \in [0, 1]^{V(G)} : \sum_{i \in C} x_i \leq 1 \text{ for every clique } C \subseteq V(G) \right\},$$

then $\text{STAB}(G) \subseteq \text{CLIQ}(G) \subseteq \text{FRAC}(G)$ for every graph G . (For the second containment, observe that every edge is a clique of size 2.) However, while $\text{CLIQ}(G)$ is a stronger relaxation than $\text{FRAC}(G)$, it is not tractable in general.

In this manuscript, we focus on semidefinite relaxations of $\text{STAB}(G)$ produced by LS_+ , a lift-and-project operator devised by Lovász and Schrijver [LS91] which we fully define in Section 2. (The operator has also been referred to as N_+ in the literature.) Given a graph G , the LS_+ operator generates a sequence of relaxations $\text{LS}_+^k(G)$ which satisfies

$$\text{FRAC}(G) =: \text{LS}_+^0(G) \supseteq \text{LS}_+^1(G) \supseteq \text{LS}_+^2(G) \supseteq \cdots \supseteq \text{LS}_+^{|V(G)|}(G) = \text{STAB}(G).$$

(We will usually refer to $\text{LS}_+^1(G)$ as simply $\text{LS}_+(G)$.) When $k = O(1)$, $\text{LS}_+^k(G)$ can be described as the feasible region of a semidefinite program whose number of variables and constraints are polynomial in the size of the number of vertices and edges in G , and thus the relaxation is indeed tractable in this case. Moreover, the first relaxation $\text{LS}_+(G)$ already satisfies many well-known families of valid inequalities of $\text{STAB}(G)$, including (among others) the aforementioned clique inequalities, orthogonality constraints imposed by the Lovász theta body [Lov79], as well as odd hole, odd antihole, and odd wheel constraints [LS91].

The hierarchy of relaxations generated by LS_+ gives rise to the notion of the LS_+ -rank of a graph G , which is defined to be the smallest integer k where $\text{LS}_+^k(G) = \text{STAB}(G)$, and gives us a measure of how difficult the stable set problem is for the LS_+ operator. It is well-known that a graph G has LS_+ -rank 0 (i.e., satisfies $\text{FRAC}(G) = \text{STAB}(G)$) if and only if G is bipartite. Some families of graphs that are known to have LS_+ -rank 1 (i.e., satisfy $\text{LS}_+(G) = \text{STAB}(G)$) include — but are not limited to — odd cycles, odd antihole, odd wheels, and perfect graphs (which are defined to be the graphs with $\text{CLIQ}(G) = \text{STAB}(G)$). In the last decade, considerable progress has been made in finding a combinatorial characterization of graphs with LS_+ -rank 1 — see, for instance, [BENT13, BENT17, Wag22, BENW23].

Nevertheless, since the maximum stable problem is \mathcal{NP} -hard, there has to be graphs with unbounded LS_+ -rank. The first family of graphs that have unbounded LS_+ -rank was obtained

by Stephen and the second author [ST99], who showed that the line graph of the complete graph on $2k+1$ vertices has LS_+ -rank k , giving a family of graphs G whose LS_+ -rank is asymptotically $\Omega(\sqrt{|V(G)|})$. On the other hand, Lipták and the second author [LT03] showed the following:

Theorem 1. *For every graph G , the LS_+ -rank of G is at most $\left\lfloor \frac{|V(G)|}{3} \right\rfloor$.*

This naturally raises the following question: For every integer $\ell \geq 1$, is there a graph on 3ℓ vertices which has LS_+ -rank ℓ ? If these graphs exist, their extremal nature (in terms of being the smallest possible graphs with a given LS_+ -rank) may help reveal the critical structures that expose the limitations of these LS_+ -relaxations. This understanding could be extremely helpful when it comes to analyzing other convex relaxations of the maximum stable set problem, particularly those which are produced by other lift-and-project methods.

This direction of investigation was already set in the seminal paper [LS91] and questions about the behaviour of LS_+ -rank under simple graph operations were also raised in [GT01]. In the same general direction of research, Laurent [Lau02] analyzed the LS_+ -rank and related ranks in the context of the maximum cut problem by establishing nice behaviour (only in the context of maximum cut problems) of the underlying lift-and-project operators under graph minor operations; also see [Lau03] for an analysis of the Lasserre operator. However, as it was illustrated in some depth in [LT03], the LS_+ -rank of a graph does not behave in a nice, uniform way under the usual graph minor operations for the stable set problem. Therefore, a deeper investigation is necessary to construct the kind of graph operations which would be helpful in discovering and understanding minimal obstructions to tractable convex relaxations of the stable set polytope obtained by LS_+ or other convex optimization based lift-and-project hierarchies. Overall, the importance of the quest to understand minimal obstructions to families of SDP relaxations in particular — and convex relaxations in general — has been raised by many others. For example, Knuth, in his well-known survey “The Sandwich Theorem” [Knu94] poses six open problems in the general context of Lovász theta function. Two of the six open problems concern $LS_+(\text{FRAC}(G))$. One of them asks for finding what we call below a 2-minimal graph (answered in [LT03]).

We say that a graph G is ℓ -minimal if $|V(G)| = 3\ell$ and G has LS_+ -rank ℓ . It is known that ℓ -minimal graphs exist for $\ell \in \{1, 2, 3\}$. For $\ell = 1$, it is easy to see that the 3-cycle is the only 1-minimal graph. The first 2-minimal graph ($G_{2,1}$ in Figure 1) was found by Lipták and the second author [LT03], who also conjectured that ℓ -minimal graphs exist for all $\ell \in \mathbb{N}$. Subsequently, Escalante, Montelar, and Nasini [EMN06] showed that there is only one other 2-minimal graph ($G_{2,2}$ in Figure 1), while providing the first example of a 3-minimal graph ($G_{3,1}$ in Figure 1). (The logic behind the seemingly odd choice of vertex labels in the figures of this section will be explained in Section 4 when we introduce the vertex-stretching operation.)

In producing the first 3-minimal graph, Escalante et al. [EMN06] also showed that there does not exist an ℓ -minimal graph for any $\ell \geq 4$ if we restrict ourselves to graphs that can be obtained by starting with a complete graph and replacing every edge by a path of length at least 1. (Let K_n denote the complete graph on n vertices. Notice that $G_{2,1}$ and $G_{3,1}$ can be respectively obtained from K_4 and K_5 by replacing some edges with paths of length 3.)

Recently, the authors [AT24] discovered several family of graphs G which has LS_+ -rank $\Omega(|V(G)|)$. One of them is the family of graphs H_k , which is defined as follows. Given $k \in \mathbb{N}$, let $[k]$ denote the set $\{1, 2, \dots, k\}$. For every $k \geq 3$, let

$$V(H_k) := \{i_0, i_1, i_2 : i \in [k]\}$$

and

$$E(H_k) := \{\{i_1, i_0\}, \{i_0, i_2\} : i \in [k]\} \cup \{\{i_1, i_2\} : i, j \in [k], i \neq j\}.$$




FIGURE 1. Known 2- and 3-minimal graphs due to [LT03] and [EMN06]



FIGURE 2. Several graphs in the family H_k

Figure 2 illustrates the graphs H_k for $k = 3, 4, 5$. (Note that our vertex labels for H_k are different from those in [AT24].) The authors [AT24, Theorem 2] proved the following.

Theorem 2. *For every $k \geq 3$, The LS_+ -rank of H_k is at least $\frac{3k}{16}$.*

Theorem 2 (and other results in [AT24]) ended a 17-year lull in new hardness results for LS_+ -relaxations of the stable set problem, and provides renewed hope that ℓ -minimal graphs do exist for $\ell \geq 4$. Indeed, one of the main contributions of this work is the discovery of what we believe to be the first known instance of a 4-minimal graph ($G_{4,1}$ in Figure 3).

This paper is organized as follows. In Section 2, we define the LS_+ operator and introduce some of the tools and notation we will need for our subsequent analysis. Then, in Section 3, we discuss what we call *star-homomorphism* between graphs, and provide a template for constructing graph operations that are LS_+ -rank non-decreasing. Using this template, we define our *vertex-stretching* operation in Section 4, which generalizes similar graph operations studied previously [LT03, AEF14, BENT17]. We then show in Section 5 that every ℓ -minimal graph for $\ell \geq 2$ must be obtained from applying our vertex-stretching operation to a smaller graph, and in particular study the LS_+ -ranks of graphs obtained from stretching the vertices of a complete graph. In Section 6, we prove that $G_{4,1}$ is indeed 4-minimal and discuss some of the immediate consequences of the result, which includes the discovery of several other new 3- and 4-minimal graphs. We do remark that while the discovery of $G_{4,1}$ is largely guided by the structural results for ℓ -minimal graphs presented in Sections 3 to 5, these results are not needed for the proof of $G_{4,1}$ having LS_+ -rank 4. After that, we revisit the aforementioned families of graphs H_k

FIGURE 3. $G_{4,1}$, a 12-vertex graph with LS_+ -rank 4

in Section 7, and apply our results on vertex stretching to show that there exists a family of graphs G with maximum degree 3 and LS_+ -rank $\Omega(\sqrt{|V(G)|})$. Finally, we conclude our paper in Section 8 by mentioning some natural future research directions.

2. PRELIMINARIES

In this section, we define the lift-and-project operator LS_+ due to Lovász and Schrijver [LS91] and the convex relaxations of $\text{STAB}(G)$ it produces, as well as go over the basic tools we will use in subsequent sections to analyze the LS_+ -rank of graphs.

2.1. The LS_+ -operator. Given a set $P \subseteq [0, 1]^n$, we define the *homogenized cone* of P to be

$$\text{cone}(P) := \left\{ \begin{bmatrix} \lambda \\ \lambda x \end{bmatrix} : \lambda \geq 0, x \in P \right\}.$$

Notice that $\text{cone}(P) \subseteq \mathbb{R}^{n+1}$, and we will refer to the new coordinate with index 0. Next, given a vector x and an index i , we may refer to the i -entry in x by x_i or $[x]_i$. All vectors are column vectors by default, so x^\top , the transpose of a vector x , is a row vector. Next, let \mathbb{S}_+^n denote the set of n -by- n real symmetric positive semidefinite matrices, and $\text{diag}(Y)$ be the vector formed by the diagonal entries of a square matrix Y . We also let e_i be the i^{th} unit vector.

Given $P \subseteq [0, 1]^n$, the operator LS_+ first *lifts* P to the following set of matrices:

$$\widehat{\text{LS}}_+(P) := \{Y \in \mathbb{S}_+^{n+1} : Y e_0 = \text{diag}(Y), Y e_i, Y(e_0 - e_i) \in \text{cone}(P) \forall i \in [n]\}.$$

It then *projects* the set back down to the following set in \mathbb{R}^n :

$$\text{LS}_+(P) := \left\{ x \in \mathbb{R}^n : \exists Y \in \widehat{\text{LS}}_+(P), Y e_0 = \begin{bmatrix} 1 \\ x \end{bmatrix} \right\}.$$

Given $x \in \text{LS}_+(P)$, we say that $Y \in \widehat{\text{LS}}_+(P)$ is a *certificate matrix* for x if $Y e_0 = \begin{bmatrix} 1 \\ x \end{bmatrix}$. Also, given a set $P \subseteq [0, 1]^n$, we define

$$P_I := \text{conv}(P \cap \{0, 1\}^n)$$

to be the *integer hull* of P . The following foundational property of LS_+ was shown in [LS91].

Lemma 3. *For every set $P \subseteq [0, 1]^n$, $P_I \subseteq \text{LS}_+(P) \subseteq P$.*

Proof. For the first containment, let $x \in P \cap \{0, 1\}^n$. Observe that $Y := \begin{bmatrix} 1 \\ x \end{bmatrix} \begin{bmatrix} 1 \\ x \end{bmatrix}^\top \in \widehat{\text{LS}}_+(P)$, and so $x \in \text{LS}_+(P)$. For the second containment, let $x \in \text{LS}_+(P)$, and $Y \in \widehat{\text{LS}}_+(P)$ be a certificate matrix for x . Since $Ye_0 = Ye_i + Y(e_0 - e_i)$ for any index $i \in [n]$ and that $\widehat{\text{LS}}_+$ imposes that $Ye_i, Y(e_0 - e_i) \in \text{cone}(P)$, it follows that $Ye_0 = \begin{bmatrix} 1 \\ x \end{bmatrix} \in \text{cone}(P)$, and thus $x \in P$. \square

Therefore, $\text{LS}_+(P)$ contains the same set of integral solutions as P . Also, if P is tractable, then so is $\text{LS}_+(P)$, and it is known that $P \supset \text{LS}_+(P)$ unless $P = P_I$. Thus, $\text{LS}_+(P)$ offers a tractable relaxation of P_I that is tighter than the initial relaxation P .

Furthermore, we can apply LS_+ multiple times to obtain yet tighter relaxations. Given $k \in \mathbb{N}$, let $\text{LS}_+^k(P)$ be the set obtained from applying k successive LS_+ operations to P . (We also let $\text{LS}_+^0(P) := P$.) Then it is well-known that

$$P_I = \text{LS}_+^n(P) \subseteq \text{LS}_+^{n-1}(P) \subseteq \cdots \subseteq \text{LS}_+(P) \subseteq P.$$

Thus, LS_+ generates a hierarchy of progressively tighter convex relaxations which converge to P_I in no more than n iterations. The reader may refer to Lovász and Schrijver [LS91] for a proof of this fact and some other properties of the LS_+ operator.

2.2. Analyzing the LS_+ -rank of a graph. Recall that $\text{FRAC}(G)$, the fractional stable set polytope of a graph G , offers a simple and tractable convex relaxation of $\text{STAB}(G)$. Thus, we could apply LS_+ to obtain stronger relaxations of $\text{STAB}(G)$ than $\text{FRAC}(G)$. Given an integer $k \geq 0$, define

$$\text{LS}_+^k(G) := \text{LS}_+^k(\text{FRAC}(G)),$$

and let $r_+(G)$ denote the LS_+ -rank of G (which, again, is the smallest integer k where $\text{LS}_+^k(G) = \text{STAB}(G)$).

To show that a graph G has LS_+ -rank at least p , the standard approach is to find a point \bar{x} where $\bar{x} \notin \text{STAB}(G)$ and $\bar{x} \in \text{LS}_+^{p-1}(G)$ — this is the approach we will take when verifying that $r_+(G_{4,1}) \geq 4$. We do remark that verifying $\bar{x} \in \text{LS}_+^{p-1}(G)$ tends to get progressively more challenging as p increases, unless the symmetries of G allow for an inductive argument (which is the case for the line graphs of odd cliques [ST99], and to a lesser extent for H_k and related graphs [AT24]). Given $p \in \mathbb{N}$, we also define

$$\alpha_{\text{LS}_+^p}(G) := \max \left\{ \bar{e}^\top x : x \in \text{LS}_+^p(G) \right\},$$

where \bar{e} denotes the vector of all-ones. Notice that if $\alpha_{\text{LS}_+^p}(G) > \alpha(G)$, then $r_+(G) \geq p + 1$.

Next, the following is a well-known property of LS_+ .

Lemma 4. *Let $P \subseteq [0, 1]^n$ be a polyhedron, and F be a face of $[0, 1]^n$. Then*

$$\text{LS}_+(P \cap F) = \text{LS}_+(P) \cap F.$$

It follows from Lemma 4 that if $\bar{x} \in \text{LS}_+^p(G)$ and G' is an induced subgraph of G , then the vector obtained from \bar{x} by removing entries not in $V(G')$ is in $\text{LS}_+^p(G')$. This in turn implies that $r_+(G') \leq r_+(G)$ — see, for instance, [AT24, Lemma 5] for a proof.

We next mention several other ways of bounding $r_+(G)$ using the LS_+ -rank of graphs that are related to G . Given a graph G and $S \subseteq V(G)$, we let $G - S$ denote the subgraph of G induced by the vertices $V(G) \setminus S$, and call $G - S$ the graph obtained by the *deletion* of S . (When $S = \{i\}$ for some vertex i , we simply write $G - i$ instead of $G - \{i\}$.) Next, given $i \in V(G)$, let

$$\Gamma_G(i) := \{j \in V(G) : \{i, j\} \in E(G)\}$$

be the *open neighborhood* of i in G , and $\Gamma_G[i] := \Gamma_G(i) \cup \{i\}$ be the *closed neighborhood* of i in G . Then the graph obtained from the *destruction* of i in G is defined as

$$G \ominus i := G - \Gamma[i].$$

Then we have the following.

Theorem 5. *For every graph G ,*

- (i) [LS91, Corollary 2.16] $r_+(G) \leq \max \{r_+(G \ominus i) : i \in V(G)\} + 1$;
- (ii) [LT03, Theorem 36] $r_+(G) \leq \min \{r_+(G - i) : i \in V(G)\} + 1$.

Recall that $r_+(G) = 0$ if and only if G is bipartite (in which case $\text{FRAC}(G) = \text{STAB}(G)$). Thus, it follows immediately from Theorem 5(ii) that if G is non-bipartite but $G - i$ is bipartite for some $i \in V(G)$, then $r_+(G) = 1$ — an example for such graphs is the odd cycles. Likewise, if G is non-bipartite while $G \ominus i$ is bipartite for every $i \in V(G)$, then $r_+(G) = 1$ as well — such as when G is an odd antihole (i.e., the graph complement of an odd cycle of length at least 5), or an odd wheel (i.e., the graph obtained from joining a vertex to every vertex of an odd cycle of length at least 5).

We say that a graph G is *perfect* if $\text{CLIQ}(G) = \text{STAB}(G)$. In terms of forbidden subgraphs, G is perfect if and only if it does not contain an induced subgraph that is an odd hole (i.e., an odd cycle of length at least 5) or an odd antihole [CRST06]. Since $\text{LS}_+(G) \subseteq \text{CLIQ}(G)$ in general [LS91], it follows that $r_+(G) \leq 1$ if G is perfect.

The following is a restatement of [LT03, Lemma 5].

Proposition 6. *Let G be a graph, and $S_1, S_2, C \subseteq V(G)$ are mutually disjoint subsets such that*

- $S_1 \cup S_2 \cup C = V(G)$;
- C induces a clique in G ;
- *There is no edge $\{i, j\} \in E(G)$ where $i \in S_1, j \in S_2$.*

Then $r_+(G) = \max \{r_+(G - S_1), r_+(G - S_2)\}$.

Thus, if G has a cut clique (i.e., a clique C where $G - C$ has multiple components), then the LS_+ -rank of G is equal to that of one of its proper subgraphs.

Next, observe that $\text{FRAC}(G)$ is *lower-comprehensive* for every graph G (i.e., if $x \in \text{FRAC}(G)$, then $y \in \text{FRAC}(G)$ for all vectors y where $0 \leq y \leq x$). Also, one can show (such as using Lemma 4) that LS_+ preserves lower-comprehensiveness. Thus, it follows that $\text{LS}_+^k(G)$ is lower-comprehensive as well for every $k \in \mathbb{N}$.

Finally, in addition to preserving lower-comprehensiveness, it is also clear from the definition of LS_+ that the operator also preserves containment (i.e., if $P_1 \subseteq P_2$, then $\text{LS}_+(P_1) \subseteq \text{LS}_+(P_2)$). This implies the following.

Lemma 7. *Given graphs G, H where $V(G) = V(H)$ and $E(G) \subseteq E(H)$,*

- (i) *If $a^\top x \leq \beta$ is valid for $\text{LS}_+^p(G)$, then $a^\top x \leq \beta$ is valid for $\text{LS}_+^p(H)$;*
- (ii) *If $a^\top x \leq \beta$ is not valid for $\text{LS}_+^p(H)$, then $a^\top x \leq \beta$ is not valid for $\text{LS}_+^p(G)$.*

3. STAR-HOMOMORPHIC GRAPHS

In this section, we introduce the notion of two graphs being star-homomorphic, and describe how the LS_+ -relaxations of such a pair of graphs are related. Given a graph $G = (V(G), E(G))$, we define the graph G^\dagger where

$$\begin{aligned} V(G^\dagger) &:= \{i, \bar{i} : i \in V(G)\}, \\ E(G^\dagger) &:= E(G) \cup \{\{i, \bar{i}\} : i \in V(G)\}. \end{aligned}$$

In other words, we obtain G^\dagger from G by adding a new vertex \bar{i} for every $i \in V(G)$, and then adding an edge between \bar{i} and i . Figure 4 provides an example of constructing G^\dagger from G .




FIGURE 4. Constructing G^\dagger from G

Also, given graphs G and H , we say that $g : V(H) \rightarrow V(G)$ is a *homomorphism* if, for all $i, j \in V(H)$,

$$\{i, j\} \in E(H) \Rightarrow \{g(i), g(j)\} \in E(G).$$

Furthermore, given graphs G and H , if there exists a homomorphism $g : V(H) \rightarrow V(G^\dagger)$, then we say that H is *star-homomorphic to G* under g .




FIGURE 5. Two graphs H_1, H_2 that are star-homomorphic to G

Example 8. Consider the graphs in Figure 5. Then H_1 is star-homomorphic to G under g_1 where

$j \in V(H_1)$	1	2	3	4	5	6	7	8	9
$g_1(j) \in V(G^\dagger)$	1	2	3	4	5	6	6	6	6

Likewise, H_2 is star-homomorphic to G under g_2 where

$j \in V(H_2)$	1	2	3	4	5	6	7	8
$g_2(j) \in V(G^\dagger)$	1	2	3	4	5	6	3	6

Given that H is star-homomorphic to G under g and $x \in \mathbb{R}^{V(G)}$, we let $\tilde{g}(x) \in \mathbb{R}^{V(H)}$ be the vector where

$$(2) \quad [\tilde{g}(x)]_j := \begin{cases} x_i & \text{if } g(j) = i; \\ 1 - x_i & \text{if } g(j) = \bar{i}. \end{cases}$$

Example 9. To illustrate the function \tilde{g} , consider again the graphs G and H_1 from Figure 5, as well as the star-homomorphism $g_1 : V(H_1) \rightarrow V(G^\dagger)$ given in Example 8. Now let $x := (\frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{1}{5})^\top \in \mathbb{R}^{V(G)}$. Then $\tilde{g}_1(x) \in \mathbb{R}^{V(H_1)}$ where

$$\begin{array}{c|c|c|c|c|c|c|c|c|c} j \in V(H_1) & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ \hline [\tilde{g}_1(x)]_j & \frac{2}{5} & \frac{2}{5} & \frac{2}{5} & \frac{2}{5} & \frac{2}{5} & \frac{1}{5} & \frac{1}{5} & \frac{1}{5} & \frac{4}{5} \end{array}$$

Notice the point x given in Example 9 belongs to $\text{FRAC}(G)$, and that \tilde{g}_1 maps x to a point that belongs to $\text{FRAC}(H_1)$. This is not a coincidence, but rather a general property of \tilde{g} which follows readily from the definition of star-homomorphism:

Lemma 10. *Suppose H is star-homomorphic to G under g , and $x \in \mathbb{R}^{V(G)}$. If $x \in \text{FRAC}(G)$, then $\tilde{g}(x) \in \text{FRAC}(H)$.*

Proof. First, it is easy to see that $x \in [0, 1]^{V(G)}$ implies $\tilde{g}(x) \in [0, 1]^{V(H)}$. Now given edge $\{j_1, j_2\} \in E(H)$, $\{g(j_1), g(j_2)\}$ is either an edge in G or $\{i, \bar{i}\}$ for some $i \in V(G)$. In both cases, we see that $[\tilde{g}(x)]_{j_1} + [\tilde{g}(x)]_{j_2} \leq 1$. \square

In fact, the implication in Lemma 10 is preserved under applications of LS_+ .

Proposition 11. *Suppose H is star-homomorphic to G under g , and $x \in \mathbb{R}^{V(G)}$. If $x \in \text{LS}_+^p(G)$, then $\tilde{g}(x) \in \text{LS}_+^p(H)$.*

Proof. Suppose $x \in \text{LS}_+^p(G)$. We prove that $\tilde{g}(x) \in \text{LS}_+^p(H)$ by induction on p . The base case $p = 0$ reduces to Lemma 10. Next, suppose $p \geq 1$, and let $Y \in \widehat{\text{LS}}_+^p(G)$ be a certificate matrix. For convenience, we also extend the function \tilde{g} as follows: Given a real number $k \geq 0$, define $\tilde{g}_k : \mathbb{R}^{\{0\} \cup V(G)} \rightarrow \mathbb{R}^{\{0\} \cup V(H)}$ such that

$$[\tilde{g}_k(x)]_j := \begin{cases} x_0 & \text{if } j = 0; \\ x_i & \text{if } j \in V(H) \text{ and } g(j) = i; \\ k - x_i & \text{if } j \in V(H) \text{ and } g(j) = \bar{i}. \end{cases}$$

Notice that the function \tilde{g}_k satisfies

$$\tilde{g}_\lambda \left(\begin{bmatrix} \lambda \\ \lambda x \end{bmatrix} \right) = \begin{bmatrix} \lambda \\ \lambda \tilde{g}(x) \end{bmatrix}$$

for every $\lambda \geq 0$ and $x \in \mathbb{R}^{V(G)}$.

Since $Y \in \widehat{\text{LS}}_+^p(G)$, $Y e_i, Y(e_0 - e_i) \in \text{cone}(\text{LS}_+^{p-1}(G))$ for every $i \in V(G)$. Thus, applying the inductive hypothesis we have $\tilde{g}_{x_i}(Y e_i), \tilde{g}_{1-x_i}(Y(e_0 - e_i)) \in \text{cone}(\text{LS}_+^{p-1}(H))$.

Next, define the matrix $U \in \mathbb{R}^{\{0\} \cup V(G) \times \{0\} \cup V(H)}$ where

$$U e_j := \begin{cases} e_0 & \text{if } j = 0; \\ e_i & \text{if } j \in V(H) \text{ and } g(j) = i; \\ e_0 - e_i & \text{if } j \in V(H) \text{ and } g(j) = \bar{i}. \end{cases}$$

Then, given $z \in \mathbb{R}^{\{0\} \cup V(G)}$, $U^\top z = \tilde{g}_{z_0}(z)$.

Next, we claim that the matrix $Y' := U^\top Y U \in \widehat{\text{LS}}_+^p(H)$. First, since $Y = Y^\top$, $\text{diag}(Y) = Y e_0$, and $Y \succeq 0$, it is easy to see that the corresponding properties also hold for Y' . We next show that $Y' e_j, Y'(e_0 - e_j) \in \text{cone}(\text{LS}_+^{p-1}(H))$ for every $j \in V(H)$. First, if $g(j) = i \in V(G)$, then $Y' e_j = \tilde{g}_{x_i}(Y e_i)$, and

$$Y'(e_0 - e_j) = Y' e_0 - Y' e_j = \tilde{g}_1(Y e_0) - \tilde{g}_{x_i}(Y e_i) = \tilde{g}_{1-x_i}(Y(e_0 - e_i)).$$

To see the last equality, notice that

$$\begin{aligned} [\tilde{g}_1(Y e_0) - \tilde{g}_{x_i}(Y e_i)]_\ell &= [\tilde{g}_{1-x_i}(Y(e_0 - e_i))]_\ell \\ &= \begin{cases} 1 - x_i & \text{if } \ell = 0; \\ x_{i'} - Y[i', i] & \text{if } g(\ell) = i' \in V(G); \\ 1 - x_{i'} - x_i + Y[i', i] & \text{if } g(\ell) = \bar{i}' \text{ for some } i' \in V(G). \end{cases} \end{aligned}$$

Likewise, if $g(j) = \bar{i}$ for some $i \in V(G)$, then $Y' e_j = \tilde{g}_{1-x_i}(Y(e_0 - e_i))$ and $Y'(e_0 - e_j) = \tilde{g}_{x_i}(Y e_i)$. In all cases, it follows from the inductive hypothesis that $Y' e_j, Y'(e_0 - e_j) \in \text{cone}(\text{LS}_+^{p-1}(H))$.

Therefore, $Y' \in \widehat{\text{LS}}_+^p(H)$. Since $Y' e_0 = \tilde{g}_1(Y e_0) = \begin{bmatrix} 1 \\ \tilde{g}(x) \end{bmatrix}$, it follows that $\tilde{g}(x) \in \text{LS}_+^p(H)$. \square

Proposition 11 helps establish a framework for bounding the LS_+ -rank of a graph by that of another.

Lemma 12. *Let G and H be graphs where H is star-homomorphic to G under g . Assume that for every $x \in \mathbb{R}^{V(G)}$, the following holds:*

$$x \notin \text{STAB}(G) \Rightarrow \tilde{g}(x) \notin \text{STAB}(H).$$

Then $r_+(H) \geq r_+(G)$.

Proof. Suppose $r_+(G) = p \geq 1$ (the claim is trivial if $p = 0$). Then there exists $x \in \text{LS}_+^{p-1}(G) \setminus \text{STAB}(G)$. Then by the hypothesis and Proposition 11, $\tilde{g}(x) \in \text{LS}_+^{p-1}(H) \setminus \text{STAB}(H)$, showing that $r_+(H) \geq p$. \square

Finally, while our focus for this paper is the LS_+ operator, we remark that the framework of star-homomorphism can be extended to analyze relaxations generated by other lift-and-project operators. Again, let H be a graph that is star-homomorphic to G under g . Then notice that the function $\tilde{g} : \mathbb{R}^{V(G)} \rightarrow \mathbb{R}^{V(H)}$ can be expressed as a composition of the following four elementary operations:

- (1) Deleting a coordinate. E.g., $L : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ where

$$L(x_1, x_2, x_3, \dots, x_n) = (x_2, x_3, \dots, x_n).$$

- (2) Swapping two coordinates. E.g., $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ where

$$L(x_1, x_2, x_3, \dots, x_n) = (x_2, x_1, x_3, \dots, x_n).$$

- (3) Cloning a coordinate. E.g., $L : \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$ where

$$L(x_1, x_2, \dots, x_n) = (x_1, x_1, x_2, \dots, x_n).$$

- (4) Flipping a coordinate. E.g., $L : \mathbb{R}^n \rightarrow \mathbb{R}^n$ where

$$L(x_1, x_2, \dots, x_n) = (1 - x_1, x_2, \dots, x_n).$$

Now, let \mathcal{L} be a lift-and-project operator. If one can show that

$$(3) \quad x \in \mathcal{L}(P) \Rightarrow L(x) \in \mathcal{L}(L(P)),$$

for every function L that belongs to one of the four categories above (where $L(P)$ denotes $\{L(z) : z \in P\}$), then one can prove the analogous version of Proposition 11 for \mathcal{L} . For instance, the ideas from the proof of Proposition 11 show that (3) holds for $\mathcal{L} \in \{\text{LS}_+, \text{LS}, \text{LS}_0\}$ (where LS, LS_0 [LS91] are operators that generate polyhedral relaxations which are generally weaker than LS_+). Also, it has been shown [AT18, Proposition 1] that the Lasserre operator Las [Las01] commutes with all automorphisms of the unit hypercube, a property that is also shared by the Sherali–Adams operator SA [SA90] and one of its PSD variants SA_+ [AT16]. Thus, these operators satisfy (3) as well, and much of what we show for LS_+ in this section and the next section also applies to these operators.

4. THE (GENERALIZED) VERTEX-STRETCHING OPERATION

In this section, we introduce a graph operation that shows promise in producing relatively small graphs with high LS_+ -ranks, and study some of its properties. Given a graph G , vertex $v \in V(G)$, and non-empty sets $A_1, \dots, A_p \subset \Gamma_G(v)$ where $\bigcup_{\ell=1}^p A_\ell = \Gamma_G(v)$, we define the *stretching* of v in G by applying the following sequence of transformations to G :

- Replace v by $p + 1$ vertices: v_0, v_1, \dots, v_p ;
- For every $\ell \in [p]$, Join v_ℓ to v_0 , as well as to all vertices in A_ℓ .

We will also refer to the operation as *p-stretching* when we would like to specify p (which is necessarily at least 2). For example, Figure 6 shows the graph obtained from 2-stretching vertex 5 in K_5 (with $A_1 = \{2, 3, 4\}$ and $A_2 = \{1, 2, 3\}$). For another example, observe that in Figure 5, the graph H_1 can be obtained by 3-stretching vertex 6 in G .




FIGURE 6. Demonstrating the vertex-stretching operation

We remark that our vertex stretching operation is a generalization of the type (i) stretching operation described in [LT03] and later studied in [AEF14] (which further requires that $p = 2$ and $A_1 \cap A_2 = \emptyset$), as well as the k -stretching operation described in [BENT17] (which further requires that $p = 2$ and the vertices $A_1 \cap A_2$ induce a clique of size k in G , with $k = 0$ allowed). Also, when A_1, \dots, A_p are mutually disjoint and at most one of these p sets has size greater than 1, our vertex stretching specializes to an instance of type (ii) stretching from [LT03].

Given a graph G and an integer $p \geq 2$, we define $\mathcal{S}_p(G)$ to be the set of graphs that can be obtained from G by p -stretching one vertex. Notice that every graph $H \in \mathcal{S}_p(G)$ is star-homomorphic to G under g where

$$(4) \quad g(j) := \begin{cases} j & \text{if } j \in V(H \ominus v_0); \\ v & \text{if } j \in \{v_1, \dots, v_p\}; \\ \bar{v} & \text{if } j = v_0. \end{cases}$$

We also define $\mathcal{S}(G) := \bigcup_{p \geq 2} \mathcal{S}_p(G)$, and will show that $r_+(H) \geq r_+(G)$ for all $H \in \mathcal{S}(G)$ using Lemma 12. First, we need a tool that uses valid inequalities of $\text{STAB}(G)$ to generate potential valid inequalities for $\text{STAB}(H)$.

Lemma 13. *Let $H \in \mathcal{S}(G)$ be a graph obtained from G by p -stretching vertex $v \in V(G)$, and let $a^\top x \leq \beta$ be a valid inequality of $\text{STAB}(G)$ where $a \geq 0$. If $d \in \mathbb{R}_+^p$ satisfies $\sum_{\ell=1}^p d_\ell \geq a_v$ and*

$$(5) \quad \max \left\{ a^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup \bigcup_{\ell \in T} A_\ell \right\} \leq \beta - a_v + \sum_{\ell \notin T} d_\ell$$

for all $\emptyset \subset T \subset [p]$, then

$$(6) \quad \sum_{i \in V(H \ominus v_0)} a_i x_i + \sum_{\ell=1}^p d_\ell x_{v_\ell} + \left(\left(\sum_{\ell=1}^p d_\ell \right) - a_v \right) x_{v_0} \leq \beta - a_v + \sum_{\ell=1}^p d_\ell$$

is valid for $\text{STAB}(H)$.

Proof. Suppose $S \subseteq V(H)$ is an inclusion-wise maximal stable set in H . We define $S' \subseteq V(G)$ where

$$S' := \begin{cases} S \setminus \{v_0\} & \text{if } v_0 \in S; \\ (S \setminus \{v_1, \dots, v_p\}) \cup \{v\} & \text{if } \{v_1, \dots, v_p\} \subseteq S; \\ S \setminus \{v_1, \dots, v_p\} & \text{if } 1 \leq |\{v_1, \dots, v_p\} \cap S| \leq p-1. \end{cases}$$

In all cases, S' is a stable set in G , and $a^\top \chi_{S'} \leq \beta$ implies that χ_S satisfies (6). Note that the third case is when we require the assumption (5) with $T := \{v_1, \dots, v_p\} \cap S$. \square

Due to its similarity with the aforementioned vertex-stretching operations studied in [LT03], our vertex-stretching operation shares some similar structural properties, which we point out below.

Proposition 14. *Let $H \in \mathcal{S}(G)$ be a graph obtained from G by p -stretching vertex $v \in V(G)$. Then we have the following.*

(i) *Suppose $a^\top x \leq \beta$ is valid for $\text{STAB}(G)$ where $a \geq 0$. Then*

$$(7) \quad \sum_{i \in V(H \ominus v_0)} a_i x_i + \sum_{\ell=1}^p a_v x_{v_\ell} + (p-1)a_v x_{v_0} \leq \beta + (p-1)a_v$$

is valid for $\text{STAB}(H)$.

(ii) *Let g be as defined in (4). If $x \notin \text{STAB}(G)$, then $\tilde{g}(x) \notin \text{STAB}(H)$.*

(iii) *$r_+(H) \geq r_+(G)$.*

Proof. First, (i) follows readily from Lemma 13 with $d_\ell := a_v$ for every $\ell \in [p]$. Here, the condition (5) holds as the right hand side is at least β for all non-empty $T \subset [p]$.

For (ii), first suppose $x \notin \text{STAB}(G)$. If $x \notin [0, 1]^{V(G)}$, then $\tilde{g}(x) \notin [0, 1]^{V(H)}$ (where \tilde{g} is as defined in (2)) and the claim follows. Otherwise, there is a facet $a^\top x \leq \beta$ of $\text{STAB}(G)$ where $a \geq 0$ that is violated by x . Now notice that for every $j \in V(H)$,

$$[\tilde{g}(x)]_j = \begin{cases} x_j & \text{if } j \in V(H \ominus v_0); \\ x_v & \text{if } j \in \{v_1, \dots, v_p\}; \\ 1 - x_v & \text{if } j = v_0. \end{cases}$$

Then $\tilde{g}(x)$ violates (7), and thus does not belong to $\text{STAB}(H)$.

Finally, as we have shown that H is star-homomorphic to G under the function g as defined in (4), (iii) follows directly from Lemma 12. \square

We remark that Proposition 14 is a generalization of the corresponding results on types (i) and (ii) stretching from [LT03], and our proof uses many of the same ideas from similar arguments therein.

Next, we prove a result somewhat similar to Proposition 14(i) that derives some facets of the stable set polytope of the stretched graph.

Proposition 15. *Let $H \in \mathcal{S}(G)$ be a graph obtained from G by p -stretching vertex $v \in V(G)$, and suppose $a^\top x \leq \beta$ is a facet of $\text{STAB}(G)$ where $a \geq 0$. For every $\ell \in [p]$, define $A_\ell := \Gamma_H(v_\ell) \setminus \{v_0\}$ and*

$$(8) \quad d_\ell := a_v - \beta + \max \left\{ a^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup \bigcup_{j \in [p], j \neq \ell} A_j \right\}.$$

If the inequality (6) is valid for $\text{STAB}(H)$, then it is a facet of $\text{STAB}(H)$.

Proof. For convenience, let $n := |V(G)|$. Since $a^\top x \leq \beta$ is a facet of $\text{STAB}(G)$, there exist stable sets $S_1, \dots, S_n \subseteq V(G)$ whose incidence vectors are affinely independent and all satisfy $a^\top x \leq \beta$ with equality. Also, for every $\ell \in [p]$, let D_ℓ be a stable set that attains the maximum in the definition of d_ℓ in (8). We then define S'_1, \dots, S'_{n+p} as follows. For every $i \in [n]$,

$$S'_i := \begin{cases} S_i \cup \{v_0\} & \text{if } v \notin S_i; \\ (S_i \setminus \{v\}) \cup \{v_1, \dots, v_p\} & \text{if } v \in S_i. \end{cases}$$

We also define

$$S'_{n+i} := D_i \cup \{v_j : j \in [p], j \neq i\}$$

for all $i \in [p]$.

Observe that S'_1, \dots, S'_{n+p} must all be stable sets in H . Also, using the fact that incidence vectors of S_1, \dots, S_n are affinely independent and satisfy $a^\top x \leq \beta$ with equality, we see that the incidence vectors of S'_1, \dots, S'_{n+p} are affinely independent and all satisfy (6) with equality.

Thus, if we know that (6) is valid for $\text{STAB}(H)$, it must be a facet. \square

The special case of $p = 2$ in Proposition 15 is particularly noteworthy:

Corollary 16. *Let $H \in \mathcal{S}_2(G)$ be a graph obtained from G by 2-stretching vertex $v \in V(G)$, and suppose $a^\top x \leq \beta$ is a facet of $\text{STAB}(G)$ where $a \geq 0$. Define $A_1 := \Gamma_H(v_1) \setminus \{v_0\}$, $A_2 := \Gamma_H(v_2) \setminus \{v_0\}$, as well as the quantities*

$$\begin{aligned} d_1 &:= a_v - \beta + \max \left\{ a^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup A_2 \right\}, \\ d_2 &:= a_v - \beta + \max \left\{ a^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup A_1 \right\}. \end{aligned}$$

If $d_1 + d_2 \geq a_v$, then (6) is a facet of $\text{STAB}(H)$.

Proof. This is largely a specialization of Proposition 15 to the case $p = 2$. Notice that the additional assumption of (6) being valid is not necessary in this case because $\{1, 2\}$ has exactly two non-empty and proper subsets, and so the definition of d_1, d_2 herein are enough to guarantee that the assumption (5) is met. \square

Finally, we close this section by mentioning a “reverse” implication of Proposition 14.

Corollary 17. *Let H be a graph, and $v_0 \in V(H)$. If $\Gamma_H(v_0)$ is a stable set, then $r_+(G) \leq r_+(H)$, where G is the graph obtained from H by contracting the set of vertices $\Gamma_H[v_0]$.*

Proof. Let $v_1, \dots, v_p \in V(H)$ be the neighbours of v_0 , and define $A_i := \Gamma_H(v_i) \setminus \{v_0\}$ for every $i \in [p]$. Then there are two cases: First, if there exists $i \in [p]$ where $A_i = \bigcup_{j \in [p]} A_j$, then G is isomorphic to the subgraph of H induced by $(V(H) \setminus \Gamma_H[v_0]) \cup \{v_i\}$, and thus $r_+(G) \leq r_+(H)$.

Next, if there does not exist $i \in [p]$ where $A_i = \bigcup_{j \in [p]} A_j$, then $H \in \mathcal{S}_p(G)$, and it follows that $r_+(G) \leq r_+(H)$. \square




FIGURE 7. An example in which contracting the closed neighborhood of a vertex increases the graph’s LS_+ -rank

Example 18. In general, it is possible that contracting the closed neighborhood of a vertex results in an increase in the graph’s LS_+ -rank. For example, notice that the graph H in Figure 7 is the union of two LS_+ -rank-1 graphs whose intersection is the cut clique $\{7, 8\}$, and thus it follows from Proposition 6 that $r_+(H) = 1$. However, contracting $\Gamma_H[6]$ in H results in G , which is isomorphic to $G_{2,1}$, and so $r_+(G) = 2$.

5. LS_+ -MINIMAL GRAPHS VIA 2-STRETCHING CLIQUES

In this section, we are interested in studying graphs with the fewest number of vertices with a given LS_+ -rank. Given $\ell \in \mathbb{N}$, define $n_+(\ell)$ to be the minimum number of vertices on which there exists a graph G with $r_+(G) = \ell$. It follows immediately from Theorem 1 that $n_+(\ell) \geq 3\ell$ for every $\ell \in \mathbb{N}$. On the other hand, Theorem 2 implies that $n_+(\ell) \leq 16\ell$. Thus, we know that $n_+(\ell) = \Theta(\ell)$ asymptotically.

Recall that a graph G is ℓ -minimal if $r_+(G) = \ell$ and $|V(G)| = 3\ell$. The following result establishes a close connection between ℓ -minimal graphs and 2-stretching vertices.

Theorem 19. *Let H be an ℓ -minimal graph where $\ell \geq 2$. Then there exists a graph G where $H \in \mathcal{S}_2(G)$.*

Proof. First, since $r_+(H) = \ell$, there exists a vertex v_0 where $r_+(H \ominus v_0) = \ell - 1$. This implies that $|V(H \ominus v_0)| \geq 3\ell - 3$, and thus $\deg(v_0) \leq 2$. If $\deg(v_0) = 1$, H would contain a cut vertex, and Proposition 6 implies there would be a proper subgraph of H with the same LS_+ -rank as H . Thus, we obtain that $\deg(v_0) = 2$. Let v_1, v_2 denote the two neighbours of v_0 . If $\{v_1, v_2\} \in E(H)$, then the edge would form a cut clique in H , and Proposition 6 again implies that there would be a proper subgraph of H with the same LS_+ -rank as H . Thus, $\{v_1, v_2\}$ must be a stable set in H .

Next, define G to be the graph obtained from H by contracting $\{v_0, v_1, v_2\}$, and label the new vertex v . We claim that H can be obtained from G by 2-stretching $v \in V(G)$ with $A_1 := \Gamma_H(v_1) \setminus \{v_0\}$ and $A_2 := \Gamma_H(v_2) \setminus \{v_0\}$. To prove this, it suffices to show that A_1, A_2 are both proper subsets of $A_1 \cup A_2$. Equivalently, we need to show that $\Gamma_H(v_1) \setminus \Gamma_H(v_2)$ and $\Gamma_H(v_2) \setminus \Gamma_H(v_1)$ are non-empty.

Let $a^\top x \leq \beta$ be a facet of $\text{STAB}(H)$ of LS_+ -rank ℓ . Then there must be stable sets $S_1, \dots, S_{3\ell} \subseteq V(H)$ whose incidence vectors are affinely independent and all satisfy $a^\top x \leq \beta$ with equality. Also, since H is ℓ -minimal, a must have full support. Therefore, S_i must be inclusion-wise maximal for all $i \in [3\ell]$, and hence belongs to one of the following cases:

- (1) $S_i \cap \{v_0, v_1, v_2\} = \{v_0\}$;
- (2) $S_i \cap \{v_0, v_1, v_2\} = \{v_1\}$;
- (3) $S_i \cap \{v_0, v_1, v_2\} = \{v_2\}$;
- (4) $S_i \cap \{v_0, v_1, v_2\} = \{v_1, v_2\}$.

Since $\chi_{S_1}, \dots, \chi_{S_{3\ell}}$ are affinely independent, one of these stable sets contains v_0 and belongs to Case (1), so assume without loss of generality that $v_0 \in S_1$. Now consider the matrix A formed by the row vectors $(\chi_{S_2} - \chi_{S_1})^\top, (\chi_{S_3} - \chi_{S_1})^\top, \dots, (\chi_{S_{3\ell}} - \chi_{S_1})^\top$. Since $S_1, \dots, S_{3\ell}$ are affinely independent, A must have linearly independent rows. This means that, if we focus on the submatrix A' of A which consists of just the three columns corresponding to v_0, v_1 , and v_2 , A' must have rank 3. This implies that there must exist at least one S_i belonging to each of Cases (2), (3), and (4).

Now consider a stable set S_i that belongs to Case (2). Since S_i is inclusion-wise maximal, it must contain a vertex that is adjacent to v_2 and not v_1 , and so we have found a vertex that belongs to $\Gamma_H(v_2) \setminus \Gamma_H(v_1)$. The same argument applied to an S_i from Case (3) gives a vertex that belongs to $\Gamma_H(v_1) \setminus \Gamma_H(v_2)$. This finishes the proof. \square

Thus, for the remainder of this section, we will focus on 2-stretching vertices, and study when that helps (and does not help) in generating ℓ -minimal graphs. Since we will be studying graphs obtained from applying a sequence of 2-stretching operations, we recursively define

$$\mathcal{S}_2^k(G) := \bigcup_{G' \in \mathcal{S}_2^{k-1}(G)} \mathcal{S}_2(G')$$

for every graph G and integer $k \geq 1$. That is, $\mathcal{S}_2^k(G)$ is the set of graphs that can be obtained from G by a sequence of k 2-stretching operations. We also let $\mathcal{S}_2^0(G) := \{G\}$. The following is a basic property of the graphs in $\mathcal{S}_2^k(G)$.

Lemma 20. *Let G be a graph, and let $H \in \mathcal{S}_2^k(G)$. Then $\alpha(H) = \alpha(G) + k$.*

Proof. Let $H \in \mathcal{S}_2(G)$ be a graph obtained from G by 2-stretching vertex $v \in V(G)$. To prove our claim, it suffices to show that $\alpha(H) = \alpha(G) + 1$. Consider a set of vertices $S \subseteq V(G)$. If $v \in S$, then S is a stable set in G if and only if $(S \setminus \{v\}) \cup \{v_1, v_2\}$ is a stable set in H . If $v \notin S$, then S is a stable set in G if and only if $S \cup \{v_0\}$ is a stable set in H . Thus, we see that $\alpha(H) = \alpha(G) + 1$. \square

Recall the graphs $G_{2,1}$, $G_{2,2}$, and $G_{3,1}$. In Figure 1, we labelled the vertices of these graphs to highlight the fact that all three graphs can be obtained from applying a number of 2-stretching operations to a complete graph. In fact, every known ℓ -minimal graph to date — the 3-cycle and the three graphs in Figure 1 — belongs to $\mathcal{S}^{\ell-1}(K_{\ell+2})$. Thus, for the remainder of this section, we focus on graphs obtained from 2-stretching vertices of a complete graph, and prove some results about the LS_+ -ranks of these graphs.

Generally, to prove that a given graph G has LS_+ -rank at least p , one needs to find a point $x \in \text{LS}_+^{p-1}(G) \setminus \text{STAB}(G)$. In particular, showing that $x \in \text{LS}_+^{p-1}(G)$ often requires finding a corresponding certificate matrix Y and showing that the matrix is positive semidefinite (among satisfying other properties imposed by LS_+). Thus, we first provide a framework for easily and reliably verifying the positive semidefiniteness of specific matrices. Given a symmetric matrix $Y \in \mathbb{R}^{n \times n}$, we say that $U, V \in \mathbb{Z}^{n \times n}$ is a *UV-certificate* of Y if

- (i) $kY = U^\top U + V$ for some $k \in \mathbb{N}$, and,
- (ii) V is diagonally dominant (i.e., $V_{ii} \geq \sum_{j \neq i} |V_{ij}|$ for every $i \in [n]$).

Observe that the existence of a *UV-certificate* implies that $Y \succeq 0$ (these certificates are sum-of-squares certificates, and every rational matrix $Y \in \mathbb{S}_+^n$ admits such certificates). Conversely, since every entry in U and V is an integer, verifying (i) and (ii) for a given *UV-certificate* only involves elementary numerical operations on whole numbers.

Next, we show that if we 2-stretch a vertex in a complete graph, the result is always a graph with LS_+ -rank 2.

Proposition 21. *Let $n \geq 4$. Then $r_+(H) = 2$ for all $H \in \mathcal{S}_2(K_n)$.*

Proof. Let $H \in \mathcal{S}_2(K_n)$, and assume without loss of generality that H is obtained from K_n by 2-stretching vertex n . Also, let G_n be the graph obtained from 2-stretching vertex n in K_n with $A_1 := [n-1] \setminus \{1\}$ and $A_2 := [n-2]$. (For example, G_4 is the graph $G_{2,2}$ from Figure 1 and G_5 is shown in Figure 6.) Now since the sets A_1, A_2 used in stretching vertex n to obtain H must satisfy $A_1, A_2 \subset [n-1]$ and $A_1 \cup A_2 = [n-1]$, we see that H must be isomorphic to a subgraph of G_n .

Next, we show that $\alpha_{\text{LS}_+}(G_4) > 2$. Consider the certificate matrix

$$Y := \begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4_1 & 4_0 & 4_2 \end{matrix} \\ \begin{bmatrix} 200 & 78 & 12 & 78 & 78 & 78 & 78 \\ 78 & 78 & 0 & 0 & 39 & 39 & 0 \\ 12 & 0 & 12 & 0 & 0 & 12 & 0 \\ 78 & 0 & 0 & 78 & 0 & 39 & 39 \\ 78 & 39 & 0 & 0 & 78 & 0 & 39 \\ 78 & 39 & 12 & 39 & 0 & 78 & 0 \\ 78 & 0 & 0 & 39 & 39 & 0 & 78 \end{bmatrix} \end{matrix}$$

Note that the columns of Y are labelled by the vertices in G_4 they correspond to (the rows of Y follow the same order of indexing). Observe that $Y \succeq 0$ — a *UV-certificate* for Y is

$$U := \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -73 & -26 & -141 & -26 & 60 & 132 & 61 \\ 0 & 124 & 0 & -124 & -200 & 0 & 200 \\ 27 & -181 & 247 & -181 & 51 & 159 & 51 \\ 0 & -527 & 0 & 527 & -326 & 0 & 326 \\ 1 & -166 & -73 & -166 & 449 & -556 & 449 \\ 1224 & 482 & 60 & 482 & 482 & 485 & 482 \end{bmatrix}, \quad V := \begin{bmatrix} 2765 & 917 & 91 & 917 & 316 & -11 & 389 \\ 917 & 1308 & 3 & -212 & -136 & 10 & 29 \\ 91 & 3 & 601 & 3 & -280 & 71 & -139 \\ 917 & -212 & 3 & 1308 & 3 & 10 & -110 \\ 316 & -136 & -280 & 3 & 1328 & -155 & -45 \\ -11 & 10 & 71 & 10 & -155 & 664 & -287 \\ 389 & 29 & -139 & -110 & -45 & -287 & 1207 \end{bmatrix},$$

which gives $7535Y = U^\top U + V$. One can also check that $Ye_i, Y(e_0 - e_i) \in \text{FRAC}(G_4)$ for every $i \in V(G_4)$. This shows that $\bar{x} := \frac{1}{200}(78, 12, 78, 78, 78, 78)^\top \in \text{LS}_+(G_4)$. Now since

$\bar{e}^\top \bar{x} = 2.01 > 2 = \alpha(G_4)$, we see that $r_+(G_4) \geq 2$. Since G_n contains G_4 as an induced subgraph for all $n \geq 4$, we conclude that $\alpha_{\text{LS}_+}(G_n) > 2$. Then Lemma 7(ii) implies that $\alpha_{\text{LS}_+}(H) > 2$. Since $\alpha(H) = 2$, it follows that $r_+(H) \geq 2$.

Finally, notice that $H - n_0$ must be a perfect graph, so $r_+(H - n_0) \leq 1$ and consequently $r_+(H) \leq 2$. Thus, we conclude that $r_+(H) = 2$. \square

We remark that, to obtain the numerical certificates in the proof above, we first found a matrix Y_0 by optimizing over $\text{LS}_+(G_4)$ using CVX, a package for specifying and solving convex programs [GB14, GB08], as well as the SDP solver SeDuMi [Stu99]. Then we let $V_0 := \lambda_0 I$, where λ_0 is the least eigenvalue of Y_0 , and solved for a matrix U_0 where $U_0^\top U_0 + V_0 = Y_0$. From there, we “discretize” U_0, V_0 in a somewhat trial-and-error manner until we obtain a suitable UV -certificate. The numerical certificates for the 4-minimal graph $G_{4,1}$ presented in the proof of Theorem 26 and Lemmas 39, 40, and 41 are found in a similar manner. It would be interesting to look into the prospect of generating these certificates via a more systematic and efficient approach.

Next, we also remark that in the proof for $r_+(G_{2,2}) \geq 2$ in [EMN06], the following certificate matrix was given:

$$\frac{1}{2688} \begin{bmatrix} & 4_2 & 1 & 3 & 4_1 & 4_0 & 2 \\ 2688 & 769 & 769 & 769 & 769 & 769 & 1538 \\ 769 & 769 & 0 & 336 & 413 \frac{7}{13} & 0 & 0 \\ 769 & 0 & 769 & 0 & 336 & 384 & 0 \\ 769 & 336 & 0 & 769 & 0 & 384 & 0 \\ 769 & 413 \frac{7}{13} & 336 & 0 & 769 & 0 & 896 \\ 769 & 0 & 384 & 384 & 0 & 769 & 0 \\ 1538 & 0 & 0 & 0 & 896 & 0 & 1538 \end{bmatrix}$$

However, the certificate is incorrect: $Y[2, 4_1] = \frac{896}{2688} = \frac{1}{3} > Y[0, 4_1]$, and thus violates $Ye_{4_1} \in \text{cone}(\text{FRAC}(G_{2,2}))$. In fact, since the vector $\frac{1}{2688}(769, 769, 769, 769, 769, 1538)^\top$ contains only one entry greater than $\frac{1}{3}$, any certificate matrix for this vector cannot contain an off-diagonal entry of $\frac{1}{3}$ (which would have to appear in at least 2 columns in the certificate). Still, the claim that $r_+(G_{2,2}) = 2$ is correct, as shown in the proof of Proposition 21.

Next, while all graphs in $\mathcal{S}_2(K_n)$ have LS_+ -rank 2, we show that not all graphs in $\mathcal{S}_2^2(K_n)$ have LS_+ -rank 3. Given a graph G , we say that a path in G is *sparse* if at most one of the vertices in the path has degree greater than 2 in G . For example, in Figure 8, the graph on the left contains a sparse path $4_0, 4_1, 5_1, 5_0, 5_2$ of length 4, while the graph on the right also contains a sparse path $4_0, 4_2, 3, 5_2, 5_0$ of length 4. Then we have the following.




FIGURE 8. Two graphs in $\mathcal{S}_2^2(K_5)$ with sparse paths

Proposition 22. *Let $\ell \geq 2$. If $G \in \mathcal{S}_2^{\ell-1}(K_{\ell+2})$ contains a sparse path of length at least 3, then G is not ℓ -minimal.*

Proof. First, observe that each 2-stretching operation adds 2 vertices to a given graph, and so every $G \in \mathcal{S}_2^{\ell-1}(K_{\ell+2})$ has $\ell + 2 + 2(\ell - 1) = 3\ell$ vertices. We show that if G contains a sparse path of length at least 3, then $r_+(G) < \ell$.

Let U be a set of vertices that induces a sparse path in G where $|U| \geq 4$. Then there is a vertex $u_1 \in U$ where $\deg(v) = 2$ for all $v \in U \setminus \{u_1\}$. Now notice that in the graph $G - u_1$, there must exist $u_2 \in U \setminus \{u_1\}$ where $\deg(u_2) = 1$. Thus, Proposition 6 implies that $G - u_1$ and $G - \{u_1, u_2\}$ must have the same LS_+ -rank.

Again, since U induce a sparse path in G , there must be a vertex $u_3 \in U \setminus \{u_1, u_2\}$ such that $\deg(u_3) = 1$ in $G - \{u_1, u_2\}$, and so $G - \{u_1, u_2\}$ and $G - \{u_1, u_2, u_3\}$ must have the same LS_+ -rank. Iterating this argument until we remove all vertices of U from G , we see that $r_+(G - u_1) = r_+(G - U)$.

Since $|U| \geq 4$, $G - U$ has $3\ell - |U| < 3(\ell - 1)$ vertices, and so $r_+(G - u_1) = r_+(G - U) \leq \ell - 2$. As a result, $r_+(G) \leq \ell - 1$, and thus G is not ℓ -minimal. \square

Thus, both graphs in Figure 8 have LS_+ -rank at most 2. (In fact, they have rank 2 as they both contain $G_{2,1}$ as an induced subgraph.) Next, we show that if we 2-stretch a vertex in K_n , and then 2-stretch one of the three new vertices in the stretched graph, the resulting graph cannot have LS_+ -rank 3.

Proposition 23. *Let $n \geq 4$. Suppose $G_1 \in \mathcal{S}_2(K_n)$ is obtained by stretching vertex n in K_n , and $G_2 \in \mathcal{S}_2^2(K_n)$ is obtained by stretching vertex n_0, n_1 , or n_2 in G_1 . Then $r_+(G_2) = 2$.*

Proof. First, if G_2 is obtained from G_1 by stretching n_0 , then n_{00}, n_{01}, n_{02} all have degree 2. Notice that $G_2 - n_{00}$ must be a perfect graph, and so $r_+(G_2 - n_{00}) \leq 1$, which implies $r_+(G_2) \leq 2$ in this case.

Otherwise, assume without loss of generality that G_2 is obtained from G_1 by stretching n_1 , and that $\{n_{12}, n_0\} \in E(G_2)$. (Note that $\{n_{11}, n_0\}$ may or may not be an edge.) Now notice that $G_2 - n_{12}$ is a perfect graph, and thus, $r_+(G_2) \leq 2$ in this case as well. Finally, since $r_+(G_1) = 2$ (from Proposition 21) and $G_2 \in \mathcal{S}(G_1)$, Proposition 14(iii) implies that $r_+(G_2) \geq 2$. \square

Thus, to obtain a graph with LS_+ -rank 3 in $\mathcal{S}_2^2(K_n)$, it is necessary that we stretch two of the original vertices of K_n . (That is not sufficient though, as shown for the graphs in Figure 8.)

Next, observe that if G is an ℓ -minimal graph, then $\text{STAB}(G)$ must have a facet with LS_+ -rank ℓ , and it is necessary that such a facet has full support. (Otherwise it follows from Lemma 4 that G has a proper subgraph with the same LS_+ -rank, which implies the existence of a graph on fewer than 3ℓ vertices with LS_+ -rank ℓ .) We provide more circumstantial evidence that 2-stretching a number of original vertices of a complete graph is a promising approach for generating ℓ -minimal graphs by showing that the stable set polytope of many of these graphs have a full-support facet. Let $H \in \mathcal{S}_2^k(K_n)$ be a graph that is obtained from 2-stretching the vertices $D \subseteq V(K_n)$ (and thus $|D| = k$). For every $v \in V(K_n)$, we define the vertices *associated* with v to be v_0, v_1 , and v_2 if $v \in D$, and just the unstretched vertex v if $v \notin D$. Notice that every vertex in H is associated with exactly one $v \in V(K_n)$. Then we have the following.

Proposition 24. *Let k, n be integers where $n \geq 3$ and $n \geq k \geq 0$. Suppose $H \in \mathcal{S}_2^k(K_n)$ is obtained from K_n by 2-stretching the vertices $D \subseteq V(K_n)$. Furthermore, suppose H satisfies the following:*

- (*) *For every $v \in D$ and for every $j \in \{1, 2\}$, there exists $u \in V(K_n) \setminus \{v\}$ where v_j is not adjacent with any vertex associated with u .*

Then $\sum_{i \in V(H)} x_i \leq k + 1$ is a facet of $\text{STAB}(H)$.

Proof. We prove our claim by induction on k . When $k = 0$, $H = K_n$, and the claim obviously holds. Next, assume $1 \leq k \leq n$. Since $H \in \mathcal{S}_2^k(K_n)$, there exists $G \in \mathcal{S}_2^{k-1}(K_n)$ where $H \in \mathcal{S}_2(G)$. (So there exists $v \in D$ where H is obtained from G by stretching v .)

Since H satisfies $(*)$ by assumption, so must G , and so it follows from the inductive hypothesis that $\sum_{i \in V(G)} x_i \leq k$ is a facet of $\text{STAB}(G)$. To prove our claim, we make use of Proposition 15 and show that $d_1 = d_2 = 1$. To do so, let $A_1 := \Gamma_H(v_1) \setminus \{v_0\}$, $A_2 := \Gamma_H(v_2) \setminus \{v_0\}$,

$$c_1 := \max \left\{ \bar{e}^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup A_2 \right\},$$

$$c_2 := \max \left\{ \bar{e}^\top x : x \in \text{STAB}(G), x_i = 0 \ \forall i \in \{v\} \cup A_1 \right\}.$$

Then it suffices to prove that $c_1 = c_2 = k$, which would then imply that $d_1 = d_2 = 1$.

First, it is obvious that $c_1, c_2 \leq k$ since $\alpha(G) = k$. Next, consider $\Gamma_H(v_1)$. By the assumption that H satisfies $(*)$, at least one of the following must hold:

- There exists $u \in V(K_n)$ where $u \notin D$ (so $u \in V(H)$) and $u \notin \Gamma_H(v_1)$. Then

$$S := \{u\} \cup \{w_0 : w \in D \setminus \{v\}\}$$

is a stable set that gives $c_1 = k$.

- There exists $u \in V(K_n)$, $u \neq v$ where $u \in D$ (so $u_0, u_1, u_2 \in V(H)$) and $u_0, u_1, u_2 \notin \Gamma_H(v_1)$. Then

$$S := \{u_1, u_2\} \cup \{w_0 : w \in D \setminus \{u, v\}\}$$

is a stable set that gives $c_1 = k$.

The same argument shows that $c_2 = k$, and this finishes the proof. \square

We remark that the assumption of stretching only the original vertices of K_n in Proposition 24 is necessary, as shown in the following example.

Example 25. Recall the graph $G_{2,2}$, introduced in Figure 1 and reproduced below in Figure 9 (left). Observe that $G_{2,2} \in \mathcal{S}_2(K_4)$, and that $\bar{e}^\top x \leq 2$ is a facet of $\text{STAB}(G_{2,2})$. Now, we 2-stretch the vertex $4_2 \in V(G_{2,2})$ to obtain $H \in \mathcal{S}_2^2(K_4)$ as shown in Figure 9 (right). Observe that the subgraph of H induced by vertices $1, 2, 3, 4_1, 4_0, 4_{21}$ is isomorphic to $G_{2,1}$ from Figure 1. Thus,

$$(9) \quad x_1 + x_2 + x_3 + x_{4_1} + x_{4_0} + x_{4_{21}} \leq 2$$

is valid for $\text{STAB}(H)$. (In fact, one can show that it is a facet of $\text{STAB}(G)$ using Proposition 15.) This implies that $\sum_{i \in V(H)} x_i \leq 3$, which is the sum of (9) and the edge inequality $x_{4_{20}} + x_{4_{22}} \leq 1$, is not a facet of $\text{STAB}(H)$.

6. EXISTENCE OF 4-MINIMAL GRAPHS

Recall the graph $G_{4,1}$ (Figure 3), which was introduced in Section 1. We show in this section that $r_+(G_{4,1}) = 4$, providing what we believe to be the first known example of a 4-minimal graph (and the first advance in this direction since 2006 [EMN06]). Notice that $G_{4,1}$ dovetails nicely with the structural results on ℓ -minimal graphs developed in Sections 3 to 5 — in particular, observe from its drawing in Figure 3 that $G_{4,1} \in \mathcal{S}_2^3(K_6)$, and is obtained from stretching three of the original vertices in K_6 . On the other hand, while these structural results were helpful in our discovery of $G_{4,1}$ and putting the graph into perspective, the proof we present for $r_+(G_{4,1}) = 4$ is largely based on numerical certificates and the symmetries of $G_{4,1}$.




FIGURE 9. A graph in $H \in \mathcal{S}_2^2(K_4)$ (right) where $\bar{e}^\top x \leq \alpha(H)$ is not a facet of $\text{STAB}(H)$ (see Example 25)




FIGURE 10. An alternative drawing of $G_{4,1}$ to highlight its automorphisms

To that end, we point out two important automorphisms of $G_{4,1}$ that will be useful in simplifying our analysis of its LS_+ -rank. Consider the alternative drawing of $G_{4,1}$ in Figure 10, and define the functions $f_1, f_2 : V(G_{4,1}) \rightarrow V(G_{4,1})$ as follows:

i	1	2	3	4 ₁	4 ₀	4 ₂	5 ₁	5 ₀	5 ₂	6 ₁	6 ₀	6 ₂
$f_1(i)$	2	3	1	5 ₁	5 ₀	5 ₂	6 ₁	6 ₀	6 ₂	4 ₁	4 ₀	4 ₂
$f_2(i)$	1	3	2	5 ₂	5 ₀	5 ₁	4 ₂	4 ₀	4 ₁	6 ₂	6 ₀	6 ₁

Visually, f_1 corresponds to rotating the graph $G_{4,1}$ in Figure 10 counterclockwise by $\frac{2\pi}{3}$, and f_2 corresponds to reflecting the figure along the centre vertical line. Now we are ready to prove the main result of this section.

Theorem 26. *The LS_+ -rank of $G_{4,1}$ is 4.*

Proof. For convenience, let $G := G_{4,1}$ throughout this proof. Since G has 12 vertices, by Theorem 1 it suffices to show that $r_+(G) \geq 4$. Consider the matrix Y_0 defined as follows:

$$Y_0 := \begin{bmatrix} 100000 & 25340 & 25340 & 25340 & 16500 & 75020 & 16500 & 16500 & 75020 & 16500 & 16500 & 75020 & 16500 \\ 25340 & 25340 & 0 & 0 & 0 & 17502 & 7838 & 7838 & 17502 & 0 & 0 & 25340 & 0 \\ 25340 & 0 & 25340 & 0 & 0 & 25340 & 0 & 0 & 17502 & 7838 & 7838 & 17502 & 0 \\ 25340 & 0 & 0 & 25340 & 7838 & 17502 & 0 & 0 & 25340 & 0 & 0 & 17502 & 7838 \\ 16500 & 0 & 0 & 7838 & 16500 & 0 & 8073 & 589 & 15911 & 0 & 589 & 15419 & 1081 \\ 75020 & 17502 & 25340 & 17502 & 0 & 75020 & 0 & 15419 & 51150 & 15911 & 15911 & 51150 & 15419 \\ 16500 & 7838 & 0 & 0 & 8073 & 0 & 16500 & 1081 & 15419 & 589 & 0 & 15911 & 589 \\ 16500 & 7838 & 0 & 0 & 589 & 15419 & 1081 & 16500 & 0 & 8073 & 589 & 15911 & 0 \\ 75020 & 17502 & 17502 & 25340 & 15911 & 51150 & 15419 & 0 & 75020 & 0 & 15419 & 51150 & 15911 \\ 16500 & 0 & 7838 & 0 & 0 & 15911 & 589 & 8073 & 0 & 16500 & 1081 & 15419 & 589 \\ 16500 & 0 & 7838 & 0 & 589 & 15911 & 0 & 589 & 15419 & 1081 & 16500 & 0 & 8073 \\ 75020 & 25340 & 17502 & 17502 & 15419 & 51150 & 15911 & 15911 & 51150 & 15419 & 0 & 75020 & 0 \\ 16500 & 0 & 0 & 7838 & 1081 & 15419 & 589 & 0 & 15911 & 589 & 8073 & 0 & 16500 \end{bmatrix}.$$

Again, the columns of Y_0 are labelled by the vertices in G they correspond to, with the rows of Y_0 following the same order of indexing.

We prove our claim by showing that $Y_0 \in \widehat{LS}_+^3(G)$. First, one can check that $Y_0 \succeq 0$ (a UV -certificate is provided in Table 1). Moreover, observe that for all $i, j \in V(G)$,

$$Y_0[i, j] = Y_0[f_1(i), f_1(j)] = Y_0[f_2(i), f_2(j)],$$

and thus the entries of Y_0 exhibit the same symmetries of the graph that are exposed by the automorphisms f_1 and f_2 . Hence, to show that $Y_0 \in \widehat{LS}_+^3(G)$, it suffices to verify the conditions $Y_0 e_i, Y_0(e_0 - e_i) \in \text{cone}(LS_+^2(G))$ for $i \in \{1, 4_1, 6_0\}$, since for every other vertex j there is an automorphism of G that would map j to one of these three vertices.

Next, notice that

$$\begin{aligned} Y_0 e_1 &\leq 17502 \begin{bmatrix} 1 \\ \chi_{\{1, 4_0, 5_0, 6_0\}} \end{bmatrix} + 7838 \begin{bmatrix} 1 \\ \chi_{\{1, 4_2, 5_1, 6_0\}} \end{bmatrix}, \\ Y_0 e_{4_1} &\leq 7838 \begin{bmatrix} 1 \\ \chi_{\{3, 4_1, 5_0, 6_0\}} \end{bmatrix} + 589 \begin{bmatrix} 1 \\ \chi_{\{4_1, 4_2, 5_1, 6_0\}} \end{bmatrix} + 6992 \begin{bmatrix} 1 \\ \chi_{\{4_1, 4_2, 5_0, 6_0\}} \end{bmatrix} \\ &\quad + 492 \begin{bmatrix} 1 \\ \chi_{\{4_1, 4_2, 5_0, 6_2\}} \end{bmatrix} + 589 \begin{bmatrix} 1 \\ \chi_{\{4_1, 5_0, 6_1, 6_2\}} \end{bmatrix}, \\ Y_0(e_0 - e_{6_0}) &\leq 7366 \begin{bmatrix} 1 \\ \chi_{\{2, 4_0, 5_0, 6_1\}} \end{bmatrix} + 476 \begin{bmatrix} 1 \\ \chi_{\{2, 4_0, 5_2, 6_1\}} \end{bmatrix} + 476 \begin{bmatrix} 1 \\ \chi_{\{3, 4_1, 5_0, 6_2\}} \end{bmatrix} \\ &\quad + 7366 \begin{bmatrix} 1 \\ \chi_{\{3, 4_0, 5_0, 6_2\}} \end{bmatrix} + 605 \begin{bmatrix} 1 \\ \chi_{\{4_1, 4_2, 5_0, 6_2\}} \end{bmatrix} + 605 \begin{bmatrix} 1 \\ \chi_{\{4_0, 5_1, 5_2, 6_1\}} \end{bmatrix} \\ &\quad + 8058 \begin{bmatrix} 1 \\ \chi_{\{4_0, 5_0, 6_1, 6_2\}} \end{bmatrix} + 28 \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \end{aligned}$$

Notice that all incidence vectors above correspond to stable sets in G . Since $\text{cone}(\text{STAB}(G))$ is a lower-comprehensive convex cone, we obtain that $Y_0 e_1, Y_0 e_{4_1}, Y_0(e_0 - e_{6_0}) \in \text{cone}(\text{STAB}(G)) \subseteq \text{cone}(LS_+^2(G))$. The details for $Y_0(e_0 - e_1), Y_0 e_{6_0}, Y_0(e_0 - e_{4_1}) \in \text{cone}(LS_+^2(G))$ are provided, respectively, in the proofs of Lemmas 39, 40, and 41 in Appendix A.

Finally, let \bar{x} be the vector such that $Y_0 e_0 = 100000 \begin{bmatrix} 1 \\ \bar{x} \end{bmatrix}$. Since $Y_0 \in \widehat{LS}_+^3(G)$, we have $\bar{x} \in LS_+^3(G)$. Thus, we see that

$$\alpha_{LS_+^3}(G) \geq \bar{e}^\top \bar{x} = 4.0008 > 4 = \alpha(G).$$

This shows $\bar{x} \notin \text{STAB}(G)$ and thus $r_+(G) \geq 4$. \square

Notice that $G_{4,1}$ contains 24 edges. Also, notice that $G_{4,1} \in \mathcal{S}_2^3(K_6)$ and satisfies $(*)$ from Proposition 24. Consequently, every graph $G \in \mathcal{S}_2^3(K_6)$ which is a subgraph of $G_{4,1}$ also satisfies $(*)$, and so Proposition 24 assures that the inequality $\bar{e}^\top x \leq 4$ is a facet of $\text{STAB}(G)$. Thus, by Lemma 7(ii) it follows that every graph in $\mathcal{S}_2^3(K_6)$ which is a subgraph of $G_{4,1}$ (which can have as few as 21 edges) also has LS_+ -rank 4, giving more examples of 4-minimal graphs. The six non-isomorphic proper subgraphs of $G_{4,1}$ that belong to $\mathcal{S}_2^3(K_6)$ are listed in Figure 11.




FIGURE 11. The six non-isomorphic proper subgraphs of $G_{4,1}$ that belong to $\mathcal{S}_2^3(K_6)$ (and thus are 4-minimal)

Moreover, the fact that $G_{4,1}$ is 4-minimal also provides some new examples of 3-minimal graphs.

Corollary 27. *Let $G_{3,2} := G_{4,1} \ominus 6_0$. Then $G_{3,2}$ is a 3-minimal graph.*

Proof. Since $G_{4,1}$ is 4-minimal, there exists vertex $i \in V(G_{4,1})$ where $G \ominus i$ has LS_+ -rank 3 and $\deg(i) = 2$. Thus, $i \in \{4_0, 5_0, 6_0\}$. Now observe that $G_{4,1} \ominus 4_0, G_{4,1} \ominus 5_0, G_{4,1} \ominus 6_0$ are all isomorphic to each other. Thus, $G_{3,2}$ is 3-minimal. \square

By Lemma 7(ii) again, every graph in $\mathcal{S}_2^2(K_5)$ that is a subgraph of $G_{3,2}$ is 3-minimal. Figure 12 illustrates $G_{3,2}$ (top left) and its five non-isomorphic proper subgraphs that belong to

$\mathcal{S}_2^2(K_5)$. Notice that one of these graphs (top right of Figure 12) is isomorphic to $G_{3,1}$, the first 3-minimal graph discovered in [EMN06].




FIGURE 12. The graph $G_{3,2}$ (top left) and its five non-isomorphic proper sub-graphs that belong to $\mathcal{S}_2^2(K_5)$ (and thus are 3-minimal)

We close the section by showing that there are no 3-minimal graphs with fewer edges than $G_{3,1}$.

Proposition 28. *Suppose G is a 3-minimal graph. Then $|E(G)| \geq 14$.*

Proof. Since G is 3-minimal, there must exist vertex v_0 where $r_+(G \ominus v_0) = 2$. This implies that $|V(G \ominus v_0)| \geq 6$, and thus $\deg(v_0) \leq 2$. Since ℓ -minimal graphs cannot have cut vertices, we see that $\deg(v_0) = 2$ and $|V(G \ominus v)| = 6$, and so $G \ominus v$ is isomorphic to either $G_{2,1}$ (8 edges) or $G_{2,2}$ (9 edges).

Let v_1, v_2 be the two neighbours of v_0 , and let $A := \{v_0, v_1, v_2\}$ and $B := V(G) \setminus A$. Observe that

$$(10) \quad |E(G)| = \delta(A) + \delta(B) + \delta(A, B).$$

Since $|E(G)| \leq 13$, $\delta(A) = 2$, and $\delta(B) \geq 8$, we obtain $\delta(A, B) \leq 3$. Again, G being 3-minimal implies that $\deg(v_1), \deg(v_2) \geq 2$, and so we obtain that $2 \leq \delta(\{v_1\}, B) + \delta(\{v_2\}, B) \leq 3$. Thus, we may assume without loss of generality that $\delta(\{v_1\}, B) = 1$, and let u be the only neighbour of v_1 in B .

If $\delta(\{v_2\}, B) = 1$, then u, v_1, v_0, v_2 form a sparse path of length 3 (with $\deg(v_1) = \deg(v_0) = \deg(v_2) = 2$), and Proposition 22 implies that G is not 3-minimal. Now suppose $\delta(\{v_2\}, B) = 2$. This means that $\delta(A, B) = 3$, and so from (10) we know that $|E(G)| = 13$, $\delta(B) = 8$, and $G - A$ is indeed isomorphic to $G_{2,1}$ and not $G_{2,2}$.

Next, since $r_+(G) = 3$, we obtain that $r_+(G - u) \geq 2$. However, notice that v_0 is a cut vertex in $G - u$. Thus, if we let $A' := \{u, v_1, v_0\}$ and $B' := V(G) \setminus A'$, then we see that $r_+(G - A') \geq 2$. Since $G - A'$ has 6 vertices, it must be isomorphic to $G_{2,1}$ or $G_{2,2}$. Thus, we see that $\delta(B') \geq 8$. Also, $\delta(A') = 2$ and

$$\delta(A', B') = \delta(\{v_0\}, B') + \delta(\{u\}, B') = 1 + (\deg(u) - 1) = \deg(u).$$

Since $13 = |E(G)| = \delta(A') + \delta(A', B') + \delta(B')$, we obtain that $\deg(u) = \delta(A', B') = 3$, and $\delta(B') = 8$. Thus, $G - A'$ is also isomorphic to $G_{2,1}$ and not $G_{2,2}$. For both $G - A$ and $G - A'$ to be isomorphic to $G_{2,1}$, v_2 must be adjacent to the two neighbours of u in $G \ominus v_0$. Thus, G is isomorphic to the graph shown in Figure 13.




FIGURE 13. Illustrating the proof of Proposition 28

However, notice that $G - w$ has LS_+ -rank 1, which contradicts $r_+(G) = 3$. This completes the proof. \square

7. REVISITING H_k AND CONSTRUCTING SPARSE GRAPHS WITH HIGH LS_+ -RANK

In this section, we revisit the graphs H_k defined in Section 1, and obtain other related graphs with high LS_+ -ranks by applying some of our results on vertex stretching. First, we point out that the LS_+ -rank lower bound in Theorem 2 also applies to some particular subgraphs of H_k . For every $k \geq 3$, define

$$H'_k := H_k - \{1_0, 1_2, 2_0, 2_1\}.$$




FIGURE 14. Several graphs in the family H'_k

Figure 14 illustrates the graphs H'_k for $k = 3, 4, 5$. Notice that $H'_k \in \mathcal{S}_2^{k-2}(K_k)$ for all $k \geq 3$ — this is apparent if one takes the drawings of H'_k from Figure 14 and relabels the vertices 1_1 and 2_2 by 1 and 2 respectively. Then we have the following.

Proposition 29. *For every $k \geq 3$, $r_+(H'_k) \geq \frac{3k}{16}$.*

Proof. For convenience, let $p := \lceil \frac{3k}{16} \rceil - 1$ throughout this proof. Also, given $a, b \in \mathbb{R}$, we define the vector $w_k(a, b) \in \mathbb{R}^{V(H_k)}$ such that

$$[w_k(a, b)]_j := \begin{cases} a & \text{if } j \in \{i_1, i_2 : i \in [k]\}; \\ b & \text{if } j \in \{i_0 : i \in [k]\}. \end{cases}$$

In [AT24], it was shown that there exists real numbers a, b where $w_k(a, b)$ is contained in $\text{LS}_+^p(H_k)$ and violates the inequality

$$(11) \quad w_k(k-1, k-2)^\top x \leq k(k-1),$$

which is valid for $\text{STAB}(H_k)$ [AT24, Lemma 9(ii)].

Now let $w_k(a, b)' \in \mathbb{R}^{V(H'_k)}$ be the vector obtained from $w_k(a, b)$ by removing the four entries that correspond to vertices which are not in H'_k . Then by Lemma 4, we have $w_k(a, b)' \in \text{LS}_+^p(H'_k)$. On the other hand, the fact that $w_k(a, b)$ violates (11) implies that $(k-1)(2ka) + (k-2)(kb) > k(k-1)$, which implies that

$$\alpha_{\text{LS}_+^p}(H'_k) \geq \bar{e}^\top w_k(a, b)' = (2k-2)a + (k-2)b > k-1.$$

However, since $H'_k \in \mathcal{S}_2^{k-2}(K_k)$, it follows from Lemma 20 that $\alpha(H'_k) = k-1$. This implies that $w_k(a, b)' \in \text{LS}_+^p(H'_k) \setminus \text{STAB}(H'_k)$, and that $r_+(H'_k) \geq p+1 \geq \frac{3k}{16}$. \square

In fact, we can use the argument above to find many subgraphs of H'_k for which the LS_+ -rank lower bound given in Proposition 29 applies.

Proposition 30. *Let $G \in \mathcal{S}_2^{k-2}(K_k)$ be a subgraph of H'_k . Then $r_+(G) \geq \frac{3k}{16}$.*

Proof. Again, let $p := \lceil \frac{3k}{16} \rceil - 1$, and let $G \in \mathcal{S}_2^{k-2}(K_k)$ be a subgraph of H'_k . Since $\alpha_{\text{LS}_+^p}(H'_k) > k-1$ (as shown in the proof of Proposition 29, Lemma 7(ii) implies that $\alpha_{\text{LS}_+^p}(G) > k-1$. But then Lemma 20 implies that $\alpha(G) = k-1$. Thus, $r_+(G) \geq p+1 \geq \frac{3k}{16}$. \square

Given a graph G , define the *edge density* of G to be $d(G) := \frac{|E(G)|}{\binom{|V(G)|}{2}}$. For instance, $d(G) = 1$ for complete graphs, and $d(G) = 0$ for empty graphs. An interesting contrast that has emerged in the study of lift-and-project relaxations of the stable set polytope of graphs is that dense graphs tend to have high lift-and-project ranks with respect to operators that produce polyhedral relaxations, whereas graphs from both ends of the density spectrum tend to be of small lift-and-project ranks with respect to semidefinite operators. Thus, it is interesting to note that

$$d(H'_k) = \frac{k^2 - k - 1}{\binom{3k-4}{2}} = \frac{2}{9} + o(k).$$

Moreover, it follows from Proposition 30 that the LS_+ -rank lower bound we showed for H'_k also applies for many subgraphs of H'_k with lower edge densities. For an example, given $k \geq 3$, we define the graph H''_k where $V(H''_k) := V(H'_k)$, with $E(H''_k)$ consisting of the following edges:

- (i) $\{1_1, 2_2\}$;
- (ii) $\{1_1, i_2\}, \{2_2, i_1\}, \{i_0, i_1\}$, and $\{i_0, i_2\}$ for every $i \in \{3, \dots, k\}$;
- (iii) $\{i_2, j_1\}$ for all $i, j \in \{3, \dots, k\}$ where $(j-i) \bmod (k-2) < \frac{k-2}{2}$;

(iv) $\{i_2, j_1\}$ for all $i, j \in \{3, \dots, k\}$ where $j - i = \frac{k-2}{2}$.

Observe that (iv) only contributes edges when k is even. Also, for every $k \geq 3$, notice that H_k'' is a subgraph of H_k' , and that $H_k'' \in \mathcal{S}_2^{k-2}(K_k)$ (see Figures 15 and 16, respectively, for drawings of H_5'' and H_6''). Furthermore, H_k'' has the fewest edges among all graphs in $\mathcal{S}_2^{k-2}(K_k)$. To see this, suppose we start with a complete graph K_k with vertex labels $1_1, 2_2, 3, 4, \dots, k$, and 2-stretch the vertices $3, 4, \dots, k$ to obtain a graph $G \in \mathcal{S}_2^{k-2}(K_k)$. If we define the sets $S_1 := \{1_1\}$, $S_2 := \{2_2\}$, and $S_i := \{i_0, i_1, i_2\}$ for all $i \in \{3, \dots, k\}$, then there must be at least one edge in G joining S_i and S_j for all distinct $i, j \in [k]$. To minimize the number of edges in G , one can ensure that the sets A_1, A_2 are disjoint in each vertex stretching operation. This would result in a graph with exactly one edge joining S_i, S_j for all distinct $i, j \in [k]$, which is indeed the case for H_k'' .

It is easy to check that $|E(H_k'')| = \frac{k^2+3k-8}{2}$, and thus $d(H_k'') = \frac{1}{9} + o(k)$. Thus, we see that there are many subgraphs of H_k' with edge densities between $\frac{1}{9}$ and $\frac{2}{9}$ for which the rank lower bound in Proposition 30 applies.




FIGURE 15. H'_5 (left), H''_5 (centre), and another subgraph of H'_5 in $\mathcal{S}_2^3(K_5)$ with the fewest possible edges (right)

Finally, we point out that we can further stretch the vertices of H_k'' to obtain very sparse graphs with arbitrarily high LS_+ -ranks. Given a graph G and a vertex $v \in V(G)$, define

$$w(v) := \begin{cases} 1 & \text{if } \deg(v) \leq 3; \\ \deg(v) - 1 & \text{if } \deg(v) \geq 4. \end{cases}$$

We also define $w(G) := \sum_{v \in V(G)} w(v)$. Then we have the following.

Lemma 31. *For every graph G , there exists a graph H that can be obtained from G by a sequence of vertex stretching operations where $\deg(v) \leq 3$ for all $v \in V(H)$, and $|V(H)| \leq w(G)$.*

Proof. First, if every vertex in G has degree at most 3, then $H = G$ suffices, so we now assume there exists $v \in V(G)$ with $\deg(v) \geq 4$. Next, we define

$$w_1(G) := |\{v \in V(G) : \deg(v) \geq 4\}|,$$

$$w_2(G) := \sum_{i \in V(G)} \max\{\deg(i) - 3, 0\}.$$

Then $w(G) = |V(G)| + w_1(G) + w_2(G)$ for every graph G . It is helpful to think of $w_2(G)$ as the total “excess” vertex degree in G , and $w_2(G) = 0$ if and only if $\deg(v) \leq 3$ for all $v \in V(G)$. Now notice that

- If $v \in V(G)$ has $\deg(v) = 4$, we can 2-stretch it with $|A_1| = |A_2| = 2$. In this case, we obtain $H \in \mathcal{S}(G)$ with $|V(H)| = |V(G)| + 2$, $w_1(H) = w_1(G) - 1$, and $w_2(H) = w_2(G) - 1$.

- If $v \in V(G)$ has $\deg(v) = 5$, we can 3-stretch it with $|A_1| = |A_2| = 2$ and $|A_3| = 1$. In this case, we obtain $H \in \mathcal{S}(G)$ with $|V(H)| = |V(G)| + 3$, $w_1(H) = w_1(G) - 1$, and $w_2(H) = w_2(G) - 2$.
- If $v \in V(G)$ with $\deg(v) = p \geq 6$, we can 3-stretch it with $|A_1| = |A_2| = 2$ and $|A_3| = p-4$. In this case, we obtain $H \in \mathcal{S}(G)$ with $|V(H)| = |V(G)| + 3$, $w_1(H) \leq w_1(G)$, and $w_2(H) = w_2(G) - 3$. (More precisely, notice that $w_1(H) = w_1(G) - 1$ if $p = 6$ and $w_1(H) = w_1(G)$ if $p \geq 7$.)

In all cases, we see that given a graph G with $w_2(G) > 0$, we can apply a stretching operation to obtain $H \in \mathcal{S}(G)$ such that $w(H) \leq w(G)$ and $w_2(H) < w_2(G)$. Iterating this process would result in a graph H with $w_1(H) = w_2(H) = 0$, which would satisfy $|V(H)| = w(H) \leq w(G)$. \square

Then we have the following.

Theorem 32. *For every $k \geq 5$, there exists a graph G on $k^2 - 4$ vertices such that $\deg(i) \leq 3$ for every $i \in V(G)$, and $r_+(G) \geq r_+(H_k'')$.*

Proof. Given $k \geq 5$, consider the graph H_k'' . Notice that $\deg(1_1) = \deg(2_2) = k - 1$. Moreover, for every $i \in \{3, \dots, k\}$, we have $\deg(i_0) = 2$, $\deg(i_1), \deg(i_2) \geq 3$, and $\deg(i_1) + \deg(i_2) = k + 1$.

Thus, using notation from the proof of Lemma 31, we obtain that $|V(H_k'')| = 3k - 4$, $w_1(H_k'') \leq 2k - 2$, and $w_2(H_k'') = k^2 - 5k + 2$ (as each of $1_1, 2_2$ contributes $k - 4$ to the sum, while i_1 and i_2 together contribute $k - 5$ for every $i \in \{3, \dots, k\}$). Therefore, $w(H_k'') \leq k^2 - 4$. Thus, we can apply Lemma 31 to obtain a graph G from stretching vertices of H_k'' where $|V(G)| \leq k^2 - 4$ and $\deg(v) \leq 3$ for all $v \in V(G)$. Since stretching a vertex cannot decrease the LS_+ -rank of a graph (Proposition 14), the claim follows. \square

Note that the bound $w_1(H_k'') \leq 2k - 2$ is not tight for $k = 5$ and $k = 6$. In those cases, we can obtain a yet better bound as $w(H_5'') = 15$ and $w(H_6'') = 28$. Figure 16 illustrates H_6'' (left), and a stretched graph with $w(H_6'') = 28$ vertices which has maximum degree 3 (right). Note that we suppressed the vertex labels in this figure to reduce cluttering.




FIGURE 16. H_6'' (left), and a 28-vertex graph with maximum degree 3 obtained from stretching H_6'' (right)

Also, since $r_+(H_k'') = \Theta(k)$, it follows from Theorem 32 that there exists a family of graphs G with maximum degree 3 where $r_+(G) = \Omega(\sqrt{|V(G)|})$. This bound asymptotically matches the previously known bound achieved by line graphs of odd cliques, whose vertex degrees grow without bound.

Finally, it was pointed to us by a reviewer that the family of graphs H_k' coincide with the family of graphs G_k in [DV15, page 675] (also studied recently in [Var23, Section 5.5]). It is

very interesting that the families of graphs H'_k have been considered as challenging instances for other but related convex relaxations of the stable set polytope. These graphs are also related to four graphs G_8 , G_{11} , G_{14} and G_{17} considered as minimal obstructions in [PnVZ07] to the hierarchies discussed there which are related to the hierarchy proposed in [dKP02]. These four graphs, as well as the graph H_8 recently studied in [LV23] and [Var23], are related to our family H''_k as they all are among graphs in $\mathcal{S}_2^{k-2}(K_k)$ with the fewest possible edges. These connections raise some more hope that some of our techniques and approaches in this paper may be useful for analyzing other convex relaxations of the stable set polytope.

8. SOME FUTURE RESEARCH DIRECTIONS

In this section, we mention some follow-up questions to our work in this manuscript that could lead to interesting future research.

Problem 33. Is there an ℓ -minimal graph G in $\mathcal{S}_2^{\ell-1}(K_{\ell+2})$ for all $\ell \in \mathbb{N}$?

Results from [LT03, EMN06] show that the answer is “yes” for $\ell \in \{1, 2, 3\}$. Our 4-minimal graph $G_{4,1}$ shows that this is also true for $\ell = 4$. Does the pattern continue for larger ℓ ? And more importantly, how can we verify the LS_+ -rank of these graphs analytically, as opposed to primarily relying on specific numerical certificates?

Problem 34. Given $\ell \in \mathbb{N}$, what are the maximum and minimum possible edge densities of ℓ -minimal graphs?

Given $\ell \in \mathbb{N}$, let $d^+(\ell)$ (resp. $d^-(\ell)$) be the maximum (resp. minimum) possible edge density of an ℓ -minimal graph. It was previously known that $d^+(1) = d^-(1) = 1$ (attained by the 3-cycle), $d^+(2) = \frac{3}{5}$ ($G_{2,2}$), $d^-(2) = \frac{8}{15}$ ($G_{2,1}$), and $d^-(3) \leq \frac{7}{18}$ ($G_{3,1}$). In this work we showed that $d^-(3) = \frac{7}{18}$ (Proposition 28) and $d^+(3) \geq \frac{4}{9}$ ($G_{3,2}$). For $\ell = 4$, the discovery of $G_{4,1}$ and the other 4-minimal graphs presented in Figure 11 show that $d^-(4) \leq \frac{7}{22}$ and $d^+(4) \geq \frac{4}{11}$. Can we prove tight bounds for $d^+(\ell)$ and/or $d^-(\ell)$ in general?

Problem 35. How many non-isomorphic ℓ -minimal graphs are there for each $\ell \geq 1$?

Given $\ell \in \mathbb{N}$, let $c(\ell)$ denote the number of non-isomorphic ℓ -minimal graphs. We know that $c(1) = 1$ (the triangle) and $c(2) = 2$ ($G_{2,1}$ and $G_{2,2}$). We showed in Section 6 that $c(3) \geq 6$ ($G_{3,2}$ and its subgraphs in Figure 12) and $c(4) \geq 7$ ($G_{4,1}$ and its subgraphs in Figure 11). Does $c(\ell)$ grow without bound as ℓ increases? If so, at what rate asymptotically?

Problem 36. What is the fastest growing function f such that there exist graphs G with maximum degree at most three and $r_+(G) = \Theta(f(|V(G)|))$?

Problem 37. What is the fastest growing function f such that there exist cubic graphs G with $r_+(G) = \Theta(f(|V(G)|))$?

We proved in Section 7 that there exist very sparse graphs (maximum degree at most three) with $r_+(G) = \Omega(\sqrt{|V(G)|})$. Since all graphs G with maximum degree at most two satisfy $r_+(G) \leq 1$, Problems 36 and 37 are really about the sparsest graphs with high LS_+ -ranks.

Problem 38. What can we say about the lift-and-project ranks of graphs for other positive semidefinite lift-and-project operators? To start with some concrete questions for this research problem, what are the solutions of Problems 33-37 when we replace LS_+ with Las , BZ_+ , Θ_k , or SA_+ ? (For Problem 33, we may have different sets \mathcal{S} , based on different graph operations, for different lift-and-project operators.)

After LS_+ , many stronger semidefinite lift-and-project operators (such as Las [Las01], BZ_+ [BZ04], Θ_k [GPT10], and SA_+ [AT16]) have been proposed. While these stronger operators are capable of producing tighter relaxations than LS_+ , these SDP relaxations can also be more computationally challenging to solve. For instance, while the LS_+^k -relaxation of a set $P \subseteq [0, 1]^n$ involves $O(n^k)$ PSD constraints of order $O(n)$, the operators Las^k , BZ_+^k and SA_+^k all impose one (or more) PSD constraint of order $\Omega(n^k)$ in their formulations. We briefly mentioned at the end of Section 3 that some of our tools for analyzing LS_+ relaxations can be extended to these other operators, as well as pointed out some related work on other convex relations of the stable set polytope at the end of Section 7. It would be interesting to determine the corresponding properties of graphs which are minimal with respect to the stronger lift-and-project operators.

DECLARATIONS

Conflict of interest: The authors declare that they have no conflict of interest.

REFERENCES

- [AEF14] Néstor E. Aguilera, Mariana S. Escalante, and Pablo G. Fekete. On the facets of lift-and-project relaxations under graph operations. *Discrete Appl. Math.*, 164(part 2):360–372, 2014.
- [AT16] Yu Hin Au and Levent Tunçel. A comprehensive analysis of polyhedral lift-and-project methods. *SIAM J. Discrete Math.*, 30(1):411–451, 2016.
- [AT18] Yu Hin Au and Levent Tunçel. Elementary polytopes with high lift-and-project ranks for strong positive semidefinite operators. *Discrete Optim.*, 27:103–129, 2018.
- [AT24] Yu Hin Au and Levent Tunçel. Stable set polytopes with high lift-and-project ranks for the Lovász–Schrijver SDP operator. *Math. Program.*, 2024.
- [BENT13] Silvia M. Bianchi, Mariana S. Escalante, Graciela L. Nasini, and Levent Tunçel. Lovász-Schrijver SDP-operator and a superclass of near-perfect graphs. *Electronic Notes in Discrete Mathematics*, 44:339–344, 2013.
- [BENT17] Silvia M. Bianchi, Mariana S. Escalante, Graciela L. Nasini, and Levent Tunçel. Lovász-Schrijver SDP-operator, near-perfect graphs and near-bipartite graphs. *Math. Program.*, 162(1-2, Ser. A):201–223, 2017.
- [BENW23] Silvia M. Bianchi, Mariana S. Escalante, Graciela L. Nasini, and Annegret K. Wagler. Lovász-Schrijver PSD-operator and the stable set polytope of claw-free graphs. *Discrete Appl. Math.*, 332:70–86, 2023.
- [BZ04] Daniel Bienstock and Mark Zuckerberg. Subset algebra lift operators for 0-1 integer programming. *SIAM J. Optim.*, 15(1):63–95, 2004.
- [CRST06] Maria Chudnovsky, Neil Robertson, Paul Seymour, and Robin Thomas. The strong perfect graph theorem. *Ann. of Math. (2)*, 164(1):51–229, 2006.
- [dKP02] Etienne de Klerk and Dmitrii V. Pasechnik. Approximation of the stability number of a graph via copositive programming. *SIAM J. Optim.*, 12(4):875–892, 2002.
- [DV15] Cristian Dobre and Juan Vera. Exploiting symmetry in copositive programs via semidefinite hierarchies. *Math. Program.*, 151(2):659–680, 2015.
- [EMN06] Mariana S. Escalante, M. S. Montelar, and Graciela L. Nasini. Minimal N_+ -rank graphs: progress on Lipták and Tunçel’s conjecture. *Oper. Res. Lett.*, 34(6):639–646, 2006.
- [GB08] Michael C. Grant and Stephen P. Boyd. Graph implementations for nonsmooth convex programs. In *Recent advances in learning and control*, volume 371 of *Lect. Notes Control Inf. Sci.*, pages 95–110. Springer, London, 2008.
- [GB14] Michael Grant and Stephen Boyd. CVX: Matlab software for disciplined convex programming, version 2.1. <http://cvxr.com/cvx>, March 2014.
- [GPT10] João Gouveia, Pablo A. Parrilo, and Rekha R. Thomas. Theta bodies for polynomial ideals. *SIAM J. Optim.*, 20(4):2097–2118, 2010.
- [GT01] Michel X. Goemans and Levent Tunçel. When does the positive semidefiniteness constraint help in lifting procedures? *Math. Oper. Res.*, 26(4):796–815, 2001.
- [Knu94] Donald E. Knuth. The sandwich theorem. *Electron. J. Combin.*, 1:Article 1, approx. 48, 1994.

- [Las01] Jean B. Lasserre. An explicit exact SDP relaxation for nonlinear 0-1 programs. In *Integer programming and combinatorial optimization (Utrecht, 2001)*, volume 2081 of *Lecture Notes in Comput. Sci.*, pages 293–303. Springer, Berlin, 2001.
- [Lau02] Monique Laurent. Tighter linear and semidefinite relaxations for max-cut based on the Lovász-Schrijver lift-and-project procedure. *SIAM J. Optim.*, 12(2):345–375, 2001/02.
- [Lau03] Monique Laurent. Lower bound for the number of iterations in semidefinite hierarchies for the cut polytope. *Math. Oper. Res.*, 28(4):871–883, 2003.
- [Lov79] László Lovász. On the Shannon capacity of a graph. *IEEE Trans. Inform. Theory*, 25(1):1–7, 1979.
- [LS91] L. Lovász and A. Schrijver. Cones of matrices and set-functions and 0-1 optimization. *SIAM J. Optim.*, 1(2):166–190, 1991.
- [LT03] László Lipták and Levent Tunçel. The stable set problem and the lift-and-project ranks of graphs. *Math. Program.*, 98(1-3, Ser. B):319–353, 2003. Integer programming (Pittsburgh, PA, 2002).
- [LV23] Monique Laurent and Luis F. Vargas. Exactness of parrilo’s conic approximations for copositive matrices and associated low order bounds for the stability number of a graph. *Mathematics of Operations Research*, 48(2):1017–1043, 2023.
- [PnVZ07] Javier Peña, Juan Vera, and Luis F. Zuluaga. Computing the stability number of a graph via linear and semidefinite programming. *SIAM J. Optim.*, 18(1):87–105, 2007.
- [SA90] Hanif D. Sherali and Warren P. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM J. Discrete Math.*, 3(3):411–430, 1990.
- [ST99] Tamon Stephen and Levent Tunçel. On a representation of the matching polytope via semidefinite liftings. *Math. Oper. Res.*, 24(1):1–7, 1999.
- [Stu99] Jos F. Sturm. Using SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones. *Optim. Methods Softw.*, 11/12(1-4):625–653, 1999. Interior point methods.
- [Var23] Luis F. Vargas. *Sum-of-squares representations for copositive matrices and independent sets in graphs*. PhD thesis, Tilburg University, 2023.
- [Wag22] Annegret K. Wagler. On the Lovász-Schrijver PSD-operator on graph classes defined by clique cutsets. *Discrete Appl. Math.*, 308:209–219, 2022.

APPENDIX A. PROOFS OF LEMMAS 39, 40, AND 41

The following lemmas provide the deferred technical details from the proof of Theorem 26. To reduce cluttering, given $S \subseteq [n]$ we will let $\hat{\chi}_S$ denote the vector $\begin{bmatrix} 1 \\ \chi_S \end{bmatrix} \in \mathbb{R}^{n+1}$.

Lemma 39. *Let Y_0 be as defined in the proof of Theorem 26. Then*

$$Y_0(e_0 - e_1) \in \text{cone}(\text{LS}_+^2(G_{4,1})).$$

Proof. First, notice that $[Y_0(e_0 - e_1)]_1 = 0$. Thus, let $G' := G_{4,1} - 1$ and v be the restriction of $Y_0(e_0 - e_1)$ to the coordinates indexed by $\text{cone}(\text{LS}_+^2(G'))$. Then, by Lemma 4, it suffices to show that $v \in \text{cone}(\text{LS}_+^2(G'))$. Consider the matrix

$$Y_2 := \begin{bmatrix} & 2 & 3 & 4_1 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 & 6_1 & 6_0 & 6_2 \\ 74660 & 25340 & 25340 & 16500 & 57518 & 8662 & 8662 & 57518 & 16500 & 16500 & 49680 & 16500 \\ 25340 & 25340 & 0 & 0 & 25340 & 0 & 0 & 17166 & 8174 & 8363 & 16977 & 0 \\ 25340 & 0 & 25340 & 8174 & 17166 & 0 & 0 & 25340 & 0 & 0 & 16977 & 8363 \\ 16500 & 0 & 8174 & 16500 & 0 & 8320 & 342 & 16158 & 0 & 0 & 14067 & 2433 \\ 57518 & 25340 & 17166 & 0 & 57518 & 0 & 7678 & 41360 & 16158 & 16494 & 34971 & 14067 \\ 8662 & 0 & 0 & 8320 & 0 & 8662 & 984 & 7678 & 342 & 0 & 8662 & 0 \\ 8662 & 0 & 0 & 342 & 7678 & 984 & 8662 & 0 & 8320 & 0 & 8662 & 0 \\ 57518 & 17166 & 25340 & 16158 & 41360 & 7678 & 0 & 57518 & 0 & 14067 & 34971 & 16494 \\ 16500 & 8174 & 0 & 0 & 16158 & 342 & 8320 & 0 & 16500 & 2433 & 14067 & 0 \\ 16500 & 8363 & 0 & 0 & 16494 & 0 & 0 & 14067 & 2433 & 16500 & 0 & 8137 \\ 49680 & 16977 & 16977 & 14067 & 34971 & 8662 & 8662 & 34971 & 14067 & 0 & 49680 & 0 \\ 16500 & 0 & 8363 & 2433 & 14067 & 0 & 0 & 16494 & 0 & 8137 & 0 & 16500 \end{bmatrix}.$$

We claim that $Y_2 \in \widehat{\text{LS}}_+^2(G')$. First, one can verify that $Y_2 \succeq 0$ (a UV -certificate is provided in Table 1). Also, notice that the function f_2 (restricted to $V(G')$) is an automorphism of G' .

Moreover, observe that for all $i, j \in V(G')$, $Y_2[i, j] = Y_2[f_2(i), f_2(j)]$. Thus, by symmetry, it only remains to prove the conditions $Y_2 e_i, Y_2(e_0 - e_i) \in \text{cone}(LS_+(G'))$ for $i \in \{2, 4_1, 4_0, 4_2, 6_1, 6_0\}$.

First, notice that

- $[Y_2 e_{4_0}]_0 = [Y_2 e_{4_0}]_{4_0}, [Y_2 e_{4_0}]_{4_1} = [Y_2 e_{4_0}]_{4_2} = 0$, and that the following matrix certifies that $Y_2 e_{4_0}$ (with the entries corresponding to vertices $4_1, 4_0, 4_2$ removed) belongs to $\text{cone}(LS_+(G' \ominus 4_0))$.

$$Y_{21} := \begin{bmatrix} & 2 & 3 & 5_1 & 5_0 & 5_2 & 6_1 & 6_0 & 6_2 \\ 57518 & 25340 & 17164 & 7678 & 41360 & 16158 & 16496 & 34970 & 14068 \\ 25340 & 25340 & 0 & 0 & 19057 & 6283 & 5860 & 19444 & 0 \\ 17164 & 0 & 17164 & 0 & 17164 & 0 & 0 & 12010 & 5117 \\ 7678 & 0 & 0 & 7678 & 0 & 7678 & 3125 & 4516 & 0 \\ 41360 & 19057 & 17164 & 0 & 41360 & 0 & 10718 & 26585 & 10400 \\ 16158 & 6283 & 0 & 7678 & 0 & 16158 & 5778 & 8385 & 3668 \\ 16496 & 5860 & 0 & 3125 & 10718 & 5778 & 16496 & 0 & 8910 \\ 34970 & 19444 & 12010 & 4516 & 26585 & 8385 & 0 & 34970 & 0 \\ 14068 & 0 & 5117 & 0 & 10400 & 3668 & 8910 & 0 & 14068 \end{bmatrix}$$

- $[Y_2 e_{6_0}]_0 = [Y_2 e_{6_0}]_{6_0}, [Y_2 e_{6_0}]_{6_1} = [Y_2 e_{6_0}]_{6_2} = 0$, and that the following matrix certifies that $Y_2 e_{6_0}$ (with the entries corresponding to vertices $6_1, 6_0, 6_2$ removed) belongs to $\text{cone}(LS_+(G' \ominus 6_0))$.

$$Y_{22} := \begin{bmatrix} & 2 & 3 & 4_1 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 \\ 49680 & 16977 & 16977 & 14068 & 34970 & 8662 & 8662 & 34970 & 14068 \\ 16977 & 16977 & 0 & 0 & 16977 & 0 & 0 & 11129 & 5848 \\ 16977 & 0 & 16977 & 5848 & 11129 & 0 & 0 & 16977 & 0 \\ 14068 & 0 & 5848 & 14068 & 0 & 8220 & 442 & 13626 & 0 \\ 34970 & 16977 & 11129 & 0 & 34970 & 0 & 7578 & 21344 & 13626 \\ 8662 & 0 & 0 & 8220 & 0 & 8662 & 1084 & 7578 & 442 \\ 8662 & 0 & 0 & 442 & 7578 & 1084 & 8662 & 0 & 8220 \\ 34970 & 11129 & 16977 & 13626 & 21344 & 7578 & 0 & 34970 & 0 \\ 14068 & 5848 & 0 & 0 & 13626 & 442 & 8220 & 0 & 14068 \end{bmatrix}$$

- $[Y_2(e_0 - e_2)]_2 = 0$, and that the following matrix certifies that $Y_2(e_0 - e_2)$ (with the entry corresponding to vertex 2 removed) belongs to $\text{cone}(LS_+(G' - 2))$.

$$Y_{23} := \begin{bmatrix} & 3 & 4_1 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 & 6_1 & 6_0 & 6_2 \\ 49320 & 25340 & 16500 & 32178 & 8662 & 8662 & 40354 & 8324 & 8137 & 32703 & 16500 \\ 25340 & 25340 & 6118 & 19222 & 0 & 0 & 25340 & 0 & 0 & 19368 & 5972 \\ 16500 & 6118 & 16500 & 0 & 8107 & 595 & 15905 & 0 & 2465 & 10494 & 6006 \\ 32178 & 19222 & 0 & 32178 & 0 & 7688 & 24409 & 7769 & 5672 & 21928 & 10250 \\ 8662 & 0 & 8107 & 0 & 8662 & 974 & 7688 & 555 & 0 & 5933 & 2729 \\ 8662 & 0 & 595 & 7688 & 974 & 8662 & 0 & 8067 & 70 & 8592 & 0 \\ 40354 & 25340 & 15905 & 24409 & 7688 & 0 & 40354 & 0 & 7763 & 24111 & 16243 \\ 8324 & 0 & 0 & 7769 & 555 & 8067 & 0 & 8324 & 374 & 7950 & 257 \\ 8137 & 0 & 2465 & 5672 & 0 & 70 & 7763 & 374 & 8137 & 0 & 8067 \\ 32703 & 19368 & 10494 & 21928 & 5933 & 8592 & 24111 & 7950 & 0 & 32703 & 0 \\ 16500 & 5972 & 6006 & 10250 & 2729 & 0 & 16243 & 257 & 8067 & 0 & 16500 \end{bmatrix}$$

- $[Y_2(e_0 - e_{4_1})]_{4_1} = 0$, and that the following matrix certifies that $Y_2(e_0 - e_{4_1})$ (with the entry corresponding to vertex 4_1 removed) belongs to $\text{cone}(LS_+(G' - 4_1))$.

$$Y_{24} := \begin{bmatrix} & 2 & 3 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 & 6_1 & 6_0 & 6_2 \\ 58160 & 25340 & 17164 & 57518 & 342 & 8320 & 41360 & 16500 & 16496 & 35612 & 14068 \\ 25340 & 25340 & 0 & 25228 & 0 & 0 & 19229 & 6068 & 5788 & 19552 & 0 \\ 17164 & 0 & 17164 & 17063 & 0 & 0 & 17055 & 0 & 0 & 12187 & 4977 \\ 57518 & 25228 & 17063 & 57518 & 0 & 7946 & 41199 & 16198 & 16378 & 35219 & 13979 \\ 342 & 0 & 0 & 0 & 342 & 340 & 1 & 190 & 0 & 340 & 1 \\ 8320 & 0 & 0 & 7946 & 340 & 8320 & 0 & 8190 & 3046 & 5274 & 0 \\ 41360 & 19229 & 17055 & 41199 & 1 & 0 & 41360 & 0 & 10763 & 26612 & 10417 \\ 16500 & 6068 & 0 & 16198 & 190 & 8190 & 0 & 16500 & 5653 & 8919 & 3601 \\ 16496 & 5788 & 0 & 16378 & 0 & 3046 & 10763 & 5653 & 16496 & 0 & 9091 \\ 35612 & 19552 & 12187 & 35219 & 340 & 5274 & 26612 & 8919 & 0 & 35612 & 0 \\ 14068 & 0 & 4977 & 13979 & 1 & 0 & 10417 & 3601 & 9091 & 0 & 14068 \end{bmatrix}$$

Also, notice that $Y_{21}e_0 = Y_{21}(e_{5_0} + e_{5_2})$. Thus, if we let Y'_{21} be the matrix obtained from Y_{21} by removing the 0th row and column, then we see that $Y'_{21} \succeq 0 \Rightarrow Y_{21} \succeq 0$. The UV -certificates of Y'_{21} , Y_{22} , Y_{23} , and Y_{24} are provided in Table 1.

Next, observe that

$$\begin{aligned} Y_2e_2 &\leq 8291\hat{\chi}_{\{2,4_0,5_0,6_1\}} + 8873\hat{\chi}_{\{2,4_0,5_0,6_0\}} + 72\hat{\chi}_{\{2,4_0,5_2,6_1\}} + 8104\hat{\chi}_{\{2,4_0,5_2,6_0\}}, \\ Y_2e_{4_1} &\leq 6365\hat{\chi}_{\{3,4_1,5_0,6_0\}} + 1811\hat{\chi}_{\{3,4_1,5_0,6_2\}} + 342\hat{\chi}_{\{4_1,4_2,5_1,6_0\}} + 7361\hat{\chi}_{\{4_1,4_2,5_0,6_0\}} \\ &\quad + 617\hat{\chi}_{\{4_1,4_2,5_0,6_2\}} + 4\hat{\chi}_{\{4_1,5_0,6_1,6_2\}}, \\ Y_2e_{4_2} &\leq 642\hat{\chi}_{\{4_1,4_2,5_1,6_0\}} + 7678\hat{\chi}_{\{4_1,4_2,5_0,6_0\}} + 342\hat{\chi}_{\{4_2,5_1,5_2,6_0\}}, \\ Y_2e_{6_1} &\leq 6764\hat{\chi}_{\{2,4_0,5_0,6_1\}} + 1599\hat{\chi}_{\{2,4_0,5_2,6_1\}} + 4\hat{\chi}_{\{4_1,5_0,6_1,6_2\}} + 7300\hat{\chi}_{\{4_0,5_0,6_1,6_2\}} \\ &\quad + 833\hat{\chi}_{\{4_0,5_2,6_1,6_2\}}, \\ Y_2(e_0 - e_{4_0}) &\leq 7254\hat{\chi}_{\{3,4_1,5_0,6_0\}} + 1004\hat{\chi}_{\{3,4_1,5_0,6_2\}} + 489\hat{\chi}_{\{4_1,4_2,5_1,6_0\}} + 6472\hat{\chi}_{\{4_1,4_2,5_0,6_0\}} \\ &\quad + 1291\hat{\chi}_{\{4_1,4_2,5_0,6_2\}} + 137\hat{\chi}_{\{4_1,5_0,6_1,6_2\}} + 495\hat{\chi}_{\{4_2,5_1,5_2,6_0\}}, \\ Y_2(e_0 - e_{4_2}) &\leq 832\hat{\chi}_{\{2,4_0,5_0,6_1\}} + 3414\hat{\chi}_{\{2,4_0,5_0,6_0\}} + 496\hat{\chi}_{\{2,4_0,5_2,6_1\}} + 919\hat{\chi}_{\{2,4_0,5_2,6_0\}} \\ &\quad + 5480\hat{\chi}_{\{3,4_1,5_0,6_0\}} + 2094\hat{\chi}_{\{3,4_1,5_0,6_2\}} + 2827\hat{\chi}_{\{3,4_0,5_0,6_0\}} + 1634\hat{\chi}_{\{3,4_0,5_0,6_2\}} \\ &\quad + 700\hat{\chi}_{\{4_1,5_0,6_1,6_2\}} + 526\hat{\chi}_{\{4_0,5_1,5_2,6_1\}} + 1070\hat{\chi}_{\{4_0,5_1,5_2,6_0\}} + 690\hat{\chi}_{\{4_0,5_0,6_1,6_2\}} \\ &\quad + 455\hat{\chi}_{\{4_0,5_2,6_1,6_2\}} + 126\hat{\chi}_{\{4_2,5_1,5_2,6_0\}} + \frac{44735}{57518}Y_2e_{4_0}, \\ Y_2(e_0 - e_{6_1}) &\leq 275\hat{\chi}_{\{2,4_0,5_0,6_0\}} + 186\hat{\chi}_{\{3,4_1,5_0,6_0\}} + 2333\hat{\chi}_{\{3,4_1,5_0,6_2\}} + 186\hat{\chi}_{\{3,4_0,5_0,6_0\}} \\ &\quad + 5933\hat{\chi}_{\{3,4_0,5_0,6_2\}} + 140\hat{\chi}_{\{4_1,4_2,5_0,6_2\}} + 227\hat{\chi}_{\{4_0,5_1,5_2,6_0\}} + \frac{48880}{49680}Y_2e_{6_0}, \\ Y_2(e_0 - e_{6_0}) &\leq 7474\hat{\chi}_{\{2,4_0,5_0,6_1\}} + 978\hat{\chi}_{\{2,4_0,5_2,6_1\}} + 978\hat{\chi}_{\{3,4_1,5_0,6_2\}} + 7474\hat{\chi}_{\{3,4_0,5_0,6_2\}} \\ &\quad + 1454\hat{\chi}_{\{4_1,5_0,6_1,6_2\}} + 5168\hat{\chi}_{\{4_0,5_0,6_1,6_2\}} + 1454\hat{\chi}_{\{4_0,5_2,6_1,6_2\}}. \end{aligned}$$

Since all incidence vectors above correspond to stable sets in G' , and we already showed earlier that $Y_2e_{4_0}, Y_2e_{6_0} \in \text{cone}(\text{LS}_+(G'))$, we obtain that all the vectors above belong to $\text{cone}(\text{LS}_+(G'))$. Thus, we conclude that $Y_0(e_0 - e_1) \in \text{cone}(\text{LS}_+^2(G_{4,1}))$. \square

Lemma 40. *Let Y_0 be as defined in the proof of Theorem 26. Then*

$$Y_0e_{6_0} \in \text{cone}(\text{LS}_+^2(G_{4,1})).$$

Proof. First, notice that $[Y_0e_{6_0}]_0 = [Y_0e_{6_0}]_{6_0}$, and $[Y_0e_{6_0}]_{6_1} = [Y_0e_{6_0}]_{6_2} = 0$. Thus, let $G' := G_{4,1} \ominus 6_0$ and v be the restriction of $Y_0e_{6_0}$ to the coordinates indexed by $\text{cone}(\text{LS}_+^2(G'))$. Then,

by Lemma 4, it suffices to show that $v \in \text{cone}(LS_+^2(G'))$. Consider the matrix

$$Y_1 := \begin{bmatrix} & 1 & 2 & 3 & 4_1 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 \\ 75020 & 25340 & 17502 & 17502 & 15419 & 51150 & 15911 & 15911 & 51150 & 15419 \\ 25340 & 25340 & 0 & 0 & 0 & 17400 & 7940 & 7940 & 17400 & 0 \\ 17502 & 0 & 17502 & 0 & 0 & 17502 & 0 & 0 & 9571 & 7931 \\ 17502 & 0 & 0 & 17502 & 7931 & 9571 & 0 & 0 & 17502 & 0 \\ 15419 & 0 & 0 & 7931 & 15419 & 0 & 7488 & 396 & 14993 & 0 \\ 51150 & 17400 & 17502 & 9571 & 0 & 51150 & 0 & 15485 & 27920 & 14993 \\ 15911 & 7940 & 0 & 0 & 7488 & 0 & 15911 & 396 & 15485 & 396 \\ 15911 & 7940 & 0 & 0 & 396 & 15485 & 396 & 15911 & 0 & 7488 \\ 51150 & 17400 & 9571 & 17502 & 14993 & 27920 & 15485 & 0 & 51150 & 0 \\ 15419 & 0 & 7931 & 0 & 0 & 14993 & 396 & 7488 & 0 & 15419 \end{bmatrix}.$$

We claim that $Y_1 \in \widehat{LS}_+^2(G')$. First, one can verify that $Y_1 \succeq 0$ (a UV -certificate is provided in Table 1). Also, notice that the function f_2 (restricted to $V(G')$) is an automorphism of G' . Moreover, observe that for all $i, j \in V(G')$, $Y_1[i, j] = Y_1[f_2(i), f_2(j)]$. Thus, by symmetry, it only remains to prove the conditions $Y_1 e_i, Y_1(e_0 - e_i) \in \text{cone}(LS_+(G'))$ for $i \in \{1, 2, 4_1, 4_0, 4_2\}$.

First, notice that $[Y_1 e_{4_0}]_0 = [Y_1 e_{4_0}]_{4_0}, [Y_1 e_{4_0}]_{4_1} = [Y_1 e_{4_0}]_{4_2} = 0$, and that the following matrix certifies that $Y_1 e_{4_0}$ (with the entries corresponding to vertices $4_1, 4_0, 4_2$ removed) belongs to $\text{cone}(LS_+(G' \ominus 4_0))$. (See Table 1 for a UV -certificate.)

$$Y_{11} := \begin{bmatrix} & 1 & 2 & 3 & 5_1 & 5_0 & 5_2 \\ 51150 & 17400 & 17502 & 9571 & 15485 & 27920 & 14993 \\ 17400 & 17400 & 0 & 0 & 7544 & 9856 & 0 \\ 17502 & 0 & 17502 & 0 & 0 & 10450 & 7052 \\ 9571 & 0 & 0 & 9571 & 0 & 9571 & 0 \\ 15485 & 7544 & 0 & 0 & 15485 & 0 & 7941 \\ 27920 & 9856 & 10450 & 9571 & 0 & 27920 & 0 \\ 14993 & 0 & 7052 & 0 & 7941 & 0 & 14993 \end{bmatrix}$$

Now consider the following vectors:

$$\begin{aligned} z^{(1)} &:= [\quad 51150 \quad 17400 \quad 17502 \quad 9571 \quad 0 \quad 51150 \quad 0 \quad 5485 \quad 27920 \quad 14933 \quad]^\top \\ z^{(2)} &:= [\quad 51150 \quad 17502 \quad 17400 \quad 9571 \quad 0 \quad 51150 \quad 0 \quad 14933 \quad 27920 \quad 15485 \quad]^\top \\ z^{(3)} &:= [\quad 57518 \quad 25340 \quad 0 \quad 17164 \quad 14068 \quad 34970 \quad 16496 \quad 16158 \quad 41360 \quad 7678 \quad]^\top \\ z^{(4)} &:= [\quad 49680 \quad 0 \quad 16977 \quad 169771 \quad 40683 \quad 4970 \quad 8662 \quad 8662 \quad 34970 \quad 14068 \quad]^\top \end{aligned}$$

Notice that $z^{(1)} \in \text{cone}(LS_+(G'))$ follows from $Y_1 e_{4_0} \in \text{cone}(LS_+(G'))$ as shown above. Then it follows from the symmetry of G' that $z^{(2)} \in \text{cone}(LS_+(G'))$ as well. $z^{(3)}, z^{(4)} \in \text{cone}(LS_+(G'))$ follows respectively from $Y_2 e_{4_0}, Y_2 e_{6_0} \in \text{cone}(LS_+(G_{4,1} - 1))$, as shown in Lemma 39. Next, observe that

$$\begin{aligned} Y_1 e_1 &\leq 17400 \hat{\chi}_{\{1,4_0,5_0\}} + 7940 \hat{\chi}_{\{1,4_2,5_1\}}, \\ Y_1 e_2 &\leq 9571 \hat{\chi}_{\{2,4_0,5_0\}} + 7931 \hat{\chi}_{\{2,4_0,5_2\}}, \\ Y_1 e_{4_1} &\leq 7931 \hat{\chi}_{\{3,4_1,5_0\}} + 396 \hat{\chi}_{\{4_1,4_2,5_1\}} + 7092 \hat{\chi}_{\{4_1,4_2,5_0\}}, \\ Y_1 e_{4_2} &\leq 414 \hat{\chi}_{\{1,4_0,5_1\}} + 7523 \hat{\chi}_{\{2,4_0,5_0\}} + 7974 \hat{\chi}_{\{3,4_0,5_0\}}, \\ Y_1(e_0 - e_1) &\leq 874 \hat{\chi}_{\{2,4_0,5_0\}} + 3160 \hat{\chi}_{\{2,4_0,5_2\}} + 3160 \hat{\chi}_{\{3,4_1,5_0\}} + 874 \hat{\chi}_{\{3,4_0,5_0\}} + 1100 \hat{\chi}_{\{4_1,4_2,5_0\}} \\ &\quad + 1100 \hat{\chi}_{\{4_0,5_1,5_2\}} + \frac{39412}{49680} z^{(4)}, \\ Y_1(e_0 - e_2) &\leq 1729 \hat{\chi}_{\{1,4_0,5_1\}} + 6165 \hat{\chi}_{\{1,4_0,5_0\}} + 626 \hat{\chi}_{\{1,4_2,5_1\}} + 1749 \hat{\chi}_{\{1,4_2,5_0\}} \\ &\quad + 4298 \hat{\chi}_{\{3,4_1,5_0\}} + 2999 \hat{\chi}_{\{3,4_0,5_0\}} + 1009 \hat{\chi}_{\{4_1,4_2,5_1\}} + 1751 \hat{\chi}_{\{4_1,4_2,5_0\}} \end{aligned}$$

$$\begin{aligned}
& + 1959\hat{\chi}_{\{40,51,52\}} + 971\hat{\chi}_{\{42,51,52\}} + \frac{34235}{57518}z^{(3)} + 27\hat{\chi}_{\emptyset}, \\
Y_1(e_0 - e_{4_1}) & \leq 498\hat{\chi}_{\{1,40,51\}} + 2500\hat{\chi}_{\{1,40,50\}} + 639\hat{\chi}_{\{1,42,51\}} + 6832\hat{\chi}_{\{1,42,50\}} \\
& + 1613\hat{\chi}_{\{2,40,50\}} + 1022\hat{\chi}_{\{2,40,52\}} + 1421\hat{\chi}_{\{3,40,50\}} + 478\hat{\chi}_{\{40,51,52\}} \\
& + 952\hat{\chi}_{\{42,51,52\}} + \frac{20799}{51150}z^{(1)} + \frac{22819}{51150}z^{(2)} + 28\hat{\chi}_{\emptyset}, \\
Y_1(e_0 - e_{4_0}) & \leq 452\hat{\chi}_{\{1,40,51\}} + 7504\hat{\chi}_{\{2,40,50\}} + 7931\hat{\chi}_{\{3,41,50\}} + 7955\hat{\chi}_{\{3,40,50\}} + 28\hat{\chi}_{\emptyset}, \\
Y_1(e_0 - e_{4_2}) & \leq 234\hat{\chi}_{\{1,40,51\}} + 195\hat{\chi}_{\{2,40,52\}} + 7935\hat{\chi}_{\{3,41,50\}} + 93\hat{\chi}_{\{3,40,50\}} \\
& + \frac{46278}{51150}z^{(1)} + \frac{4354}{51150}z^{(2)} + 20\hat{\chi}_{\emptyset}.
\end{aligned}$$

Since all incidence vectors above correspond to stable sets in G' , we obtain that all the vectors above belong to $\text{cone}(\text{LS}_+(G'))$. Thus, we conclude that $Y_0e_{6_0} \in \text{cone}(\text{LS}_+^2(G_{4,1}))$. \square

Lemma 41. *Let Y_0 be as defined in the proof of Theorem 26. Then*

$$Y_0(e_0 - e_{4_1}) \in \text{cone}(\text{LS}_+^2(G_{4,1})).$$

Proof. For convenience, let $G := G_{4,1}$ throughout this proof. Using $Y_0e_{6_0} \in \text{cone}(\text{LS}_+^2(G))$ from Lemma 40 and the symmetry of G , we know that the vector

$$z := \begin{bmatrix} & 1 & 2 & 3 & 4_1 & 4_0 & 4_2 & 5_1 & 5_0 & 5_2 & 6_1 & 6_0 & 6_2 \\ 75020 & 17502 & 25340 & 17502 & 0 & 75020 & 0 & 15419 & 51150 & 15911 & 15911 & 51150 & 15419 \end{bmatrix}^\top$$

belongs to $\text{cone}(\text{LS}_+^2(G))$. Now observe that

$$\begin{aligned}
Y_0(e_0 - e_{4_1}) & \leq \frac{2}{3}z + \frac{1}{3} \left(7726\hat{\chi}_{\{1,40,51,6_0\}} + 17105\hat{\chi}_{\{1,40,50,6_0\}} + 16187\hat{\chi}_{\{1,42,50,6_0\}} \right. \\
& + 8509\hat{\chi}_{\{2,40,50,6_1\}} + 8324\hat{\chi}_{\{2,40,50,6_0\}} + 8509\hat{\chi}_{\{2,40,52,6_0\}} + 9486\hat{\chi}_{\{3,40,50,6_0\}} \\
& \left. + 8017\hat{\chi}_{\{3,40,50,6_2\}} + 7403\hat{\chi}_{\{40,50,6_1,6_2\}} + 9170\hat{\chi}_{\{42,51,52,6_0\}} + 24\hat{\chi}_{\emptyset} \right).
\end{aligned}$$

Notice that all incidence vectors above correspond to stable sets in G . Since $\text{cone}(\text{LS}_+^2(G))$ is a lower-comprehensive convex cone, it follows that $Y_0(e_0 - e_{4_1}) \in \text{cone}(\text{LS}_+^2(G))$. \square

Finally, we provide in Table 1 the UV -certificates of all PSD matrices used in Theorem 26 and Lemmas 39, 40, and 41.

	U															V														
Y ₀	-2	1	1	1	1	1	1	1	1	1	1	1	1	1	11050	1142	1601	781	-196	621	-196	624	621	624	-557	621	165			
	0	114	-8	-107	152	27	-114	202	411	348	-353	-440	-234		1142	15917	-509	-499	-1882	3390	-156	-27	432	-915	-1711	-214	1694			
	0	-57	128	-71	-320	-492	-335	292	270	69	29	221	268		1601	-509	12805	-677	958	-1473	1373	-587	-70	59	-379	235	227			
	-361	-710	-711	-711	-88	434	-88	-88	434	-88	-89	434	-87		781	-499	-677	14384	1269	58	20	75	1698	-1891	-868	901	703			
	0	-135	521	-387	477	230	31	395	-171	-771	-872	-60	741		-196	-1882	958	1269	13455	-739	-38	-706	1544	-297	-936	409	-415			
	0	525	-145	-379	731	-65	-873	-779	-167	409	48	231	462		621	3390	-1473	58	-739	11866	-186	540	-364	186	-530	381	544			
	0	0	0	0	-984	0	984	-984	0	984	-984	0	984		-196	-156	1373	20	-38	-186	8568	-1358	220	469	156	-1144	-163			
	820	-787	-788	-787	1222	-569	1222	1221	-569	1221	1222	-569	1222		624	-27	-587	75	-706	540	-1358	10158	1776	-61	476	-1550	172			
	0	1693	-2745	1052	-901	857	-628	1238	-329	-652	-337	-529	1280		621	432	-70	1698	1544	-364	220	1776	11890	-1026	1209	198	23			
	0	-2192	-370	2562	909	116	-1116	326	-801	1102	-1235	685	14		624	-915	59	-1891	-297	186	469	-61	-1026	13347	-1927	-193	1281			
0	1845	-595	-1250	308	-990	1042	1707	-2079	1128	-2014	3069	-2170		-557	-1711	-379	-868	-936	-530	156	476	1209	-1927	11530	-292	43				
0	-378	1787	-1409	-2148	2972	-1904	1341	-2343	1854	808	-629	50		621	-214	235	901	409	381	-1144	-1550	198	-193	-292	9703	1337				
	8215	2154	2154	2154	1246	6313	1246	1246	6313	1246	1246	6313	1246		165	1694	227	703	-415	544	-163	172	23	1281	43	1337	8773			
Y ₂	-2	1	1	0	0	1	1	0	0	1	1	0	0		4918	41	-26	-35	-29	-89	-233	38	-35	-130	576	-9				
	0	1	-2	72	70	-5	5	-70	-72	0	0	0	0		41	6219	172	148	1281	-511	448	-1143	908	-675	173	-80				
	-67	-50	-51	82	87	-35	-35	88	82	-57	-58	-56				-26	172	4074	-79	-16	399	-532	329	1079	54	115	-922			
	-56	-129	-129	35	-44	-250	-250	-44	35	142	278	143				-35	148	-79	5725	-112	974	-28	-20	201	-548	953	-344			
	0	-110	110	-110	141	384	-384	-141	111	52	0	-52				-29	1281	-16	-112	4852	-1238	-471	-31	-79	-27	-22	-1183			
	0	216	-216	-18	25	125	-125	-25	18	-567	0	567				-89	-511	399	974	-1238	5976	-273	-691	12	-533	26	-469			
	-144	-423	-423	-424	255	253	252	255	-424	-48	263	-48				-233	448	-532	-28	-471	-273	6481	-948	934	-803	289	-295			
	362	-556	-556	538	-221	531	531	-221	538	546	-248	546				38	-1143	329	-20	-31	-691	-948	4677	-53	-1083	36	-14			
	0	928	-928	442	-429	492	-492	429	-442	429	0	-429				-35	908	1079	201	-79	12	934	-53	5504	-279	953	-613			
	-62	82	82	-588	568	-470	-470	568	-588	1034	-1489	1034				-130	-675	54	-548	-27	-533	-803	-1083	-279	4546	77	-53			
0	-1020	1021	1031	-1020	432	-432	1020	-1031	-371	0	371				576	173	115	953	-22	26	289	36	953	77	9521	-142				
	3377	1203	1203	673	2686	338	338	2686	673	717	2294	717				-9	-80	-922	-344	-1183	-469	-295	-14	-613	-53	-142	4375			
Y'21	-44	-1	0	-20	-1	0	0	0	0						3758	-873	-519	-1	636	9	-429	-479								
	233	22	465	229	20	-410	-389	86							-873	7118	-18	2774	-91	-423	-1334	-135								
	385	793	-241	-149	305	244	-460	-545							-519	-18	1822	17	282	-78	70	-402								
	439	137	-877	-360	468	-700	-18	959							-1	2774	17	8485	-2504	-708	22	687								
	1565	-1333	-1075	528	-1287	508	-788	-327							636	-91	282	-2504	6178	-107	586	225								
	655	-1159	930	-530	1923	1995	-641	1022							9	-423	-78	-708	-107	3903	-635	-904								
	-1375	1105	-499	1255	-949	1180	-1505	1780							-429	-1334	70	22	586	-635	5449	602								
	2466	1620	307	4002	775	986	3444	835							-479	-135	-402	687	225	-904	602	5052								
Y22	5	-3	-3	-1	-1	-3	-3	-1	-1						245290	67346	43642	33983	37	8790	67640	11795	6716							
	-1	-12	9	8011	8038	-189	190	-8035	-8009						67346	262681	-47235	-60196	1386	-15331	-35125	-4652	21255							
	-7048	-7604	-7605	6746	7345	-5664	-5664	7345	6752						43642	-47235	230858	-23610	-7313	-78942	460	18674	-1875							
	-1	8338	-8339	10389	-11032	-29236	29235	11032	-10389						33983	-60196	-23610	293133	-56794	-5587	28035	3646	3783							
	5767	23052	23052	28635	-19951	-26290	-26290	-19952	28635						37	1386	-7313	-56794	116143	891	-8115	-5728	-26068							
	0	82739	-82739	16909	-15913	35612	-35612	15913	-16909						8790	-15331	-78942	-5587	891	159853	-29478	-14768	-1859							
	32757	-40385	-40385	61612	-35034	61878	61878	-35034	61613						67640	-35125	460	28035	-8115	-29478	217945	-6899	-36239							
	0	55744	-55744	-88585	87206	-48485	48485	-87206	88585						11795	-4652	18674	3646	-5728	-14768	-6899	124459	2908							
	237999	88332	88332	60363	177671	34664	34664	-177671	60363						6716	21255	-1875	3783	-26068	-1859	-36239	2908	120960							
	Y23	-1	-1	0	0	-1	0	0	-1	0	0	0				3268	-126	-24	-112	-237	198	-265	583	192	-200	631				
3		0	-10	-5	-6	-1	11	-3	-6	23	24				-126	5487	-1306	-32	22	-488	501	-233	-286	8	753					
-1		7	26	29	7	-50	-61	-4	4	18	20				-24	-1306	3523	-177	4	802	29	9	-228	-99	75					
-56		-71	75	74	-61	-45	63	55	-78	-15	-13				-112	-32	-177	2815	-3	-4	-57	928	47	597	-258					
-20		-29	71	77	-26	125	-34	-186	-29	4	12				-237	22	4	-3	4247	119	-1336	82	-822	-708	8					
3		-6	219	-154	-525	-3	-17	16	490	160	-165				198	-488	802	-4	119	3240	189	165	337	-120	68					
71		395	245	-200	-264	256	-195	248	-296	-219	283				-265	501	29	-57	-1336	189	4825	96	-29	796	459					
-382		1053	-281	-46	-387	-664	309	-650	-488	157	-530				583	-233	9	928	82	165	96	3654	262	374	-122					
-61		135	-946	906	-666	-101	25	-28	493	-776	714				192	-286	-228	47	-822	337	-29	262	4115	-118	199					
25		-59	764	-724	359	-748	769	-732	442	-855	883				-200	8	-99	597	-708	-120	796	374	-118	4024	-139					
	2625	1525	824	1782	403	373	2243	358	385	1800	822				631	753	75	-258	8	68	459	-122	199	-139	4828					
Y24	-23	-1	-1	0	0	-1	-1	0	0	-1	0	-1			3039	256	-4	11	33	616	259	212	195	517	213					
	20	7	6	-12	-25	0	-15	-19	0	5	10				256	2748	-376	-1	-168	-556	192	-60	147	-215	-197					
	-4	-1	2	44	43	-2	-42	-42	-2	-2	0				-4	-376	2699	-164												