Scale-rich network metamaterials grow by local rules on a recursive cell division

Alexei Vazquez[1, *]

[1] *Nodes & Links Ltd, Salisbury House, Station Road, Cambridge, CB1 2LA, UK*

(Dated: November 12, 2026)

Scale-rich (SR) metamaterials are planar ligament networks grown by nucleating a segment at a random point and angle and extending it until it meets the existing structure, with a thickness that decays with the time of birth. In the dual graph, where ligaments are nodes and junctions are links, the degree distribution has a k^{-3} tail. Here we show that, at zero thickness, the SR construction is Cowan's iterated division of cells with area-weighted selection, and that its dual graph grows by a face-local rule: a new node joins the two nodes that carry two sides of one face. Cowan's uniform split fractions make the cell genealogy a random binary search tree. The degree of an interior ligament equals its number of face incidences, so linear preferential attachment on incidences would give the Barabási–Albert law $p_k = 12/[k(k+1)(k+2)]$. The geometric model does not grow that way. A ligament is hit in proportion to its length, which is fixed at birth, and its degree is a Poisson variable with mean $4\ell_i(\sqrt{T}-\sqrt{i})/c_L$: at fixed length and age the variance-to-mean ratio is close to one, and early and late hits are uncorrelated. Length acts as a fitness that the cell hierarchy assigns at birth, $\ell_i \propto i^{-1/2}$, and the k^{-3} tail follows from it. Using integral geometry we compute the probability that a new ligament cuts off a corner of its cell ($2\ln 2/\pi$ for a square), which sets the triangle closure and the clustering spectrum $C(k) \simeq 2[p_0 + p(k-2)]/[k(k-1)]$. A purely topological face-splitting rule reproduces $P(k)$, $C(k)$ and $k_{nn}(k)$, but through reinforcement and with longer distances. Square-closing splits raise the shortest-path multiplicity above its value at fixed degree sequence. Thickness, jamming, angle sets, alignment and density gradients leave $\gamma = 3$, whereas coherent orientation fields, which make the hit rate grow faster than length, give $\gamma \simeq 2.4$–2.5 up to $T = 10^6$. At fixed decay exponent, the reported per-ligament energy law cannot distinguish degree from length or age.

I. INTRODUCTION

Both *et al.* [1] introduced scale-rich (SR) mechanical metamaterials, two-dimensional ligament networks in which length, thickness and connectivity span orders of magnitude. The construction has two parameters. At step $t = 1, 2, \ldots, T$ a point is drawn uniformly in the unit square together with a uniform angle, and a ligament of thickness $\lambda_t = \lambda_0 t^{-\alpha}$ is grown both ways from it until each end meets an existing ligament or the boundary. Every junction is a T. Following the road-network convention [2, 3], the authors represent each ligament as a node and each junction as a link. They report power-law distributions of thickness, length and degree, with $P(k) \sim k^{-3}$ attributed to a spatial preferential attachment, because longer ligaments offer more opportunities to be hit. They also report a per-ligament elastic energy law $U_i \sim k_i^{3/2}\lambda_i^{1/2} f(\theta_i)$ and a density scaling of the total energy controlled by α [1].

Two questions are natural from the point of view of network growth. Which growth class does the SR dual graph belong to, and which of its properties are fixed by that class rather than by the design parameters (α, λ_0)? Local growth rules, in which a new node attaches to an existing node and to some of its neighbours, generate preferential attachment without global information, together with a hierarchical clustering spectrum $C(k) \sim 1/k$ and degree correlations [4–7]. Growth with node-specific fitness generates scale-free degrees in a different way, through attractiveness fixed at birth rather than through reinforcement [8]. The geometric construction is itself not new: with zero thickness it is Cowan's iterated cell division [9], a member of the family of random tessellations that also contains the STIT model [10, 11] and related to hierarchical crack patterns formed by successive domain divisions [12, 13]. We show that the SR dual grows by a face-local rule, as local-growth models do, but that its degree law is of the fitness type: the fitness is the ligament length, assigned at birth by the cell hierarchy. Both mechanisms give $\gamma = 3$ in two dimensions; they are told apart by the fluctuations of the degree at fixed length and age. Keeping both references in view, the recursive division and the local graph rule, turns most of the reported network statistics into exact or asymptotically exact statements.

The paper is organised as follows. Section II maps the SR construction onto Cowan's division and derives the statistics of the cell hierarchy. In Sec. III we write the dual growth as a face-local rule, contrast preferential attachment with length as fitness, and test which one the geometry realises. Section IV computes the corner-cut probability by integral geometry and derives the clustering spectrum. Section V asks which properties survive when geometry is removed, and Sec. VI turns to distances and shortest-path multiplicity. Section VII identifies which rules move the degree exponent, and Sec. VIII discusses what this means for the reported structure–property relations.

* alexei@nodeslinks.com




FIG. 1. (a) An SR realisation with $T = 60$ ligaments. The line width increases with $\lambda_t = \lambda_0 t^{-1/2}$. (b) Its dual graph: each ligament is a node, placed at the ligament midpoint with a size that grows with its degree, and each junction is a link. (c) A new ligament (orange) splits one cell. Dual nodes of the cell's sides are shown in blue, and the dotted lines join sides that meet at a corner. If the new ligament cuts off a corner, it joins two adjacent sides and closes a triangle in the dual. Otherwise it joins two sides at distance two and closes a square.

II. THE SR CONSTRUCTION IS A RECURSIVE CELL DIVISION

We first set $\lambda_0 = 0$ and return to thickness at the end of this section. The current ligaments partition the square into convex polygonal cells. A uniform point falls in cell K with probability equal to its area A_K. Because K is convex and bounded by ligaments, the new ligament grown from that point stops on the boundary of K: it is the chord of K through a uniform point of K at a uniform angle. The SR model is therefore the iterated division of cells with *area-weighted selection* and the *chord through a uniform point* division rule, model (a) of Fig. 2 in Cowan [9]. Cowan rescales the window after each division, by a factor that keeps the mean cell size fixed, so that a stationary limit exists; the rescaling is a similarity and does not change the topology. STIT tessellations differ in both rules: cells are selected by perimeter and cut by isotropic uniform random lines [10, 11]. Gilbert's model [14], used as a reference architecture in Ref. [1], grows all segments simultaneously and is a different process. Hierarchical crack patterns in drying layers also divide domains one at a time and form T-junctions, but the division is deterministic or nearly so, set by the domain shape and the layer thickness [12, 13]. Throughout, the boundary of the square is not a node of the dual graph; 0.55% of ligament ends lie on it.

Corner numbers. A junction placed on the side of a neighbouring cell is a straight angle there, so the corner number n of a cell is fixed at birth. A chord ending on two sides gives each daughter the two new corners at its ends, so $n_1 + n_2 = n + 4$. Starting from the square, after t steps there are $t + 1$ cells and $4 + 4t$ corners, so the mean corner number is exactly $\langle n \rangle = 4$. Measured number fractions for $n = 3, \ldots, 7$ are $0.301, 0.452, 0.200, 0.042, 0.005$. A Markov-chain treatment of the side number under division, as done for proliferating epithelia [15], would give these fractions once the corner-cut probabilities of Sec. IV are known by side number; the rules differ from the epithelial case because T-junctions do not add corners to the neighbours.

Uniform splitting and the binary search tree. Let K be a convex body, x a uniform point in K, and θ any direction independent of x. For fixed θ, the offset s of the line through x along the normal direction has density $w(s)/A_K$, where $w(s)$ is the chord length at offset s. The area on one side of the line is $\int_{-\infty}^{s} w(s')ds'$, so the area fraction is the cumulative distribution function of s evaluated at s. It is therefore uniform on $(0, 1)$, whatever the shape of K and the law of θ. This uniformity and the resulting exponential law of rescaled areas are Cowan's results [9]. They have a compact restatement. Label each cell by the interval of $[0, 1]$ that it would occupy if areas were laid on a line. With area-weighted selection, the sequence of splits is exactly that of a binary search tree (BST) built from uniform keys, the familiar random scission of an interval: the key falls in an interval with probability equal to its length and splits it at the key [16]. The cell areas after T steps are therefore the $T + 1$ spacings of T uniform points, $P(TA > x) \to e^{-x}$ and $\sum_K A_K^2 = 2/(T + 2)$, and the genealogy of cells inherits the BST statistics: the depth of the split at step t grows as $2 \ln t$, with variance $2 \ln t$, and areas never multiscale. Our simulations give a depth slope 1.983, a variance slope 1.980 and $\sum A^2 = 9.998 \times 10^{-5}$ at $T = 2 \times 10^4$ against 9.999×10^{-5} [Fig. 2(a,b)]. The multiscaling of Krapivsky and Ben-Naim [17] arises with rectangular cuts and lives in the lengths, not in the areas. Axis-aligned chords reproduce their width exponents $\gamma(a) = (a - \sqrt{a^2 + 4})/2$ within 2%, while with isotropic angles the perimeter mo-

ments scale as $t^{1-\alpha/2}$, without multiscaling.

Lengths and the age of the host. Cells at time t have linear size $t^{-1/2}$, and the mean length of the ligament born at t is $\langle \ell_t \rangle \simeq c_\ell t^{-1/2}$ with $c_\ell = 1.16$. The total length grows as $L(t) \simeq c_L \sqrt{t}$ with $c_L = 2c_\ell = 2.33$. The first ligament has $\langle \ell_1 \rangle = 0.945$, the ℓ_0 of Ref. [1]. Section III shows that an end of ligament t lands on a given ligament with probability proportional to its length. Replacing that length by its mean, a mean-field step, the host's birth index h satisfies $P(h) \simeq \ell_h / L(t) \simeq \frac{1}{2}(ht)^{-1/2}$. Then $P(h \le xt) = \sqrt{x}$, that is, asymptotically,

$$h/t \;\to\; U^2, \qquad U \sim \text{Uniform}(0,1). \tag{1}$$

Equivalently, $\ln(t/h)$ is exponential with mean 2 [Fig. 2(c)]. Measured values of $\langle (h/t)^\alpha \rangle$ are 0.667, 0.498 and 0.330 for $\alpha = 1/4$, 1/2 and 1, against $1/(1+2\alpha) =$ 0.667, 0.500 and 0.333.

Thickness. With $\lambda_t = \lambda_0 t^{-\alpha}$, Eq. (1) gives the thickness ratio between a new ligament and its host as $\lambda_t/\lambda_h = U^{2\alpha}$, again asymptotically and in mean field. Its law does not depend on t, so every generation couples to the frame it attaches to in the same statistical way.

The fill fraction without overlaps, $\rho \simeq \lambda_0 \sum_t \langle \ell_t \rangle t^{-\alpha}$, converges only for $\alpha > 1/2$. This recovers the $\alpha = 1/2$ line of the phase diagram of Ref. [1], but not its second branch: for $\alpha > 1/2$ the structure also jams when $\lambda_0 > [\ell_0 \zeta(2\alpha)]^{-1}$ [1]. That branch comes from the depletion of free area, which shortens later ligaments and which the overlap-free estimate ignores. Bisecting λ_0 in finite-width simulations, in which ligaments nucleate only in free area and stop at the surface of the ligaments they meet ($T = 2 \times 10^4$, two realisations per λ_0), we find the threshold $\lambda_0^* = 0.233$, 0.363, 0.643 and 0.958 for $\alpha = 0.6$, 0.75, 1 and 1.5. The formula of Ref. [1] gives 0.189, 0.405, 0.643 and 0.879, while the overlap-free estimate $1/\sum_t \langle \ell_t \rangle t^{-\alpha}$ gives 0.132, 0.219, 0.375 and 0.627, too low by a factor of about 1.7. Thickness removes area and stops growth in the jammed phase, but it does not change the rule by which cells are split (Sec. VII). In what follows the network is the zero-thickness one unless stated otherwise.

III. A FACE-LOCAL GROWTH RULE: PREFERENTIAL ATTACHMENT OR FITNESS?

In the dual graph, step t reads as follows: choose a face (a cell) with probability equal to its area, choose the two sides of that face that the chord meets, and join the new node t to the two ligaments that carry them [Fig. 1(c)]. The new node never links to a node outside the chosen face. It is a local rule in the sense of Ref. [4], with faces in place of neighbourhoods. Which two sides are hit is decided by the geometry, and two different mechanisms are compatible with the bookkeeping.

Incidences equal degree. Call an incidence a pair (ligament i, cell K) in which i contributes a side to K. A new ligament has two incidences, one on each side, and two links. When a later ligament ends on i, it adds one link to i and cuts one of i's sides in two, adding one incidence. For a ligament whose ends are both interior, the number of incidences therefore equals the degree k_i at all times. Each step creates two hits and four incidences, so the total number of incidences is $4t$, in agreement with $\langle n \rangle = 4$. This identity is exact.

Mechanism A: preferential attachment on incidences. If every incidence were equally likely to be hit, node i would gain links at rate $2k_i/4t$, linear preferential attachment with

$$\frac{dk_i}{dt} = \frac{k_i}{2t}, \qquad k_i(t) = 2\sqrt{t/i}, \tag{2}$$

the Barabási–Albert rule with $m = 2$ [18]. Its master equation [19, 20] gives

$$p_k = \frac{12}{k(k+1)(k+2)}, \qquad P(k' \ge k) = \frac{6}{k(k+1)}, \tag{3}$$

for $k \ge 2$, so $\gamma = 3$ and $K^2 P(k \ge K) \to 6$. Each hit adds an incidence and therefore raises the future hit rate: early hits beget late hits.

Mechanism B: length as fitness. In the geometry, a chord ends on a side with a probability that grows with the side's length, and the total length that ligament i exposes is its length ℓ_i, fixed at birth. Being hit splits a side but adds no length, so there is no reinforcement. Two ends land per step on a total length $L(t) = c_L\sqrt{t}$, so the hit rate per unit length is $2/L(t)$, and the degree is a sum of independent rare events with mean

$$\langle k_i \rangle - 2 = \frac{4\ell_i}{c_L}\left(\sqrt{T} - \sqrt{i}\right). \tag{4}$$

This is the "more opportunities to be hit" argument of Ref. [1], made quantitative. It is a fitness model [8] in which the fitness is the birth length. The cell hierarchy assigns it: $\langle \ell_i \rangle = c_\ell i^{-1/2}$, so that, with $4c_\ell/c_L = 2$, Eq. (4) reduces on average to $k_i \simeq 2\sqrt{T/i}$, the same growth law as Eq. (2). The randomness, however, sits in the birth length, $x = \ell_i/\langle \ell_i \rangle$, rather than in the history of hits. With $k \simeq 2x\sqrt{T/i}$ and i uniform,

$$P(k \ge K) \simeq \frac{4\langle x^2 \rangle}{K^2}, \tag{5}$$

again $\gamma = 3$, with an amplitude set, in mean field, by the spread of birth lengths. A hub is predictable at birth from its length.

Which mechanism the geometry realises. Both mechanisms predict $\gamma = 3$ and the same mean growth law, so the exponent cannot decide between them. We apply three tests (three realisations with $T = 10^5$ unless stated). (i) At fixed predicted mean μ_i from Eq. (4), mechanism B makes $k_i - 2$ nearly Poisson. The measured variance-to-mean ratio is 1.05–1.06 for $\mu_i \in [2,4)$, 1.14–1.18 for $[4,8)$ and 1.36–1.44 for $[8,16)$, and $\text{var}(k_i - 2 - \mu_i)/\langle \mu_i \rangle = 0.98$–$1.09$ for ligaments born in $[100, T/20)$.




FIG. 2. The cell hierarchy of the SR construction (six realisations, $T = 2 \times 10^4$). (a) The area fraction of a daughter cell is uniform. (b) Rescaled cell areas follow e^{-x}, the spacing law of uniform points. (c) The birth index h of the ligament hit by a new ligament t follows the mean-field law $h/t \to U^2$, for which $\ln(t/h)$ has density $\frac{1}{2}e^{-x/2}$.

The excess for the largest μ_i (2–4.7 for $\mu_i > 16$, a few hundred early ligaments) reflects the local environment of the oldest ligaments, which the mean-field rate $2/L(t)$ ignores. Under mechanism A the variance at fixed age exceeds the mean by a factor that grows with the mean: in the topological surrogate of Sec. V the ratio is 10 for ligaments born at $i \simeq 1000$ (mean degree 16–17) and 51–91 for those born at $i \simeq 100$ (mean 66–68). (ii) Let e_i be the number of hits on ligament i during $[i, 2i)$ and l_i the number during $[2i, T)$. After subtracting the expectations of Eq. (4), e_i and l_i are uncorrelated ($r = -0.027$, -0.015 and 0.045). In the topological surrogate of Sec. V, which realises mechanism A, the same correlation at fixed age is 0.46–0.50. (iii) The tail amplitude does not separate the mechanisms cleanly. At $T = 10^6$ (five realisations) the measured $K^2 P(k \geq K)$ is 5.78 ± 0.02, 5.75 ± 0.03 and 5.75 ± 0.06 at $K = 32$, 64 and 128. The exact finite-K value of Eq. (3), $6K/(K+1)$, is 5.82, 5.91 and 5.95, so the data lie below preferential attachment by 5σ at $K = 64$. Drawing each degree as 2 plus a Poisson variable with the mean of Eq. (4), using the actual birth lengths ($\langle x^2 \rangle = 1.41$, so $4\langle x^2 \rangle = 5.64$), gives 5.57 ± 0.02, 5.56 ± 0.09 and 5.59 ± 0.07, below the data. The excess over the Poisson fitness model comes from the extra spread of the oldest ligaments noted in test (i).

The geometric SR dual therefore grows by mechanism B: a face-local rule in which old nodes are hit in proportion to a fitness fixed at birth. Mechanism A is what the same face-local bookkeeping produces when geometry is removed (Sec. V). In two dimensions the two coincide in the exponent because the hierarchy makes the fitness scale as $i^{-1/2}$, exactly the growth law of preferential attachment with $m = 2$. They differ in their fluctuations, and they differ in three dimensions, where the two mechanisms predict different exponents (Sec. VII).

At $T = 10^5$ the measured $K^2 P(k - 2 \geq K)$ at $K = 8, 16, 32$ is $3.53, 4.45, 5.19$ [Fig. 3(a)], and the ratio of measured to predicted degree in Eq. (4) averages 1.006, with correlation 0.95 [Fig. 3(b)]. The degree of the cohort born at i decays as $i^{-\beta}$ with $\beta = 0.50$–0.51.

IV. CORNER CUTS, TRIANGLES AND CLUSTERING

A new node closes a triangle in the dual when the two ligaments it joins are already linked. Locally this happens when the chord cuts off a single corner of its cell, since the two sides meeting at that corner belong to linked ligaments.

Integral-geometry formula. Chords through a uniform point at a uniform angle sample lines with density $w(s)\, ds\, d\theta/(\pi A_K)$. For a fixed direction θ, order the corners by their projection on the normal. The chord cuts off the lowest corner alone when its offset lies between the projections of the lowest and second-lowest corners. The integral of $w(s)$ over that range is the area $A_{\rm lo}(\theta)$ cut off by the line through the second-lowest corner, and likewise $A_{\rm hi}(\theta)$ at the other end. Hence

$$P_{\rm corner}(K) = \frac{1}{\pi A_K} \int_0^\pi \left[A_{\rm lo}(\theta) + A_{\rm hi}(\theta) \right] d\theta. \qquad (6)$$

The line measure is the standard one of integral geometry [21], weighted here by the chord length because the line passes through a uniform point. For a triangle every chord cuts off a corner, so $P_{\rm corner} = 1$. For a square, Eq. (6) gives $2 \ln 2/\pi = 0.4413$ in closed form. Numerical quadrature of Eq. (6) gives 0.303 for a 1×4 rectangle and 0.232 and 0.136 for regular pentagons and hexagons. Evaluated on the simulated SR cells, the mean of Eq. (6) by corner number $n = 3, \ldots, 7$ is 1.000, 0.461, 0.254, 0.152 and 0.10, and its area-weighted average is 0.447–0.449 (numerical). The frequency of corner cuts measured directly in the growth is 0.447–0.449. This agreement checks the formula and the code against each other; it is not an independent prediction, because the cell shapes are taken from the simulation.




FIG. 3. Dual graph of the SR construction, three realisations with $T = 10^5$. (a) Cumulative degree distribution and Eq. (3); on this scale it cannot be told apart from Eq. (5). (b) Degree against the prediction (4) from birth length and age (light dots: a random sample of ligaments; circles: binned means). (c) Clustering spectrum and Eq. (7) with $p_0 = 0.494$, $p = 0.53$. (d) Mean neighbour degree; the dashed line is $\ln T$.

Triangle bookkeeping. A corner cut creates a triangular daughter unless the parent was already a triangle, which can only split into $(3, 4)$. The number fraction of triangles is therefore $f_3 = \langle P_{\text{corner}} \rangle - w_3$, where w_3 is the area fraction of triangles. Measured values are $0.30 = 0.449 - 0.147$.

Non-local closures. The fraction of new nodes that close a triangle is 0.496, not 1/2. Corner cuts account for 0.448. The other 0.048 join two non-adjacent sides of their cell whose ligaments already meet outside it. The remaining 0.50 of new nodes join two ligaments at distance two or more around the cell: distance two closes a square, and in cells with $n \geq 6$ the two sides can be three steps apart around the face.

Clustering spectrum. Links between existing nodes are never added. A node therefore acquires triangles only at birth, with probability p_0, or when it is hit and the other end of the new ligament lands on one of its neighbours, with probability p per hit. With constant p the number of triangles at node i is $e_i = p_0 + p(k_i - 2)$, and

$$C(k) = \frac{2[p_0 + p(k-2)]}{k(k-1)} \simeq \frac{2p}{k}, \tag{7}$$

the hierarchical form produced by triadic closure [4, 5, 7]. Counting the per-hit triangle events directly, p is not constant: it is 0.41 when the hit node has degree 2, 0.48 for degrees 3–4, 0.53 for 5–9, and 0.56–0.58 for degrees 10 and above (two realisations, $T = 10^5$, each bin $\geq 10^4$ hits). Hits on hubs are more likely to land next to a neighbour because hubs are long and carry many short stems. With $p_0 = 0.494$ and a single fitted $p = 0.53$ (0.56 when the fit is restricted to $k > 7$), Eq. (7) is within 2% of the data for $k \geq 15$ [Fig. 3(c)]. The mean clustering is 0.38. At $k = 2$ the measured $C = 0.536$ exceeds p_0, because corner-cut ligaments are short and rarely hit again. The degree assortativity is $r \simeq -0.02$. The mean neighbour degree $k_{nn}(k)$ falls from 19.5 at $k = 2$ to a plateau of 11.1–11.5 for hubs, close to the Barabási–Albert value $(m/2) \ln T = \ln T = 11.5$ [Fig. 3(d)].

In the language of local rules, the SR dual mixes two moves. With probability $\simeq 0.45$ the new node attaches to both ends of a link (the corner), as in the edge-attachment model of Dorogovtsev, Mendes and Samukhin [6], except that the link is selected with probability proportional to the area of its cell rather than uniformly. With probability $\simeq 0.50$ it attaches to two nodes at distance two or more on a face cycle, mostly closing a square, and with probability 0.05 it closes a triangle non-locally. The edge-attachment model has $\gamma = 3$ and clustering 0.74 [6, 22]. The SR dual has the same γ and half the clustering; the other half of the closures went into squares. It is also the two-dimensional relative of the random Apollonian network [23, 24], in which faces are selected and a node's degree is proportional to the number of faces it bounds.

V. WHAT TOPOLOGY ALONE REPRODUCES

To separate the role of geometry, we grew a purely topological surrogate. Each face is stored as a cyclic list of the nodes that carry its sides. At each step a face is chosen uniformly. If it is a triangle, two of its sides are joined; otherwise two adjacent sides are joined with probability p_4, or two non-adjacent sides with probability $1-p_4$. The face is then split into two, with $n_1+n_2 = n+4$ as in the tessellation. No lengths or positions are used.

Because every side of the chosen face is equally likely to be hit, every incidence is equally likely to be hit, and the surrogate realises mechanism A of Sec. III exactly. Every face-local variant we tried (faces chosen uniformly, by number of sides, or by a random-split area) reproduces Eq. (3): $\gamma_{\text{MLE}}(k \geq 10) = 2.88$–$2.91$ and the same K^2P amplitudes at $K \leq 32$. The single parameter $p_4 = 0.29$ also gives a mean clustering of 0.386 (SR, 0.377) and places $C(k)$ and $k_{nn}(k)$ on the SR curves. The surrogate

reaches these statistics through reinforcement, as test (ii) of Sec. III shows, whereas the geometric model reaches them through length. The low-order degree statistics therefore do not identify the mechanism.

The surrogate differs on distances. At $T = 10^5$ its mean distance is 14.1 against 8.8 for SR. Both grow logarithmically, but with prefactors 1.43 and 0.60 [Fig. 5(a)]. In the geometric model chords hit long sides preferentially, and long sides belong to old, long ligaments, so new nodes attach close to the hubs. In the surrogate, sides are hit regardless of the age of the ligament that carries them, and repeated cuts across faces can produce long chains of squares. Geometry thus matters for the properties that depend on which particular sides are hit, such as distances, and not for $P(k)$, $C(k)$ or $k_{nn}(k)$.

VI. FEWEST-TURN PATHS AND THEIR MULTIPLICITY

A shortest path in the dual graph is a route through the material that changes ligament the fewest times. For a random graph with given degrees, the mean number of shortest paths between two nodes is $\langle\mu\rangle \simeq c_2/e$ with $c_2 = \langle k(k-1)\rangle/\langle k\rangle$, and local growth rules depart from this value in a rule-dependent way [25]. For $\gamma = 3$, c_2 grows as $\ln T$.

Table I lists the measurements for the SR dual, its degree-preserving randomisation (SR*) and the edge-attachment (DM) model [6], each averaged over six realisations and 100 sources per realisation. The randomised graph sits on the configuration-model value, $\langle\mu\rangle/(c_2/e) = 1.00$–$0.90$. The SR dual exceeds it, by a factor that grows from 2.0 ± 0.3 at $T = 10^3$ to 6.4 ± 0.9 at $T = 10^5$, and the DM model stays within 0.87–1.20. Because μ has a heavy tail, $\langle\mu\rangle$ is dominated by rare pairs; the mean logarithm is better determined. The excess $\Delta\langle\ln\mu\rangle = \langle\ln\mu\rangle_{\rm SR} - \langle\ln\mu\rangle_{\rm SR^*}$, taken at the same degree sequence, grows from 0.40 to 1.03 over the same range.

To ask where the excess comes from, we varied the fraction of new nodes that close a triangle at fixed construction and measured $\Delta\langle\ln\mu\rangle$ against each network's own randomisation [$T = 3 \times 10^4$, six realisations, Fig. 5(b)]. In the topological surrogate, raising the corner-cut probability p_4 from 0.1 to 0.9 raises the triangle-closing fraction from 0.24 to 0.97 and lowers the excess monotonically, from 2.09 ± 0.17 to 0.19 ± 0.03. In the geometric model, the angle set $\{0, \pi/2\}$ closes no triangles and gives an excess of 2.87 ± 0.13; the set $\{0, \pi/3, 2\pi/3\}$ closes 0.23 and gives 2.43 ± 0.32; uniform angles close 0.49 and give 0.83 ± 0.07. Square-closing splits therefore supply the redundancy. The fraction of squares is not the only factor, however: selecting cells uniformly ($q = 0$) raises the triangle fraction to 0.62 but also raises the excess, to 1.50 ± 0.19. Uniform selection splits small cells as often as large ones and stacks squares locally, which the area weighting of the SR rule avoids. The DM model, which closes only triangles, has no excess.

TABLE I. Shortest-path statistics of the dual graph: mean distance $\langle d\rangle$, $\langle\mu\rangle/(c_2/e)$ and $\langle\ln\mu\rangle$, mean ± standard error over six realisations (100 sources each). Each entry gives SR / SR*, where SR* is the degree-preserving randomisation of the SR dual; standard errors of the SR* entries are below 0.03.

T	c_2	$\langle d\rangle$	$\langle\mu\rangle/(c_2/e)$	$\langle\ln\mu\rangle$
10^3	7.9	5.78/4.48	$2.0 \pm 0.3/1.00$	$1.09 \pm 0.06/0.70$
3×10^3	9.3	6.60/4.98	$2.9 \pm 0.5/1.00$	$1.39 \pm 0.07/0.80$
10^4	11.0	7.18/5.31	$3.3 \pm 0.4/0.94$	$1.55 \pm 0.06/0.87$
3×10^4	12.3	7.98/5.65	$5.4 \pm 1.0/0.93$	$1.77 \pm 0.07/0.94$
10^5	14.1	8.69/6.06	$6.4 \pm 0.9/0.90$	$2.03 \pm 0.06/1.00$

VII. WHAT MOVES THE DEGREE EXPONENT

Within mechanism B the exponent follows from two scalings: how the fitness of a ligament depends on its birth index, and how its hit rate depends on its fitness. If the hit rate is proportional to ℓ_i^a and $\ell_i \propto i^{-1/2}$, the total rate $\sum_j \ell_j^a$ grows as $t^{1-a/2}$ for $a < 2$, so $k_i \propto \ell_i^a t^{a/2} \propto (t/i)^{a/2}$, giving $\beta = a/2$ and $\gamma = 1 + 1/\beta = 1 + 2/a$. In two dimensions with uniform angles $a = 1$, so $\gamma = 3$. Any rule that keeps the hit rate proportional to length and the birth length proportional to the local cell size leaves γ at 3.

Exponents are estimated with the discrete maximum-likelihood approximation $\hat\gamma = 1 + n[\sum_i \ln(k_i/(k_{\rm min} - 1/2))]^{-1}$ over the n nodes with $k \geq k_{\rm min}$ [26], with standard error $(\hat\gamma - 1)/\sqrt{n}$. For a distribution with the curvature of Eq. (3) this estimator is biased low at small $k_{\rm min}$: on 10^5 samples drawn exactly from Eq. (3) it returns 2.77 ± 0.02, 2.87 ± 0.03, 2.91 ± 0.05 and 2.91 ± 0.10 for $k_{\rm min} = 6$, 10, 20 and 40. The SR model at $T = 10^5$ gives 2.77–2.79, 2.86–2.90, 2.94–2.97 and 2.90–3.03. Statements that $\gamma = 3$ therefore rest on comparison with these controls, not on the raw value of $\hat\gamma$.

We tested the design rules of Ref. [1] at $T = 3 \times 10^4$, two realisations each [Fig. 4(a)]:

- Thickness in the tunable-density phase ($\alpha = 0.5$–1, $\lambda_0 = 0.01$–0.1, $\rho = 0.12$–0.45) gives $\gamma_{\rm MLE} = 2.85$–2.92, $\beta = 0.50$–0.52, $C = 0.376$–0.378, and a triangle closure of 0.49–0.50.

- In the jammed phase ($\alpha = 0$–0.4), growth stops when the free area is exhausted (at $t \simeq 2000$ for $\alpha = 0$, $\lambda_0 = 0.02$). Jamming truncates T; it does not change γ (2.89–2.94 wherever the statistics allow) or C (0.35–0.37).

- Angle sets change clustering but not γ. With angles $\{0, \pi/2\}$ all cells are rectangles, so $C = 0$ exactly




FIG. 4. (a) Maximum-likelihood exponent ($k \geq 10$) for the design rules of Ref. [1], $T = 3 \times 10^4$, two realisations (the jammed $\alpha = 0$ case stops at $T \simeq 2000$; 3D plates at $T = 8 \times 10^4$, $k \geq 40$). The number on the right is the mean clustering coefficient. Error bars: half the difference between the two realisations (3D: estimator standard error). Dashed line: $\gamma = 3$; dotted line: $\gamma = 2.5$. The uniform SR control gives 2.87–2.91 at this size, the downward bias of the estimator discussed in Sec. VII. (b) Exponent against the fraction of ligaments whose angle follows a circular or a wave orientation field ($T = 3 \times 10^4$); the remaining ligaments have uniform angles.




FIG. 5. (a) Mean distance in the dual graph against T for the SR construction, its degree-preserving randomisation, the edge-attachment (DM) model [6] (six realisations) and the topological face surrogate with $p_4 = 0.29$ (two to three realisations). (b) Excess of the mean log-multiplicity over each network's degree-preserving randomisation against the fraction of new nodes that close a triangle, $T = 3 \times 10^4$, six realisations: SR rule variants (angle sets $\{0, \pi/2\}$, $\{0, \pi/3, 2\pi/3\}$, uniform angles, and uniform cell selection $q = 0$) and the face surrogate with $p_4 = 0.1$, 0.29, 0.6, 0.9. Error bars: standard errors.

and $\gamma = 2.87$; this is the rectangular fragmentation of Ref. [17]. With $\{0, \pi/3, 2\pi/3\}$, $C = 0.17$.

- Global alignment (von Mises, κ up to 16), cell selection proportional to area^q ($q = 0, 0.5, 2$), and linear or radial density gradients give $\gamma = 2.86$–2.96. Selection with $q = 0$ raises C to 0.45, and $q = 2$ lowers it to 0.35.

The angle set and the selection weight therefore tune clustering between 0 and 0.45, while the exponent stays at 3.

Coherent orientation fields. The polarized rule of Ref. [1], in which the angle follows a prescribed field $\theta(x, y)$, breaks the proportionality between hit rate and length. Locally parallel newcomers cannot hit each other; they end on the ligaments that are misaligned with the local field, and for straight ligaments in a curved field the misalignment grows with length. The degree then grows faster than length: fitting $\ln(k-2)$ against $\ln \ell$ for the oldest 5% of ligaments with $k > 4$ gives slopes $a = 1.43$–1.63, against 1.03–1.05 for uniform angles. With $a \simeq 1.5$ the argument above predicts $\beta \simeq 0.75$ and $\gamma \simeq 2.3$. It captures the direction of the change but overshoots it; the measured cohort exponents are smaller (see below), because the misaligned length of a ligament is not simply a power of its total length. For a circular field and for waves $\theta = \pi(x + y)/w$ with $w = 0.125$–1 we find $\gamma \simeq 2.44$–2.51 and cohort exponents $\beta \simeq 0.6$–0.67, stable from $T = 10^4$ to 2×10^5. At $T = 10^6$ the maximum-likelihood exponent is 2.43–2.50 ($k \geq 10$ to $k \geq 40$) for the circular field and 2.36–2.47 for the shortest wave ($w = 0.125$), while the uniform control gives 2.88–2.98, so the effect does not wash out at this size. The exponent decreases continuously as the fraction of ligaments following the field increases [Fig. 4(b)].

Three dimensions. For plates in a cube, a new plate meets $\bar{m}$ existing plates (the sides of its section polygon), creates $2+\bar{m}$ incidences and hands out $\bar{m}$ links. Then $b = \bar{m}/(2+\bar{m})$ and $\gamma = 2+2/\bar{m}$. We measure $\bar{m} = 4.47$–4.57, which predicts $\gamma = 2.44$. The dimensional argument $k \sim (\ell/\ell_T)^{d-1}$ gives $\gamma = (2d-1)/(d-1) = 2.5$, the exponent

of random Apollonian networks in d dimensions [27]. At $T = 8 \times 10^4$ the measured value is 2.50 ($k \geq 40$), with $\beta = 0.66$. Three-dimensional SR materials therefore have $\gamma \simeq 2.5$, not 3.

VIII. CONSEQUENCES FOR STRUCTURE–PROPERTY RELATIONS

At fixed α, the thickness of ligament i is $\lambda_0 i^{-\alpha}$ exactly, and its degree and length both scale as $i^{-1/2}$ on average [Eqs. (2) and (4)]. The marginal laws reported in Ref. [1], an energy that decays as $i^{-(3/4+\alpha/2)}$ with the introduction time and the corresponding powers of k, ℓ and λ, are one age law written in different variables. A product $U \sim k^a \lambda^b$ reproduces the age law for every α only if $a/2 + \alpha b = 3/4 + \alpha/2$, that is $a = 3/2$ and $b = 1/2$, the reported exponents. Because $k \propto \ell$ at fixed age [Eq. (4)], the same data are described equally well by $U \sim \ell^{3/2} \lambda^{1/2} f(\theta)$. Degree, length and age cannot be separated from cohort scaling alone.

Separating them requires the per-ligament energies themselves: a within-cohort regression of U on k, ℓ and λ would show whether degree adds anything beyond length and age. In a slender-beam frame model that we used as a check (Euler–Bernoulli beams, rigid T-junctions, $T \leq 5000$), fits on degree and on length were indistinguishable ($R^2 = 0.634$ and 0.631); a frame of slender beams is not the continuum finite-element model of Ref. [1], so we do not report its exponents.

The same reasoning applies to design. Thickness is a function of birth order, and degree and length are statistically slaved to birth order, so α and λ_0 act on the material through the age of each ligament on a fixed network. Connectivity becomes a design variable only when the rule is changed so that the hit rate stops being proportional to length, as with coherent orientation fields, or when the dimension changes the relation between fitness and birth order (Sec. VII).

IX. CONCLUSIONS

The scale-rich construction joins two known structures. Geometrically it is Cowan's recursive division of cells [9]; as a graph it grows by a face-local rule in which each new node joins two nodes on the boundary of one face. The recursion explains the cell hierarchy: uniform split fractions, a binary-search-tree genealogy and exponential areas. It also assigns each ligament its length at birth, $\ell_i \propto i^{-1/2}$, and that length acts as a fitness: the degree is a nearly Poisson count of hits with mean proportional to length times remaining time, without reinforcement. The k^{-3} tail, which preferential attachment on face incidences would also give, follows from that fitness, and the two mechanisms are separated by the fluctuations of the degree rather than by its exponent. The local rule explains the rest of the network: a hierarchical clustering spectrum from corner cuts and an excess of shortest paths from square-closing splits. Keeping both references in view settles which features are fixed by the class and which can be designed. The exponent $\gamma = 3$ and the equality $\langle n \rangle = 4$ are fixed, and in mean field so are the host-age law $h/t \to U^2$ and the stationarity of the thickness ratio. Clustering is designable through the angle set and the selection weight. The degree exponent is designable only through rules that make the hit rate grow faster than length.

Several points remain open. The rate $2/L(t)$ per unit length is a mean-field rate; the excess variance of the oldest ligaments shows that their local environment matters, and it raises the tail amplitude by about 4% above the Poisson fitness prediction. The per-hit triangle probability, which rises from 0.41 to 0.57 with the degree, and the non-local closure fraction 0.048 require the joint law of cell shapes next to long ligaments. Under coherent orientation fields we observe $\gamma \simeq 2.4$–2.5 up to $T = 10^6$; the fitness argument gives the direction of the change but not its value. Finally, the growth of the shortest-path excess with T is consistent with both a log-quadratic and a weak power law over the range studied.

More broadly, the SR dual shows that a spatial growth process can reproduce the degree statistics of a non-spatial growth class by a different route. Here the area-weighted point and the convexity of the cells deliver each new node to a face, as a random walk would, but the choice of sides within the face is set by length, a property fixed at birth, rather than by accumulated links.

ACKNOWLEDGMENTS

All code is available at https://github.com/av2atgh/scalerich. The calculations were assisted by Claude Opus 5, an AI assistant developed by Anthropic. Nodes & Links Ltd provided support in the form of salary for Alexei Vazquez but did not have any additional role in the conceptualization of the study, the analysis, the decision to publish or the preparation of the manuscript.

[1] C. Both, A. Y.-J. Chen, T.-T. Gao, N. Mooij, M. Charara, J. Wadhwa, C. M. Portela, and A.-L. Barabási, Scale-rich network-based metamaterials, Nature **659**, 345 (2026).

[2] M. Rosvall, A. Trusina, P. Minnhagen, and K. Sneppen, Networks and cities: An information perspective, Physical Review Letters **94**, 028701 (2005).

[3] V. Kalapala, V. Sanwalani, A. Clauset, and C. Moore, Scale invariance in road networks, Physical Review E **73**, 026130 (2006).

[4] A. Vazquez, Growing network with local rules: Preferential attachment, clustering hierarchy, and degree correlations, Physical Review E **67**, 056104 (2003).

[5] P. Holme and B. J. Kim, Growing scale-free networks with tunable clustering, Physical Review E **65**, 026107 (2002).

[6] S. N. Dorogovtsev, J. F. F. Mendes, and A. N. Samukhin, Size-dependent degree distribution of a scale-free growing network, Physical Review E **63**, 062101 (2001).

[7] E. Ravasz and A.-L. Barabási, Hierarchical organization in complex networks, Physical Review E **67**, 026112 (2003).

[8] G. Bianconi and A.-L. Barabási, Competition and multiscaling in evolving networks, Europhysics Letters **54**, 436 (2001).

[9] R. Cowan, New classes of random tessellations arising from iterative division of cells, Advances in Applied Probability **42**, 26 (2010).

[10] W. Nagel and V. Weiß, Crack STIT tessellations: characterization of stationary random tessellations stable with respect to iteration, Advances in Applied Probability **37**, 859 (2005).

[11] C. Redenbach and T. Jung, Random tessellations – an overview of models, arXiv (2025), 2501.05166.

[12] S. Bohn, L. Pauchard, and Y. Couder, Hierarchical crack pattern as formed by successive domain divisions. I. Temporal and geometrical hierarchy, Physical Review E **71**, 046214 (2005).

[13] S. Bohn, J. Platkiewicz, B. Andreotti, M. Adda-Bedia, and Y. Couder, Hierarchical crack pattern as formed by successive domain divisions. II. From disordered to deterministic behavior, Physical Review E **71**, 046215 (2005).

[14] E. N. Gilbert, Random plane networks and needle-shaped crystals, in *Applications of Undergraduate Mathematics in Engineering*, edited by B. Noble (Macmillan, New York, 1967).

[15] M. C. Gibson, A. B. Patel, R. Nagpal, and N. Perrimon, The emergence of geometric order in proliferating metazoan epithelia, Nature **442**, 1038 (2006).

[16] L. Devroye, A note on the height of binary search trees, Journal of the ACM **33**, 489 (1986).

[17] P. L. Krapivsky and E. Ben-Naim, Scaling and multiscaling in models of fragmentation, Physical Review E **50**, 3502 (1994).

[18] A.-L. Barabási and R. Albert, Emergence of scaling in random networks, Science **286**, 509 (1999).

[19] P. L. Krapivsky and S. Redner, Organization of growing random networks, Physical Review E **63**, 066123 (2001).

[20] S. N. Dorogovtsev and J. F. F. Mendes, Evolution of networks, Advances in Physics **51**, 1079 (2002).

[21] L. A. Santaló, *Integral Geometry and Geometric Probability* (Addison-Wesley, Reading, MA, 1976).

[22] S. N. Dorogovtsev, A. V. Goltsev, and J. F. F. Mendes, Pseudofractal scale-free web, Physical Review E **65**, 066122 (2002).

[23] J. S. Andrade, H. J. Herrmann, R. F. S. Andrade, and L. R. da Silva, Apollonian networks: Simultaneously scale-free, small world, euclidean, space filling, and with matching graphs, Physical Review Letters **94**, 018702 (2005).

[24] T. Zhou, G. Yan, and B.-H. Wang, Maximal planar networks with large clustering coefficient and power-law degree distribution, Physical Review E **71**, 046141 (2005).

[25] A. Vazquez, Local network evolution rules driving shortest path multiplicity, Physical Review E 10.1103/t1sg-qhhf (2026), arXiv:2605.25237 [physics.soc-ph].

[26] A. Clauset, C. R. Shalizi, and M. E. J. Newman, Power-law distributions in empirical data, SIAM Review **51**, 661 (2009).

[27] I. Kolossváry, J. Komjáthy, and L. Vágó, Degrees and distances in random and evolving Apollonian networks, Advances in Applied Probability **48**, 865 (2016).