

# Lecture 8. Random variable

Alex Avdiushenko

Neapolis University Paphos

June 28, 2023



# Recap and problems

# Recap and problems

- ▶ Still you can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).

# Recap and problems

- ▶ Still you can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).
- ▶ We need to have pair  $(\Omega, P)$ , where  $\Omega$  is all possible answers and  $P$  is probability measure

# Recap and problems

- ▶ Still you can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).
- ▶ We need to have pair  $(\Omega, P)$ , where  $\Omega$  is all possible answers and  $P$  is probability measure
- ▶ But you may not want to work with all the  $\Omega$

# Recap and problems

- ▶ Still you can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).
- ▶ We need to have pair  $(\Omega, P)$ , where  $\Omega$  is all possible answers and  $P$  is probability measure
- ▶ But you may not want to work with all the  $\Omega$
- ▶ May be  $\Omega$  is too informative, of complicated, or huge

# Recap and problems

- ▶ Still you can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).
- ▶ We need to have pair  $(\Omega, P)$ , where  $\Omega$  is all possible answers and  $P$  is probability measure
- ▶ But you may not want to work with all the  $\Omega$
- ▶ May be  $\Omega$  is too informative, of complicated, or huge
- ▶ Instead we can ask **black box** for some specific characteristics as weight or temperature

# Definition of a Random Variable

A **random variable** is a function that assigns a real number to each outcome in the sample space of a random experiment. Formally, if  $\Omega$  is the sample space of a random experiment, a random variable  $\xi$  is a function:

$$\xi : \Omega \rightarrow \mathbb{R}$$

In essence, a random variable provides a way to map the outcomes of a random process to numerical quantities.



# Examples

1. For rolling a die  $\Omega = \{1, 2, 3, 4, 5, 6\}$  random variable  $\xi(k) = k \bmod 2$  shows us even results
2. For  $k$  fair coins  $\Omega = \{(a_1, \dots, a_k) \mid a_i \in \{0, 1\}\}$  we can count Heads:  $\xi((a_1, \dots, a_k)) = a_1 + \dots + a_k$

So any random variable  $\xi(\omega)$  is **new black box** with real numbers as samples and pair  $(\mathbb{R}, P_\xi)$ , where

$$P_\xi(A) = P(\xi \in A) = P(\{\omega \mid \xi(\omega) \in A\}) = P(\xi \text{ shows value from } A)$$

$$\xi^{-1}(A) = \{\omega \mid \xi(\omega) \in A\}$$

So we want to study different measures on  $\mathbb{R}$ .

So we want to study different measures on  $\mathbb{R}$ .

### Question

What probability measures can be on  $\mathbb{R}$  at all?

So we want to study different measures on  $\mathbb{R}$ .

### Question

What probability measures can be on  $\mathbb{R}$  at all?

Recall, that  $P : 2^{\mathbb{R}} \rightarrow [0, 1]$  and for any event  $A \mapsto P(A)$

So we want to study different measures on  $\mathbb{R}$ .

### Question

What probability measures can be on  $\mathbb{R}$  at all?

Recall, that  $P : 2^{\mathbb{R}} \rightarrow [0, 1]$  and for any event  $A \mapsto P(A)$

►  $P(\emptyset) = 0, P(\mathbb{R}) = 1$

So we want to study different measures on  $\mathbb{R}$ .

### Question

What probability measures can be on  $\mathbb{R}$  at all?

Recall, that  $P: 2^{\mathbb{R}} \rightarrow [0, 1]$  and for any event  $A \mapsto P(A)$

- ▶  $P(\emptyset) = 0, P(\mathbb{R}) = 1$
- ▶  $A_1, A_2, \dots, A_n, \dots \subseteq \mathbb{R}, A_i \cap A_j = \emptyset$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

So we want to study different measures on  $\mathbb{R}$ .

### Question

What probability measures can be on  $\mathbb{R}$  at all?

Recall, that  $P: 2^{\mathbb{R}} \rightarrow [0,1]$  and for any event  $A \mapsto P(A)$

- ▶  $P(\emptyset) = 0, P(\mathbb{R}) = 1$
- ▶  $A_1, A_2, \dots, A_n, \dots \subseteq \mathbb{R}, A_i \cap A_j = \emptyset$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

It can be proven mathematically, that sufficiently to know only  $P((a,b])$  for all  $a, b \in \mathbb{R} \Rightarrow$  there is general procedure, which builds measure  $P(A), A \subseteq \mathbb{R}$  (it is called construction of the outer Lebesgue measure, and of course here goes  $\sigma$ -algebras and so on)

# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$



# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$

## Definition

Cumulative Distribution Function (CDF) of  $\xi$  is

$$F_{\xi}(x) = P_{\xi}((-\infty, x]) = P(\xi \leq x)$$

# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$

## Definition

Cumulative Distribution Function (CDF) of  $\xi$  is

$$F_{\xi}(x) = P_{\xi}((-\infty, x]) = P(\xi \leq x)$$

Vice versa some  $F : \mathbb{R} \rightarrow \mathbb{R}$  is distribution function iff

# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$

## Definition

Cumulative Distribution Function (CDF) of  $\xi$  is

$$F_{\xi}(x) = P_{\xi}((-\infty, x]) = P(\xi \leq x)$$

Vice versa some  $F : \mathbb{R} \rightarrow \mathbb{R}$  is distribution function iff

1.  $x \leq y \Rightarrow F(x) \leq F(y)$

# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$

## Definition

Cumulative Distribution Function (CDF) of  $\xi$  is

$$F_{\xi}(x) = P_{\xi}((-\infty, x]) = P(\xi \leq x)$$

Vice versa some  $F : \mathbb{R} \rightarrow \mathbb{R}$  is distribution function iff

1.  $x \leq y \Rightarrow F(x) \leq F(y)$
2.  $\lim_{x \rightarrow -\infty} F(x) = 0, \lim_{x \rightarrow \infty} F(x) = 1$

# Cumulative Distribution Function

Actually we need to know only  $P((-\infty, x])$  as

$$P((a, b]) = P((-\infty, b]) - P((-\infty, a])$$

## Definition

Cumulative Distribution Function (CDF) of  $\xi$  is

$$F_{\xi}(x) = P_{\xi}((-\infty, x]) = P(\xi \leq x)$$

Vice versa some  $F : \mathbb{R} \rightarrow \mathbb{R}$  is distribution function iff

1.  $x \leq y \Rightarrow F(x) \leq F(y)$
2.  $\lim_{x \rightarrow -\infty} F(x) = 0, \lim_{x \rightarrow \infty} F(x) = 1$
3.  $\lim_{t \rightarrow 0, t > 0} F(x + t) = F(x)$

# Examples



1. Consider rolling a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$ , random variable  $\xi(k) = k \bmod 2$  shows us even results

# Examples

1. Consider rolling a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$ , random variable  $\xi(k) = k \bmod 2$  shows us even results
2.  $\Omega = [0, 1]$ ,  $A \subseteq [0, 1]$ ,  $P(A) = \int_A 1 dx = \text{length of } A$


# Examples

1. Consider rolling a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$ , random variable  $\xi(k) = k \bmod 2$  shows us even results
2.  $\Omega = [0, 1]$ ,  $A \subseteq [0, 1]$ ,  $P(A) = \int_A 1 dx = \text{length of } A$






# Link between plots of $\xi(\omega)$ and $F_\xi(x)$




# Link between plots of $\xi(\omega)$ and $F_\xi(x)$



So the vertical  $\xi(\omega)$ -axis is x-axis  
for CDF  $F_\xi(x)$


# Link between plots of $\xi(\omega)$ and $F_\xi(x)$



So the vertical  $\xi(\omega)$ -axis is x-axis for CDF  $F_\xi(x)$

Moreover, different random variables  $\xi$  may correspond to one and the same CDF. For instance, you could reverse the  $\Omega$ -axis.

# Link between plots of $\xi(\omega)$ and $F_\xi(x)$




So the vertical  $\xi(\omega)$ -axis is x-axis for CDF  $F_\xi(x)$

Moreover, different random variables  $\xi$  may correspond to one and the same CDF. For instance, you could reverse the  $\Omega$ -axis.


## Question

How to find CDF break points on this plot?

# Probability atoms




# Probability atoms



$$P(\{c\}) = ?, \quad P(1) = ?$$


# Probability atoms



$$P(\{c\}) = ?, \quad P(1) = ?$$

$$\begin{aligned} P(\{c\}) &= \lim_{\delta \rightarrow 0} P((c - \delta, c + \delta]) = \\ &= \lim_{\delta \rightarrow 0} (F(c + \delta) - F(c - \delta)) = 0 \end{aligned}$$

# Probability atoms



$$P(\{c\}) = ?, \quad P(1) = ?$$

$$\begin{aligned} P(\{c\}) &= \lim_{\delta \rightarrow 0} P((c - \delta, c + \delta]) = \\ &= \lim_{\delta \rightarrow 0} (F(c + \delta) - F(c - \delta)) = 0 \end{aligned}$$

$$P(1) = F(1) - F(1-) =$$

size of the jump discontinuity



# Classification of measures

Mathematics	Engineers
Discrete	Discrete
Continuous	–
Absolutely Continuous	Continuous

# Classification of measures

<b>Mathematics</b>	<b>Engineers</b>
Discrete	Discrete
Continuous	–
Absolutely Continuous	Continuous

We are using the engineers terminology.

# Classification of measures

Mathematics	Engineers
Discrete	Discrete
Continuous	–
Absolutely Continuous	Continuous

We are using the engineers terminology.

1. Discrete consists of atoms  $a_i$  only:  $\sum_{i=1}^{\infty} p_i = 1$

# Classification of measures

Mathematics	Engineers
Discrete	Discrete
Continuous	–
Absolutely Continuous	Continuous

We are using the engineers terminology.

1. Discrete consists of atoms  $a_i$  only:  $\sum_{i=1}^{\infty} p_i = 1$
2. Continuous is defined by its density function  $p(x) > 0$ :

$$\int_{\mathbb{R}} p(x) dx = 1, \quad F'(x) = p(x), \quad F(x) = \int_{-\infty}^x p(t) dt$$

# Cantor Function

The *Cantor function*, also known as the *Devil's Staircase*, is a peculiar function that was discovered by Georg Cantor.

# Cantor Function

The *Cantor function*, also known as the *Devil's Staircase*, is a peculiar function that was discovered by Georg Cantor.

- ▶ It is defined on the interval  $[0,1]$

# Cantor Function

The *Cantor function*, also known as the *Devil's Staircase*, is a peculiar function that was discovered by Georg Cantor.

- ▶ It is defined on the interval  $[0,1]$
- ▶ It is constant on the set of numbers that are not in the Cantor set, which has Lebesgue measure zero

# Cantor Function

The *Cantor function*, also known as the *Devil's Staircase*, is a peculiar function that was discovered by Georg Cantor.

- ▶ It is defined on the interval  $[0,1]$
- ▶ It is constant on the set of numbers that are not in the Cantor set, which has Lebesgue measure zero
- ▶ It is increasing: if  $0 \leq x < y \leq 1$ , then  $f(x) \leq f(y)$



# Cantor Function


The *Cantor function*, also known as the *Devil's Staircase*, is a peculiar function that was discovered by Georg Cantor.

- ▶ It is defined on the interval  $[0,1]$
- ▶ It is constant on the set of numbers that are not in the Cantor set, which has Lebesgue measure zero
- ▶ It is increasing: if  $0 \leq x < y \leq 1$ , then  $f(x) \leq f(y)$
- ▶ **It is continuous, but not absolutely continuous**

Mathematically, it can be defined as:

$$f(x) = \frac{1}{2} \sum_{n=0}^{\infty} \frac{2}{3^n} \chi_{C_n}(x)$$

where  $C_n$  is the  $n$ -th stage in the construction of the Cantor set, and  $\chi$  is the characteristic function.



Source: Wikipedia, the free encyclopedia

# Histogramm or how to understand density

A *histogram* is a graphical representation of the distribution of a dataset.

# Histogramm or how to understand density

A *histogram* is a graphical representation of the distribution of a dataset.

- ▶ It is an estimate of the probability distribution of a continuous variable

# Histogramm or how to understand density

A *histogram* is a graphical representation of the distribution of a dataset.


- ▶ It is an estimate of the probability distribution of a continuous variable
- ▶ To construct a histogram, the first step is to "bin" the range of values — that is, divide the entire range of values into a series of intervals

# Histogramm or how to understand density

A *histogram* is a graphical representation of the distribution of a dataset.

- ▶ It is an estimate of the probability distribution of a continuous variable
- ▶ To construct a histogram, the first step is to "bin" the range of values — that is, divide the entire range of values into a series of intervals
- ▶ Then count how many values fall into each interval

An example of a histogram:



# Mathematical Expectation

We like numbers more than “strange” random variables.



# Mathematical Expectation

We like numbers more than “strange” random variables.

## Definition

For a discrete random variable  $\xi$  with probability mass function  $p(x)$ , the mathematical expectation  $E\xi$  is defined as:

$$E\xi = \sum_x x \cdot p(x)$$

For a continuous random variable  $\xi$  with probability density function  $p(x)$ , the mathematical expectation  $E\xi$  is defined as:

$$E\xi = \int_{-\infty}^{\infty} x \cdot p(x) dx = \int_{\mathbb{R}} x dP_{\xi} = \int_{\Omega} \xi dP$$

# Mathematical Expectation

We like numbers more than “strange” random variables.

## Definition

For a discrete random variable  $\xi$  with probability mass function  $p(x)$ , the mathematical expectation  $E\xi$  is defined as:

$$E\xi = \sum_x x \cdot p(x)$$

For a continuous random variable  $\xi$  with probability density function  $p(x)$ , the mathematical expectation  $E\xi$  is defined as:

$$E\xi = \int_{-\infty}^{\infty} x \cdot p(x) dx = \int_{\mathbb{R}} x dP_{\xi} = \int_{\Omega} \xi dP$$

The mathematical expectation can be thought of as the “average” or “mean” value of the random variable.

# First example


$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = a_1 + \dots + a_n$$

# First example

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = a_1 + \dots + a_n$$



## Second example

$\Omega$  is unit square,  $P(A)$  is area of  $A$

$$\xi(x, y) = |x - y|$$


## Second example

$\Omega$  is unit square,  $P(A)$  is area of  $A$


$$\xi(x, y) = |x - y|$$

$$E\xi = ?$$

Plot of  $\xi(x, y) = |x - y|$



Plot of  $\xi(x, y) = |x - y|$



$$E\xi = \int_{\square} |x - y| dx dy$$



# Three Situations for Calculating Expectation

$$E\xi = \int_{\mathbb{R}} x dP_{\xi}$$

We need to know  $P_{\xi}$

# Three Situations for Calculating Expectation

$$E\xi = \int_{\mathbb{R}} x dP_{\xi}$$

We need to know  $P_{\xi}$

If  $F_{\xi}(x)$  is CDF:

$$E\xi = \int_{\mathbb{R}} x dF_{\xi}(x) = \sum_{i=1}^{\infty} a_i p_i + \int_{\mathbb{R}} x F'_{\xi}(x) dx,$$

where  $p_i = F_{\xi}(a_i) - F_{\xi}(a_i -)$  for the jumps discontinuity  $a_i$

# Three Situations for Calculating Expectation

$$E\xi = \int_{\mathbb{R}} x dP_{\xi}$$

We need to know  $P_{\xi}$

If  $F_{\xi}(x)$  is CDF:

$$E\xi = \int_{\mathbb{R}} x dF_{\xi}(x) = \sum_{i=1}^{\infty} a_i p_i + \int_{\mathbb{R}} x F'_{\xi}(x) dx,$$

where  $p_i = F_{\xi}(a_i) - F_{\xi}(a_i -)$  for the jumps discontinuity  $a_i$

► If  $\xi$  is discrete then  $E\xi = \sum_{i=1}^{\infty} a_i p_i$

# Three Situations for Calculating Expectation

$$E\xi = \int_{\mathbb{R}} x dP_{\xi}$$

We need to know  $P_{\xi}$

If  $F_{\xi}(x)$  is CDF:

$$E\xi = \int_{\mathbb{R}} x dF_{\xi}(x) = \sum_{i=1}^{\infty} a_i p_i + \int_{\mathbb{R}} x F'_{\xi}(x) dx,$$

where  $p_i = F_{\xi}(a_i) - F_{\xi}(a_i -)$  for the jumps discontinuity  $a_i$

- ▶ If  $\xi$  is discrete then  $E\xi = \sum_{i=1}^{\infty} a_i p_i$
- ▶ If  $\xi$  is continuous then  $E\xi = \int_{\mathbb{R}} x p(x) dx$

# Three Situations for Calculating Expectation

$$E\xi = \int_{\mathbb{R}} x dP_{\xi}$$

We need to know  $P_{\xi}$

If  $F_{\xi}(x)$  is CDF:

$$E\xi = \int_{\mathbb{R}} x dF_{\xi}(x) = \sum_{i=1}^{\infty} a_i p_i + \int_{\mathbb{R}} x F'_{\xi}(x) dx,$$

where  $p_i = F_{\xi}(a_i) - F_{\xi}(a_i -)$  for the jumps discontinuity  $a_i$

- ▶ If  $\xi$  is discrete then  $E\xi = \sum_{i=1}^{\infty} a_i p_i$
- ▶ If  $\xi$  is continuous then  $E\xi = \int_{\mathbb{R}} x p(x) dx$
- ▶  $Ef(\xi) = \sum_{i=1}^{\infty} f(a_i) p_i + \int_{\mathbb{R}} f(x) F'_{\xi}(x) dx$

# Expectation Properties

Let  $\xi$  and  $\eta$  be random variables, and  $a$  and  $b$  be constants, then the following properties hold:

1.  $E[a] = a$ , where  $a$  is a constant
2. **Linearity of Expectation:**  $E[a\xi + b\eta] = aE[\xi] + bE[\eta]$

# Expectation Properties

Let  $\xi$  and  $\eta$  be random variables, and  $a$  and  $b$  be constants, then the following properties hold:

1.  $E[a] = a$ , where  $a$  is a constant
2. **Linearity of Expectation:**  $E[a\xi + b\eta] = aE[\xi] + bE[\eta]$
3.  $E[\xi - E\xi] = 0$

# Expectation Properties

Let  $\xi$  and  $\eta$  be random variables, and  $a$  and  $b$  be constants, then the following properties hold:

1.  $E[a] = a$ , where  $a$  is a constant
2. **Linearity of Expectation:**  $E[a\xi + b\eta] = aE[\xi] + bE[\eta]$
3.  $E[\xi - E\xi] = 0$
4. If  $\xi \geq 0$  a.s. then  $E[\xi] \geq 0$
5. If  $\xi \leq \eta$  a.s. then  $E[\xi] \leq E[\eta]$
6. (Jensen's inequality)  
If  $\varphi$  is a convex function, then  $E[\varphi(\xi)] \geq \varphi(E[\xi])$



# Expectation Properties

Let  $\xi$  and  $\eta$  be random variables, and  $a$  and  $b$  be constants, then the following properties hold:

1.  $E[a] = a$ , where  $a$  is a constant
2. **Linearity of Expectation:**  $E[a\xi + b\eta] = aE[\xi] + bE[\eta]$
3.  $E[\xi - E\xi] = 0$
4. If  $\xi \geq 0$  a.s. then  $E[\xi] \geq 0$
5. If  $\xi \leq \eta$  a.s. then  $E[\xi] \leq E[\eta]$
6. (Jensen's inequality)  
If  $\varphi$  is a convex function, then  $E[\varphi(\xi)] \geq \varphi(E[\xi])$
7. (in the future) If  $\xi$  and  $\eta$  are independent, then

$$E[\xi\eta] = E[\xi]E[\eta]$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

Better to invent new random variables:

$$\xi_{ij}(a_1, \dots, a_n) = \begin{cases} 1, & a_i a_j = 01 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{So } E\xi =$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

Better to invent new random variables:

$$\xi_{ij}(a_1, \dots, a_n) = \begin{cases} 1, & a_i a_j = 01 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{So } E\xi = E \sum_i \xi_{ii+1} = \sum_i E\xi_{ii+1} =$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

Better to invent new random variables:

$$\xi_{ij}(a_1, \dots, a_n) = \begin{cases} 1, & a_i a_j = 01 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{So } E\xi = E \sum_i \xi_{ii+1} = \sum_i E\xi_{ii+1} = \sum_i P(\xi_{ii+1} = 1) =$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

Better to invent new random variables:

$$\xi_{ij}(a_1, \dots, a_n) = \begin{cases} 1, & a_i a_j = 01 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{So } E\xi = E \sum_i \xi_{ii+1} = \sum_i E\xi_{ii+1} = \sum_i P(\xi_{ii+1} = 1) = \sum_i \frac{1}{4} =$$

# Task 1

$$\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}, \quad P((a_1, \dots, a_n)) = \frac{1}{2^n}$$

$$\xi(a_1, \dots, a_n) = \#(\text{ of occurrences of } 01)$$

$$E\xi = ?$$

Of course we can calculate  $E\xi = \sum_{k=0}^{n-1} k \cdot P(k \text{ occurrences})$ , but it's hard.

Better to invent new random variables:

$$\xi_{ij}(a_1, \dots, a_n) = \begin{cases} 1, & a_i a_j = 01 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{So } E\xi = E \sum_i \xi_{ii+1} = \sum_i E\xi_{ii+1} = \sum_i P(\xi_{ii+1} = 1) = \sum_i \frac{1}{4} = \frac{n-1}{4}$$



## Task 2

$\Omega = S_n$ , choose a random permutation uniformly

Number is stable, if  $k = \sigma(k)$ . Random variable  $\xi : \Omega \rightarrow \mathbb{R}$ , find  $E\xi$ :

$$\xi(\sigma) = \#(\text{stable numbers})$$

## Task 2

$\Omega = S_n$ , choose a random permutation uniformly

Number is stable, if  $k = \sigma(k)$ . Random variable  $\xi : \Omega \rightarrow \mathbb{R}$ , find  $E\xi$ :

$$\xi(\sigma) = \#(\text{stable numbers})$$

### Solution

Again consider new random variables:

$$\xi_k(\sigma) = \begin{cases} 1, & \text{if } \sigma(k) = k \\ 0, & \text{otherwise} \end{cases}$$

## Task 2

$\Omega = S_n$ , choose a random permutation uniformly

Number is stable, if  $k = \sigma(k)$ . Random variable  $\xi : \Omega \rightarrow \mathbb{R}$ , find  $E\xi$ :

$$\xi(\sigma) = \#(\text{stable numbers})$$

### Solution

Again consider new random variables:

$$\xi_k(\sigma) = \begin{cases} 1, & \text{if } \sigma(k) = k \\ 0, & \text{otherwise} \end{cases}$$

Then

$$E\xi = E\xi_1 + \dots + E\xi_n = \sum P(\xi_k = 1) = n \cdot \frac{(n-1)!}{n!} = 1$$

# How far the answers are scattered from expectation?

The variance of a random variable  $\xi$  is a measure of its dispersion. It is denoted as  $Var(\xi)$  or  $\sigma^2$  and it is defined as:

$$Var(\xi) = E[(\xi - E\xi)^2]$$

Alternatively, it can be computed using the formula:

$$Var(\xi) = E[\xi^2] - (E\xi)^2$$

The standard deviation of a random variable is the square root of its variance and it is denoted by  $\sigma$ .

# How far the answers are scattered from expectation?


The variance of a random variable  $\xi$  is a measure of its dispersion. It is denoted as  $Var(\xi)$  or  $\sigma^2$  and it is defined as:

$$Var(\xi) = E[(\xi - E\xi)^2]$$

Alternatively, it can be computed using the formula:

$$Var(\xi) = E[\xi^2] - (E\xi)^2$$

The standard deviation of a random variable is the square root of its variance and it is denoted by  $\sigma$ .



$Var(a\xi + b) = a^2 Var(\xi)$  – it can be proven by definition

# Gaussian Distribution

The Gaussian distribution, also known as the normal distribution, is a continuous probability distribution for a real-valued random variable. Its probability density function is given by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where:

# Gaussian Distribution

The Gaussian distribution, also known as the normal distribution, is a continuous probability distribution for a real-valued random variable. Its probability density function is given by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where:

- ▶  $\mu$  is the mean or expectation of the distribution (and also its median and mode)

# Gaussian Distribution

The Gaussian distribution, also known as the normal distribution, is a continuous probability distribution for a real-valued random variable. Its probability density function is given by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where:

- ▶  $\mu$  is the mean or expectation of the distribution (and also its median and mode)
- ▶  $\sigma$  is the standard deviation



# Gaussian Distribution

The Gaussian distribution, also known as the normal distribution, is a continuous probability distribution for a real-valued random variable. Its probability density function is given by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where:

- ▶  $\mu$  is the mean or expectation of the distribution (and also its median and mode)
- ▶  $\sigma$  is the standard deviation
- ▶  $\sigma^2$  is the variance

# Gaussian Distribution

The Gaussian distribution, also known as the normal distribution, is a continuous probability distribution for a real-valued random variable. Its probability density function is given by:


$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

where:

- ▶  $\mu$  is the mean or expectation of the distribution (and also its median and mode)
- ▶  $\sigma$  is the standard deviation
- ▶  $\sigma^2$  is the variance

The standard Gaussian distribution has a mean of 0 and a standard deviation of 1. It is denoted as  $\xi \sim N(\mu, \sigma^2)$

# Gaussian Distribution Plot



where  $\xi \sim N(\mu = 0, \sigma^2 = 1)$  and:

- ▶  $x$  is a normal random variable
- ▶  $f(x)$  is the probability density function

# What if we have several random variables?

$(\Omega, P)$  and  $\xi_1, \xi_2, \dots, \xi_k$

## Question

How can we describe all the values?

# What if we have several random variables?

$(\Omega, P)$  and  $\xi_1, \xi_2, \dots, \xi_k$

## Question

How can we describe all the values?

$\vec{\xi} : \Omega \rightarrow \mathbb{R}^k, \quad \omega \mapsto (\xi_1(\omega), \xi_2(\omega), \dots, \xi_k(\omega))$

So we have pair  $(\mathbb{R}^k, P_{\vec{\xi}})$  and  $P_{\vec{\xi}}(A) = P\left(\left\{\omega \mid \vec{\xi}(\omega) \in A\right\}\right)$  —  
probability distribution of random vector

# What if we have several random variables?

$(\Omega, P)$  and  $\xi_1, \xi_2, \dots, \xi_k$

## Question

How can we describe all the values?

$\vec{\xi}: \Omega \rightarrow \mathbb{R}^k, \quad \omega \mapsto (\xi_1(\omega), \xi_2(\omega), \dots, \xi_k(\omega))$


So we have pair  $(\mathbb{R}^k, P_{\vec{\xi}})$  and  $P_{\vec{\xi}}(A) = P\left(\left\{\omega \mid \vec{\xi}(\omega) \in A\right\}\right)$  — probability distribution of random vector

How to define probability  $P$  in  $\mathbb{R}^k$ ? Again we need “good” sets, and for instance in  $\mathbb{R}^2$  it is  $\{(x, y) \mid x \leq a, y \leq b\}$ . So we need CDF as

$$F(x_1, \dots, x_k) = P((-\infty, x_1] \times \dots \times (-\infty, x_k])$$


# Task 3

Given  $F$  in  $\mathbb{R}^2$  compute  $P((a, b] \times (c, d])$ .



# Task 3

Given  $F$  in  $\mathbb{R}^2$  compute  $P((a, b] \times (c, d])$ .




1. As in 1D we use limit to define the probability of rectangle



# Task 3


Given  $F$  in  $\mathbb{R}^2$  compute  $P((a, b] \times (c, d])$ .



1. As in 1D we use limit to define the probability of rectangle
2. There is discrete and continuous random variables

# Task 3

Given  $F$  in  $\mathbb{R}^2$  compute  $P((a, b] \times (c, d])$ .



1. As in 1D we use limit to define the probability of rectangle
2. There is discrete and continuous random variables
3.  $P_{\xi}(A) = \int_A p(x_1, \dots, x_n) dx_1 \dots dx_n$