

Lecture 7. Probability spaces

Alex Avdiushenko

Neapolis University Paphos

June 21, 2023



Intro to the philosophy of probability theory

- ▶ The first part of the probability is some kind of mental model on how to properly **understand and apply** it
- ▶ The second part is a complicated mathematical formalism that is needed to make the theory consistent, filtering some “bad” sets and functions (here is σ -algebras, measurable functions, Lebesgue integrals etc.)

Intro to the philosophy of probability theory

- ▶ The first part of the probability is some kind of mental model on how to properly **understand and apply** it
- ▶ The second part is a complicated mathematical formalism that is needed to make the theory consistent, filtering some “bad” sets and functions (here is σ -algebras, measurable functions, Lebesgue integrals etc.)
- ▶ In real life and applications, it seems that there are no non-measurable sets and all these theoretical contradictions at all
- ▶ Therefore, we can build the entire theory of probability not completely rigorously, but cutting mathematical corners and getting the intuition necessary to use probability

Mental model of random variable

You can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).

1. More formally, a random variable is a function X from a sample space S to the set of elementary outcomes Ω . This function assigns to each element s in the sample space a unique value $X(s)$.

Mental model of random variable

You can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).

1. More formally, a random variable is a function X from a sample space S to the set of elementary outcomes Ω . This function assigns to each element s in the sample space a unique value $X(s)$.

The two main types of random variables are:

Mental model of random variable

You can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).

1. More formally, a random variable is a function X from a sample space S to the set of elementary outcomes Ω . This function assigns to each element s in the sample space a unique value $X(s)$.

The two main types of random variables are:

- ▶ **Discrete Random Variables:** These are variables that take on a countable number of distinct values. Examples include flipping a coin, rolling a die, etc.

Mental model of random variable

You can think of random variable as a variable whose possible values are some outcomes of a random phenomenon (some **black box**, answering your questions).

1. More formally, a random variable is a function X from a sample space S to the set of elementary outcomes Ω . This function assigns to each element s in the sample space a unique value $X(s)$.

The two main types of random variables are:

- ▶ **Discrete Random Variables:** These are variables that take on a countable number of distinct values. Examples include flipping a coin, rolling a die, etc.
- ▶ **Continuous Random Variables:** These are variables that can take on any value in a given range or interval. Examples include the height of a person, the weight of an object, etc.

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P: 2^\Omega \rightarrow [0, 1]$

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P: 2^\Omega \rightarrow [0, 1]$
For any event $A \mapsto P(A)$

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P: 2^\Omega \rightarrow [0, 1]$
For any event $A \mapsto P(A)$
 - ▶ $P(\emptyset) = 0, P(\Omega) = 1$

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P: 2^\Omega \rightarrow [0, 1]$
For any event $A \mapsto P(A)$
 - ▶ $P(\emptyset) = 0, P(\Omega) = 1$
 - ▶ $A \cap B = \emptyset \Rightarrow P(A \cup B) = P(A) + P(B)$

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P: 2^\Omega \rightarrow [0, 1]$

For any event $A \mapsto P(A)$

- ▶ $P(\emptyset) = 0, P(\Omega) = 1$
- ▶ $A \cap B = \emptyset \Rightarrow P(A \cup B) = P(A) + P(B)$
- ▶ (as above plus some continuity)
 $A_1, A_2, \dots, A_n, \dots \subseteq \Omega, A_i \cap A_j = \emptyset$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

2. Event is some subset of the set of elementary outcomes Ω
3. Probability (probability measure) is function $P : 2^\Omega \rightarrow [0, 1]$

For any event $A \mapsto P(A)$

- ▶ $P(\emptyset) = 0, P(\Omega) = 1$
- ▶ $A \cap B = \emptyset \Rightarrow P(A \cup B) = P(A) + P(B)$
- ▶ (as above plus some continuity)
 $A_1, A_2, \dots, A_n, \dots \subseteq \Omega, A_i \cap A_j = \emptyset$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

Technical problem

If Ω is more than countable, then there is no such probability measures ;)

So we will ignore this technical and mathematical problem and we will work with pair (Ω, P) instead of triple (Ω, Σ, P) as our Probability Space

Remarks

The probability distribution of a random variable describes how the probabilities are distributed over the values of the random variable.

Remarks

The probability distribution of a random variable describes how the probabilities are distributed over the values of the random variable.

Question

What does it mean for events, that $A \subseteq B$?

The probability distribution of a random variable describes how the probabilities are distributed over the values of the random variable.

Question

What does it mean for events, that $A \subseteq B$?

It means, that $A \Rightarrow B$, so $P(A) \leq P(B)$ and $P(B \setminus A) = P(B) - P(A)$
For $A, B \subseteq \Omega$ in general case $P(B \setminus A) = P(B) - P(A \cap B)$

The probability distribution of a random variable describes how the probabilities are distributed over the values of the random variable.

Question

What does it mean for events, that $A \subseteq B$?

It means, that $A \Rightarrow B$, so $P(A) \leq P(B)$ and $P(B \setminus A) = P(B) - P(A)$
For $A, B \subseteq \Omega$ in general case $P(B \setminus A) = P(B) - P(A \cap B)$

Also there is negation $\bar{A} = \Omega \setminus A$ and $P(A) + P(\bar{A}) = 1$
For events \cap is AND, \cup is OR, so

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Example

For rolling a die $\Omega = \{1, 2, 3, 4, 5, 6\}$

Example

For rolling a die $\Omega = \{1, 2, 3, 4, 5, 6\}$

$P : 2^\Omega \rightarrow [0, 1]$, but for any event $A = \{1, 2, 5\}$

$$P(A) = P(\{1\}) + P(\{2\}) + P(\{5\})$$

Example

For rolling a die $\Omega = \{1, 2, 3, 4, 5, 6\}$

$P : 2^\Omega \rightarrow [0, 1]$, but for any event $A = \{1, 2, 5\}$

$$P(A) = P(\{1\}) + P(\{2\}) + P(\{5\})$$

So we need to know only p_1, p_2, \dots, p_6 , such that $\sum_{i=1}^6 p_i = 1$.

Example

For rolling a die $\Omega = \{1, 2, 3, 4, 5, 6\}$

$P : 2^\Omega \rightarrow [0, 1]$, but for any event $A = \{1, 2, 5\}$

$$P(A) = P(\{1\}) + P(\{2\}) + P(\{5\})$$

So we need to know only p_1, p_2, \dots, p_6 , such that $\sum_{i=1}^6 p_i = 1$.

This is why for any finite or countable set Ω we only need to define the probabilities of elements and then:

$$P(A) = \sum_{x \in A} P(\{x\})$$

Example with continuum set

Suppose you have some real value, i.e. $\Omega = \mathbb{R}$

$$A \subseteq \mathbb{R} \mapsto P(A)$$

Example with continuum set

Suppose you have some real value, i.e. $\Omega = \mathbb{R}$

$$A \subseteq \mathbb{R} \mapsto P(A)$$

Natural way

Is to define probability density as function $p: \mathbb{R} \rightarrow \mathbb{R}$:

Example with continuum set

Suppose you have some real value, i.e. $\Omega = \mathbb{R}$

$$A \subseteq \mathbb{R} \mapsto P(A)$$

Natural way

Is to define probability density as function $p: \mathbb{R} \rightarrow \mathbb{R}$:

- ▶ $\forall x \ p(x) \geq 0$

Example with continuum set

Suppose you have some real value, i.e. $\Omega = \mathbb{R}$

$$A \subseteq \mathbb{R} \mapsto P(A)$$

Natural way

Is to define probability density as function $p: \mathbb{R} \rightarrow \mathbb{R}$:

- ▶ $\forall x \ p(x) \geq 0$
- ▶ $\int_{\mathbb{R}} p(x) dx = 1$

Example with continuum set

Suppose you have some real value, i.e. $\Omega = \mathbb{R}$

$$A \subseteq \mathbb{R} \mapsto P(A)$$


Natural way

Is to define probability density as function $p: \mathbb{R} \rightarrow \mathbb{R}$:

- ▶ $\forall x \ p(x) \geq 0$
- ▶ $\int_{\mathbb{R}} p(x) dx = 1$

Then

$$P(A) = \int_A p(x) dx$$



If we think about $P(A)$ as «mass», then $p(x)$ is density.

Frequency interpretation

Frequency interpretation

Question

How to think about probability distribution P in pair (Ω, P) from the perspective of **black box**, answering your questions?

Frequency interpretation

Question

How to think about probability distribution P in pair (Ω, P) from the perspective of **black box**, answering your questions?

Suppose we have potentially infinite process of events generation:

$$\omega_1, \omega_2, \dots, \omega_n, \dots$$

Frequency interpretation

Question

How to think about probability distribution P in pair (Ω, P) from the perspective of **black box**, answering your questions?

Suppose we have potentially infinite process of events generation:

$$\omega_1, \omega_2, \dots, \omega_n, \dots$$

Then frequency limit

$$\lim_{n \rightarrow \infty} \frac{\#\{i : \omega_i \in A \mid 1 \leq i \leq n\}}{n} = P(A)$$

Random Permutations

A permutation of a set is an arrangement of its elements in a particular order. For a set of n elements, there are $n!$ (read as 'n factorial') possible permutations.

Random Permutations

A permutation of a set is an arrangement of its elements in a particular order. For a set of n elements, there are $n!$ (read as 'n factorial') possible permutations.

Definition (Random Permutation)

A random permutation of a set of n elements σ is a permutation chosen uniformly at random from the set of all $n!$ permutations, denoted as S_n .

In other words, each of the $n!$ permutations is equally likely to be chosen.

Random Permutations

A permutation of a set is an arrangement of its elements in a particular order. For a set of n elements, there are $n!$ (read as 'n factorial') possible permutations.

Definition (Random Permutation)

A random permutation of a set of n elements σ is a permutation chosen uniformly at random from the set of all $n!$ permutations, denoted as S_n .

In other words, each of the $n!$ permutations is equally likely to be chosen.

Example

For a set with three elements $\{1, 2, 3\}$, there are $3! = 6$ permutations:

$$(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1)$$

In a random permutation, each of these six arrangements is equally likely, so $P(\sigma) = \frac{1}{6}$ for any permutation $\sigma \in S_3$.

Task 1

Let event A be « σ is cycle of length n ». What is $P(A)$?

Task 1

Let event A be « σ is cycle of length n ». What is $P(A)$?

Solution

This is a purely combinatorial problem and

$$P(A) = \frac{\#(\text{of } n\text{-length cycles})}{n!}$$

Task 1

Let event A be « σ is cycle of length n ». What is $P(A)$?

Solution

This is a purely combinatorial problem and

$$P(A) = \frac{\#(\text{of } n\text{-length cycles})}{n!}$$

The numerator equals $\frac{n!}{n}$, so $P(A) = \frac{1}{n}$.

Task 2

Let event A be « σ is cycle of length $k < n$ ». What is $P(A)$?

Task 2

Let event A be « σ is cycle of length $k < n$ ». What is $P(A)$?

Solution

We need to chose k elements from all to form a cycle — this is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \text{ variants.}$$

Task 2

Let event A be « σ is cycle of length $k < n$ ». What is $P(A)$?

Solution

We need to chose k elements from all to form a cycle — this is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \text{ variants.}$$

Then we have $(k-1)!$ ways to build a certain cycle on k elements.

Task 2

Let event A be « σ is cycle of length $k < n$ ». What is $P(A)$?

Solution

We need to choose k elements from all to form a cycle — this is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \text{ variants.}$$

Then we have $(k-1)!$ ways to build a certain cycle on k elements.

The numerator equals $\binom{n}{k}(k-1)!$, so $P(A) = \frac{1}{k(n-k)!}$. Compare the probability with the previous one.

Conditional Probability

Suppose we ignore all the events from our **black box**, which is not from B .

Conditional Probability

Suppose we ignore all the events from our **black box**, which is not from B .

Definition

The conditional probability of an event A given that another event B has occurred is denoted by $P(A|B)$ and is defined as:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

where $P(B) > 0$.

Conditional Probability

Suppose we ignore all the events from our **black box**, which is not from B .

Definition

The conditional probability of an event A given that another event B has occurred is denoted by $P(A|B)$ and is defined as:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

where $P(B) > 0$.

- ▶ Conditional probability gives the probability of an event given that another event has occurred.

Conditional Probability

Suppose we ignore all the events from our **black box**, which is not from B .

Definition

The conditional probability of an event A given that another event B has occurred is denoted by $P(A|B)$ and is defined as:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

where $P(B) > 0$.

- ▶ Conditional probability gives the probability of an event given that another event has occurred.
- ▶ If events A and B are independent, then $P(A|B) = P(A)$.

Example

If the probability that it rains today is 0.5 (event B) and the probability that it rains both today and tomorrow is 0.3 (event A), then the conditional probability of it raining tomorrow given that it rains today, $P(A|B)$, is $\frac{0.3}{0.5} = 0.6$.

When events A and B are independent?

The area of event A with respect to Ω is the same as the area of event $A \cap B$ with respect to B . So

$$P(A) = P(A|B) \Rightarrow P(A \cap B) = P(A)P(B) \Leftrightarrow A \perp\!\!\!\perp B$$

Remarks

When events A and B are independent?

The area of event A with respect to Ω is the same as the area of event $A \cap B$ with respect to B . So

$$P(A) = P(A|B) \Rightarrow P(A \cap B) = P(A)P(B) \Leftrightarrow A \perp\!\!\!\perp B$$

Remarks

► $A \perp\!\!\!\perp B \Leftrightarrow A \perp\!\!\!\perp \overline{B}$

When events A and B are independent?

The area of event A with respect to Ω is the same as the area of event $A \cap B$ with respect to B . So

$$P(A) = P(A|B) \Rightarrow P(A \cap B) = P(A)P(B) \Leftrightarrow A \perp\!\!\!\perp B$$

Remarks

- ▶ $A \perp\!\!\!\perp B \Leftrightarrow A \perp\!\!\!\perp \overline{B}$
- ▶ If $P(A) > 0$ and $P(B) > 0$, but $P(A \cap B) = 0$ then they are dependent

When events A and B are independent?

The area of event A with respect to Ω is the same as the area of event $A \cap B$ with respect to B . So

$$P(A) = P(A|B) \Rightarrow P(A \cap B) = P(A)P(B) \Leftrightarrow A \perp\!\!\!\perp B$$

Remarks

- ▶ $A \perp\!\!\!\perp B \Leftrightarrow A \perp\!\!\!\perp \overline{B}$
- ▶ If $P(A) > 0$ and $P(B) > 0$, but $P(A \cap B) = 0$ then they are dependent
- ▶ If $A \subseteq B$ and $P(B) \neq 1$ then they are dependent

When events A and B are independent?

The area of event A with respect to Ω is the same as the area of event $A \cap B$ with respect to B . So

$$P(A) = P(A|B) \Rightarrow P(A \cap B) = P(A)P(B) \Leftrightarrow A \perp\!\!\!\perp B$$

Remarks

- ▶ $A \perp\!\!\!\perp B \Leftrightarrow A \perp\!\!\!\perp \overline{B}$
- ▶ If $P(A) > 0$ and $P(B) > 0$, but $P(A \cap B) = 0$ then they are dependent
- ▶ If $A \subseteq B$ and $P(B) \neq 1$ then they are dependent
- ▶ On contrary, \emptyset and Ω are independent with any other $A \subseteq \Omega$

Mutual Independence

Definition

A collection of events $\{A_i\}_{i=1}^n$ is said to be *Mutually Independent* if for every subset $\{A_{i_j}\}_{j=1}^k$ of the collection, the joint probability of all events in the subset is equal to the product of their individual probabilities:

$$P\left(\bigcap_{j=1}^k A_{i_j}\right) = \prod_{j=1}^k P(A_{i_j})$$

Mutual Independence

Definition

A collection of events $\{A_i\}_{i=1}^n$ is said to be *Mutually Independent* if for every subset $\{A_{i_j}\}_{j=1}^k$ of the collection, the joint probability of all events in the subset is equal to the product of their individual probabilities:

$$P\left(\bigcap_{j=1}^k A_{i_j}\right) = \prod_{j=1}^k P(A_{i_j})$$

- Mutual independence is a stronger condition than pairwise independence

Mutual Independence

Definition

A collection of events $\{A_i\}_{i=1}^n$ is said to be *Mutually Independent* if for every subset $\{A_{i_j}\}_{j=1}^k$ of the collection, the joint probability of all events in the subset is equal to the product of their individual probabilities:

$$P\left(\bigcap_{j=1}^k A_{i_j}\right) = \prod_{j=1}^k P(A_{i_j})$$

- ▶ Mutual independence is a stronger condition than pairwise independence
- ▶ If events are mutually independent, then they are pairwise independent, but not vice versa

Examples

First one

Tossing a fair coin multiple times. The outcome of each toss is an independent event. The outcome of any collection of tosses is the product of the probabilities of the individual tosses.


Examples

First one

Tossing a fair coin multiple times. The outcome of each toss is an independent event. The outcome of any collection of tosses is the product of the probabilities of the individual tosses.

Second one

Suppose we have two fair coins.




Examples

First one

Tossing a fair coin multiple times. The outcome of each toss is an independent event. The outcome of any collection of tosses is the product of the probabilities of the individual tosses.

Second one



Suppose we have two fair coins.
Pairwise independent


$$P(A) = P(B) = P(C) = \frac{1}{2}$$

Examples

First one

Tossing a fair coin multiple times. The outcome of each toss is an independent event. The outcome of any collection of tosses is the product of the probabilities of the individual tosses.

Second one



Suppose we have two fair coins.
Pairwise independent

$$P(A) = P(B) = P(C) = \frac{1}{2}$$

But not mutually independent


$$P(A)P(B)P(C) \neq P(A \cap B \cap C)$$

Examples

First one

Tossing a fair coin multiple times. The outcome of each toss is an independent event. The outcome of any collection of tosses is the product of the probabilities of the individual tosses.

Second one



Suppose we have two fair coins.
Pairwise independent

$$P(A) = P(B) = P(C) = \frac{1}{2}$$

But not mutually independent

$$P(A)P(B)P(C) \neq P(A \cap B \cap C)$$

$$A \text{ and } B \Rightarrow C$$

Task 3

Consider n coins, i.e. $\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}$. If they are mutually independent, then

$$P((a_1, \dots, a_n)) = P(a_1) \cdot \dots \cdot P(a_n) = \frac{1}{2^n}$$

Task 3

Consider n coins, i.e. $\Omega = \{(a_1, \dots, a_n) \mid a_i \in \{0, 1\}\}$. If they are mutually independent, then

$$P((a_1, \dots, a_n)) = P(a_1) \cdot \dots \cdot P(a_n) = \frac{1}{2^n}$$

This way we can model the two librarians problem. Suppose two persons independently decide if they pick up each of 10 books:

$$\begin{pmatrix} a_1 & \dots & a_{10} \\ b_1 & \dots & b_{10} \end{pmatrix}$$

So they simply toss 20 fair coins.

What is the probability of event, when they didn't pick the same book at the same time?

Solution

Solution

It means we must have at least one 0 in each column.

Solution

It means we must have at least one 0 in each column.

i.e. $P = \left(\frac{3}{4}\right)^{10} \sim 0.056$

Solution

It means we must have at least one 0 in each column.

I.e. $P = \left(\frac{3}{4}\right)^{10} \sim 0.056$

More general case

$X = \{1, \dots, n\}$, $A_1, \dots, A_k \subseteq X$ — random independent equiprobable subsets

What is $P(A_1 \cap \dots \cap A_k = \emptyset)$?

Solution

It means we must have at least one 0 in each column.

$$\text{i.e. } P = \left(\frac{3}{4}\right)^{10} \sim 0.056$$

More general case

$X = \{1, \dots, n\}$, $A_1, \dots, A_k \subseteq X$ — random independent equiprobable subsets

What is $P(A_1 \cap \dots \cap A_k = \emptyset)$?

Here we have $\Omega = \underbrace{2^X \times \dots \times 2^X}_k$

So we need $P(\forall \text{ column has } 0) = P(0 \in c_1 \cap \dots \cap 0 \in c_n)$

It turns out that these events are mutually independent, hence we have $P(0 \in c_1) \cdot \dots \cdot P(0 \in c_n) = \left(1 - \frac{1}{2^k}\right)^n$

Law of Total Probability

Law of Total Probability

Let B_1, B_2, \dots, B_n be a partition of the event (sample) space Ω , such that $P(B_i) > 0$ for all i . The Law (or Formula) of Total Probability states that for any event A , we have:

$$P(A) = \sum_{i=1}^n P(A|B_i) \cdot P(B_i)$$

Law of Total Probability

Let B_1, B_2, \dots, B_n be a partition of the event (sample) space Ω , such that $P(B_i) > 0$ for all i . The Law (or Formula) of Total Probability states that for any event A , we have:

$$P(A) = \sum_{i=1}^n P(A|B_i) \cdot P(B_i)$$

The law of total probability is a fundamental rule relating marginal probabilities and conditional probabilities.

It is easy to see, because $P(A) = \sum_{i=1}^n P(A \cap B_i)$

Law of Total Probability

Let B_1, B_2, \dots, B_n be a partition of the event (sample) space Ω , such that $P(B_i) > 0$ for all i . The Law (or Formula) of Total Probability states that for any event A , we have:

$$P(A) = \sum_{i=1}^n P(A|B_i) \cdot P(B_i)$$

The law of total probability is a fundamental rule relating marginal probabilities and conditional probabilities.

It is easy to see, because $P(A) = \sum_{i=1}^n P(A \cap B_i)$

Example

Suppose we have two boxes, one with 2 red and 3 green balls (box 1), and another with 3 red and 1 green ball (box 2). If we choose a box at random and then pick a ball, the law of total probability can be used to calculate the probability of drawing a red ball.

$$P(\text{Red}) = P(\text{Red}|\text{Box 1}) \cdot P(\text{Box 1}) + P(\text{Red}|\text{Box 2}) \cdot P(\text{Box 2})$$

Bayes' Theorem

Bayes' Theorem

Bayes' theorem is a fundamental principle in the field of statistics and probability. It describes how to update the probabilities of hypotheses when given evidence.

The theorem is stated mathematically as the following equation:

$$P(H|E) = \frac{P(E|H) \cdot P(H)}{P(E)}$$

where:

Bayes' Theorem

Bayes' theorem is a fundamental principle in the field of statistics and probability. It describes how to update the probabilities of hypotheses when given evidence.

The theorem is stated mathematically as the following equation:

$$P(H|E) = \frac{P(E|H) \cdot P(H)}{P(E)}$$

where:

- ▶ $P(H|E)$ is the probability of hypothesis H given the evidence E (a posteriori)

Bayes' Theorem

Bayes' theorem is a fundamental principle in the field of statistics and probability. It describes how to update the probabilities of hypotheses when given evidence.

The theorem is stated mathematically as the following equation:

$$P(H|E) = \frac{P(E|H) \cdot P(H)}{P(E)}$$

where:

- ▶ $P(H|E)$ is the probability of hypothesis H given the evidence E (a posteriori)
- ▶ $P(E|H)$ is the probability of the evidence given that the hypothesis is true (likelihood)

Bayes' Theorem

Bayes' theorem is a fundamental principle in the field of statistics and probability. It describes how to update the probabilities of hypotheses when given evidence.

The theorem is stated mathematically as the following equation:

$$P(H|E) = \frac{P(E|H) \cdot P(H)}{P(E)}$$

where:

- ▶ $P(H|E)$ is the probability of hypothesis H given the evidence E (a posteriori)
- ▶ $P(E|H)$ is the probability of the evidence given that the hypothesis is true (likelihood)
- ▶ $P(H)$ is the prior probability of H being true before the evidence is accounted

Bayes' Theorem

Bayes' theorem is a fundamental principle in the field of statistics and probability. It describes how to update the probabilities of hypotheses when given evidence.

The theorem is stated mathematically as the following equation:

$$P(H|E) = \frac{P(E|H) \cdot P(H)}{P(E)}$$

where:

- ▶ $P(H|E)$ is the probability of hypothesis H given the evidence E (a posteriori)
- ▶ $P(E|H)$ is the probability of the evidence given that the hypothesis is true (likelihood)
- ▶ $P(H)$ is the prior probability of H being true before the evidence is accounted
- ▶ $P(E)$ is the total probability of the evidence

Example

Suppose a disease affects 1% of a population. A test for the disease is 99% accurate. If a person tests positive for the disease, Bayes' theorem can be used to find out the probability that they actually have the disease.

Example

Suppose a disease affects 1% of a population. A test for the disease is 99% accurate. If a person tests positive for the disease, Bayes' theorem can be used to find out the probability that they actually have the disease.

$$P(\text{Disease}|\text{Positive Test}) = \frac{P(\text{Positive Test}|\text{Disease}) \cdot P(\text{Disease})}{P(\text{Positive Test})}$$


Example

Suppose a disease affects 1% of a population. A test for the disease is 99% accurate. If a person tests positive for the disease, Bayes' theorem can be used to find out the probability that they actually have the disease.

$$P(\text{Disease}|\text{Positive Test}) = \frac{P(\text{Positive Test}|\text{Disease}) \cdot P(\text{Disease})}{P(\text{Positive Test})}$$


$$\frac{0.99 \cdot 0.01}{0.99 \cdot 0.01 + 0.01 \cdot 0.99} = \frac{1}{2}$$

Task 4 (Casino)




If you are lucky to score N , then you win and leave. If you lose all the money to zero, then the game also ends.

Task 4 (Casino)



If you are lucky to score N , then you win and leave. If you lose all the money to zero, then the game also ends.
What is Ω here?


Task 4 (Casino)



If you are lucky to score N , then you win and leave. If you lose all the money to zero, then the game also ends.

What is Ω here? All the paths!

Task 4 (Casino)



If you are lucky to score N , then you win and leave. If you lose all the money to zero, then the game also ends.

What is Ω here? All the paths!

Question

Find the probability of winning.

Solution

Solution

$$P(k \text{ dollars and win since } 0 \text{ step}) = P(k \text{ dollars and win since } i \text{ step})$$

Solution

$P(k \text{ dollars and win since } 0 \text{ step}) = P(k \text{ dollars and win since } i \text{ step})$

Let $p_k = P(k \text{ dollars and win})$ and we will try to find them all. So

Solution

$P(k \text{ dollars and win since } 0 \text{ step}) = P(k \text{ dollars and win since } i \text{ step})$

Let $p_k = P(k \text{ dollars and win})$ and we will try to find them all. So

$$p_0 = 0, p_N = 1$$

$$p_k = P(k \text{ dollars and win} | \text{won on the first step})p + \\ P(k \text{ dollars and win} | \text{lost on the first step})(1-p)$$

$$p_k = p_{k+1}p + p_{k-1}(1-p) \Rightarrow \lambda = \lambda^2 p + (1-p)$$

$$\lambda^2 p - \lambda + (1-p) = 0 \Rightarrow \lambda_1 = \frac{1-p}{p}, \lambda_2 = 1$$

Solution

$P(k \text{ dollars and win since } 0 \text{ step}) = P(k \text{ dollars and win since } i \text{ step})$

Let $p_k = P(k \text{ dollars and win})$ and we will try to find them all. So

$$p_0 = 0, p_N = 1$$

$$p_k = P(k \text{ dollars and win} | \text{won on the first step})p + \\ P(k \text{ dollars and win} | \text{lost on the first step})(1-p)$$

$$p_k = p_{k+1}p + p_{k-1}(1-p) \Rightarrow \lambda = \lambda^2 p + (1-p)$$

$$\lambda^2 p - \lambda + (1-p) = 0 \Rightarrow \lambda_1 = \frac{1-p}{p}, \lambda_2 = 1$$

$$p \neq \frac{1}{2}$$

$$p_k = c_1 \left(\frac{1-p}{p} \right)^k + c_2$$

From the boundary conditions:

$$p_k = \frac{p^N - p^{N-k}(1-p)^k}{p^N - (1-p)^N}$$

Solution

$P(k \text{ dollars and win since } 0 \text{ step}) = P(k \text{ dollars and win since } i \text{ step})$

Let $p_k = P(k \text{ dollars and win})$ and we will try to find them all. So

$$p_0 = 0, p_N = 1$$

$$p_k = P(k \text{ dollars and win} | \text{won on the first step})p + \\ P(k \text{ dollars and win} | \text{lost on the first step})(1-p)$$

$$p_k = p_{k+1}p + p_{k-1}(1-p) \Rightarrow \lambda = \lambda^2 p + (1-p)$$

$$\lambda^2 p - \lambda + (1-p) = 0 \Rightarrow \lambda_1 = \frac{1-p}{p}, \lambda_2 = 1$$

$$p \neq \frac{1}{2}$$

$$p_k = c_1 \left(\frac{1-p}{p} \right)^k + c_2$$

From the boundary conditions:

$$p_k = \frac{p^N - p^{N-k}(1-p)^k}{p^N - (1-p)^N}$$

$$p = \frac{1}{2}$$

$$p_k = (c_1 + c_2 k) \lambda^k = c_1 + c_2 k$$

From the boundary conditions:

$$c_1 = 0, c_2 = \frac{1}{N} \Rightarrow p_k = \frac{k}{N}$$

Continuous Probability Distributions

Continuous Probability Distributions

Definition

A random variable X has a *continuous distribution* if there exists a function $f(x)$ called the probability density function (pdf) of X such that for any two numbers a and b with $a \leq b$,

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

Continuous Probability Distributions

Definition

A random variable X has a *continuous distribution* if there exists a function $f(x)$ called the probability density function (pdf) of X such that for any two numbers a and b with $a \leq b$,

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

- ▶ $f(x) \geq 0$ for all x , and $\int_{-\infty}^{\infty} f(x) dx = 1$

Continuous Probability Distributions

Definition

A random variable X has a *continuous distribution* if there exists a function $f(x)$ called the probability density function (pdf) of X such that for any two numbers a and b with $a \leq b$,

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

- ▶ $f(x) \geq 0$ for all x , and $\int_{-\infty}^{\infty} f(x) dx = 1$
- ▶ Unlike a discrete distribution, $P(X = x) = 0$ for all x in a continuous distribution

Continuous Probability Distributions

Definition

A random variable X has a *continuous distribution* if there exists a function $f(x)$ called the probability density function (pdf) of X such that for any two numbers a and b with $a \leq b$,

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

- ▶ $f(x) \geq 0$ for all x , and $\int_{-\infty}^{\infty} f(x) dx = 1$
- ▶ Unlike a discrete distribution, $P(X = x) = 0$ for all x in a continuous distribution
- ▶ The expected value (or mean) μ of X is given by

$$\mu = \int_{-\infty}^{\infty} xf(x) dx$$

Examples

Examples


- ▶ The Uniform distribution on an interval $[a, b]$ is a simple example of a continuous distribution

Examples

- ▶ The Uniform distribution on an interval $[a, b]$ is a simple example of a continuous distribution
- ▶ The Normal or Gaussian distribution is another common continuous distribution



Task 5

$P(\triangle ABC \text{ is obtuse triangle}) = ?$



Task 5

$P(\triangle ABC \text{ is obtuse triangle}) = ?$



One slide about Pseudo-Random Processes

Definition

A pseudo-random process is a process that appears to be random but is generated by a deterministic process.

One slide about Pseudo-Random Processes

Definition

A pseudo-random process is a process that appears to be random but is generated by a deterministic process.

- ▶ Pseudo-random sequences are not truly random because they are generated using deterministic methods (computer memory is always limited)

One slide about Pseudo-Random Processes

Definition

A pseudo-random process is a process that appears to be random but is generated by a deterministic process.

- ▶ Pseudo-random sequences are not truly random because they are generated using deterministic methods (computer memory is always limited)
- ▶ They are often used in computer simulations and cryptography due to their predictability, repeatability, and efficiency

One slide about Pseudo-Random Processes

Definition

A pseudo-random process is a process that appears to be random but is generated by a deterministic process.

- ▶ Pseudo-random sequences are not truly random because they are generated using deterministic methods (computer memory is always limited)
- ▶ They are often used in computer simulations and cryptography due to their predictability, repeatability, and efficiency

Example

Consider a linear congruential generator (LCG), a type of pseudo-random number generator, defined by the recurrence relation:

$$X_{n+1} = (aX_n + c) \mod m$$

where X is the sequence of pseudo-random values, and a , c , m are constants. Despite the appearance of randomness in the sequence X , it is entirely determined by the choice of initial seed X_0 .

Fibonacci Generator

Fibonacci Generator

Definition

A Fibonacci generator is a type of pseudo-random number generator that generates numbers using the Fibonacci sequence.

Fibonacci Generator

Definition

A Fibonacci generator is a type of pseudo-random number generator that generates numbers using the Fibonacci sequence.

- ▶ It uses the relation: $X_{n+2} = (X_{n+1} + X_n) \bmod m$
- ▶ It appears to generate random numbers, but the numbers are completely determined by the initial values X_0 and X_1 .

Fibonacci Generator

Definition

A Fibonacci generator is a type of pseudo-random number generator that generates numbers using the Fibonacci sequence.

- ▶ It uses the relation: $X_{n+2} = (X_{n+1} + X_n) \bmod m$
- ▶ It appears to generate random numbers, but the numbers are completely determined by the initial values X_0 and X_1 .

Example

Suppose we start with $X_0 = 0$ and $X_1 = 1$ and choose $m = 10$. The generated sequence would be: 0, 1, 1, 2, 3, 5, 8, 3, 1, 4, 5, 9, 4, 3, 7, 0, 7, 7, 4, 1, 5, 6, 1, 7, 8, 5, 3, 8, 1, 9, 0, 9, 9, 8, 7, 5, 2, 7, 9, 6, 5, 1, 6, 7, 3, 0, 3, 3, 6, 9, 5, 4, 9, 3, 2, 5, 7, 2, 9, 1, 0, 1, 1, 2, 3, 5, 8, 3, 1, 4, 5, 9, 4, 3, 7, 0, 7, 7, 4, 1, 5, 6, 1, 7, 8, 5, 3, 8, 1, 9, 0, ...