Supporting Information File

This PDF file includes:

Material and methods
Table S1 – S2
References

Material and methods

We conducted systematic and snowball literature reviews to gather empirical data and examine how human activity affected the activity and overlap of predator-prey dyads. For the systematic review, we used Web of Science to search the published literature from 2000 – 2020, using the query "TS = ("activity pattern*" OR "temporal*" OR "diel" OR "nocturnal*" OR "diurnal*" OR "time of day" OR "camera trap*" OR "overlap*" OR "co-occurrence" OR "cooccurrence" OR "occupancy") AND ("human*" OR "people" OR "anthropogenic" OR "touris*" OR "recreat*" OR "disturb*") AND ("wildlife" OR "mammal*" OR "animal*" OR "*carnivor*" OR "herbivor*" OR "ungulate*" OR "predator*" OR "prey" OR "commun*") NOT ("marine*" OR "*fish*")" to match titles, abstracts, and keywords within the topics of Environmental Sciences & Ecology, Zoology, or Biodiversity & Conservation. We read the resulting paper titles and abstracts (n = 6,646) and read the full study if it showed potential to meet our criteria (n = 405).

We retained studies that (1) measured the diel activity of at least one terrestrial mammal predator and terrestrial mammal prey dyad <1 kg in body mass, and (2) used camera traps to compare animal activity at paired sites or treatments of high and low human use. We limited our review to camera trap studies given that the temporal dimension is often overlooked, easily standardized, and eliminates confounding lethal or density effects of spatial indices. For our analysis, we defined "human activity" as sustained, non-lethal presence of humans. Targeting studies that measured non-lethal human activity allowed us to examine behavioral effects rather than the numerical effects of humans on wildlife. Our definition of human activity was intentionally broad (e.g., people on foot, motorized and non-motorized recreation, pastoralism, energy infrastructure, housing, roads, urban areas) as our aim was to capture and compare the multiple pathways of animal response to disturbance. We distinguished high and low human use according to author descriptions in each peer-reviewed paper (**Table S1**). We designated predator-prey dyads based on predator diet as listed in the paper, expert advice, or the Encyclopedia of Life (Parr et al. 2014) (see **Table S2** and **Table S3** for the list of included predator-prey dyads).

To examine predator-prey temporal activity, we extracted day and night image counts of each species at each high and low human use to calculate the ratio of diurnal activity. When these metrics were not available, we used the raw data with author permission, filtered for image independence as greater than 30-minutes apart (Sollmann et al. 2018), and classified images

based on sun-time (Nouvellet et al. 2012; see **Table S1** for detailed methods on each study). We discarded any dyads for which there were fewer than 10 total images of a species. We centered the diurnal activity ratio on zero and standardized it so that -1 represented complete nocturnality and 1 complete diurnality. Then, we set the rescaled diurnal activity ratios at low human use to zero and calculated the relative difference for diurnal activity ratios at high human use across predator-prey dyads. This allowed us to simultaneously compare the relative magnitude of diurnal activity changes across all predator-prey dyads (**Fig. 2a**). We made the assumption that the majority of human activity was diurnal across studies, thus, we considered any difference in activity between low and high human use to be in response to diurnal human activity.

To examine predator-prey temporal overlap, we compared temporal overlap coefficients of each species pair at low and high human use. Overlap coefficients (Δ) are widely used metrics to report temporal partitioning of animals from camera trap surveys and are calculated using kernel density estimation that range from 0 (no overlap) to 1 (complete overlap) (Ridout & Linkie 2009). When these metrics were not available, we used published datasets or the raw data with author permission to estimate Δ using the 'overlap' R package. We calculated Δ using Dhat4, according to the protocol of Ridout & Linkie (2009). We were able to include or calculate overlap coefficients for 17 of the studies and 172 dyads. We calculated the change in overlap as the difference between predator-prey overlap at high and low human use (**Fig. 2b)**.

Out of the 405 abstracts that warranted a full review, we excluded 155 studies that did not use camera trap array study designs, 75 studies for lacking concurrent data on mammal predators and prey, 80 studies that did not adequately distinguish between high and low human use, 28 studies that had fewer than 10 camera sites or did not include temporal data, and 48 studies with temporal data in the wrong format for our analysis. Ultimately, there were 19 studies that had data that fit our criteria for inclusion, with 178 predator-prey dyads included in the temporal activity analysis (**Fig. 2a**). Of these, we included 167 predator-prey dyads from 17 studies in the temporal overlap analysis (**Fig. 2b**). We eliminated 11 dyads for the overlap analysis, given that 5 dyads did not include data to calculate overlap coefficients, and 6 dyads exhibited a neutral activity response and thus did not warrant further testing via our framework's approach.

We examined how functional traits (prey order, body size, predator hunting mode, trophic level, predator guild, and circadian rhythm) influenced temporal activity of predator and prey species (unique to each study) between settings of high and low human activity (**Fig. 3**).

Table S1. Site descriptions, human activity level, and method used to calculate predator-prey activity ratio and overlap coefficient of each study in analysis (n = 19).

Reference	Latitude, Longitude	Location	Habitat Type	Human activity type	Data acquisition and human activity	Activity ratio	Overlap coefficient
Baker & Leberg 2018	32.28, -111.15	USA	Desert	Human presence (on foot, vehicles, human density)	Acquired record table from authors. We used the author's classification of SAGU site as high human activity and CHIR site as low human activity.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Barrueto *et al.* 2014	51.2, -115.6	Canada	Forest	Human presence (on foot), Roads	Acquired data from Figure 2. We used the authors classification of backcountry and wildlife crossings as low and high human activity.	We digitized percent data from Figure 2, transformed it into image counts using the image totals from the paper, then calculated day/night activity ratios based on sun-time for the study period.	We digitized percent data from Figure 2, transformed it into image counts using the image totals from the paper, then estimated the overlap coefficient with the 'overlap' package in R (delta 4, bootstrap 1000).
Caldwell *et al.* 2020	39.68, -120.02	USA	Shrubland, Woodland	Human presence (on foot, vehicles)	Accquired record table from authors. We classified low human activity as less than 25 total human and/or vehicle detections (2 sites, cameras 35 & 33), and high human activity as greater than 25 total human and/or vehicle detections (4 sites, remaining).	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to calculate hourly counts and estimate the overlap coefficient with the 'overlap' package in R (delta 4, bootstrap 1000).
Carter *et al.* 2015	27.56, 84.38	Nepal	Forest	Inside vs. outside National Park	Acquired record table from authors. We used the author's classification of high human activity as sites outside the park, and low human activity as sites inside the park, as well as the author's classification of day/night.	We used the raw record table to calculate day/night image counts by species for 2010 & 2011 to calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Dawson *et al.* 2018	-17.90, 122.83	Australia	Savanna, Woodland	Energy infrastructure (seismic lines)	Acquired data from Table S3 and Figure 6. We used the author's classification of control and seismic line camera placement.	The author provided image counts from Appendix S2, as broken down by day/night to calculate diel activity ratios.	We used values in Table S3 and Figure 6.
Díaz-Ruíz *et al.* 2016	38.56, -3.28	Spain	Savanna	Agriculture, Housing	We used the author's classification of agricultural sites as high human activity and shrubland sites as low human activity.	We digitized Figure 3 to estimate the proportion of hourly activity, we averaged hourly activity for agricultural and shrubland sites which we used to calculate day/night activity ratios.	We used data from Table 2. We averaged the overlap coefficients for agricultural sites with both species present (1, 3, 4, and 8) and shrubland sites (5, 6, 10, and 11)
Gallo *et al.* 2019	41.88, -87.63	USA	Urban	Urban housing	Acquired data from Dryad. We replicated the authors' procedure of distinguishing high/low human activity sites using a PCA of the site covariates mean canopy, mean housing, and mean impervious, and counted values <= 0 as high human activity.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Gray *et al.* 2011	12.75, 106.95	Cambodia	Forest	Human presence (on foot), Resource extraction	Activity ratios aqcuired with permission from Gaynor et al. 2018.	Acquired activity ratios with permission from Gaynor et al. 2018.	No overlap data available.
Kays *et al.* 2017	,	USA	Forest	Human presence (on foot, vehicles, dogs, horses)	Accquired study data from Dryad. We classified low human activity as 'off trail' sites (50 or 200m from trails), and high human activity as 'on trail' sites.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Lendrum *et al.* 2017	39.92, -108.20	USA	Forest	Human presence, Energy infrastructure	Accquired study data from Table 2 and Figure 2. We used the authors site classifications as developed (high human activity) and undeveloped (low human activity) based on the presence of energy infrastructure and human activity.	We used activity data from Table 2. We evenly divided dawn/dusk crepuscular activity into the nocturnal/diurnal categories.	We digitized the plots in Figure 2 to estimate the overlap of the density curves using the overlapTrue() function in the 'overlap' package in R. (CIs could not be calculated without raw data).

Mills & Harris 2020	11.52, 1.97	Burkina Faso, Benin, Niger	Savanna, Woodland	Human presence (on foot)	Accquired study data from Figure 3 and Figure 4. We used the author's classification of high human activity as > 0.54 mean occupancy of humans.	We extracted the nocturnal activity for each species by digitizing Figure 4, we then calculated the proportion of diurnal activity as 1 - nocturnal activity.	We digitized the plots in Figure 3 to estimate the overlap of the density curves using the overlapTrue() function in the 'overlap' package in R. (CIs could not be calculated without raw data).
Nix *et al.* 2018	40.03, -111.50	USA	Forest	Human presence (on foot, camping)	Record table acquired with permission from Gaynor et al. 2018. We used the authors classification of low human activity as the weekday (Monday - Thursday), and high human activity as the weekend (Friday - Sunday).	We used the record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the record table to filter independent detections and estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Oberosler *et al.* 2017	46.11, 10.93	Italy	Forest	Human presence, Logging, Roads	Acquired data from Figure 4.	We digitized Figure 4 to estimate the proportion of hourly activity at sites of high and low human activity, which we used to calculate day/night activity ratios.	No overlap data available.
Reilly *et al.* 2017	37.77, -122.42	USA	Savanna	Human presence (on foot, horses, vehicles)	Acquired record table from authors. We used the author's classification of high and low human activity, but combined high/medium and low/none categories.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Rich *et al.* 2016	-19.52, 23.63	Botswana	Savanna	Human presence (on foot, livestock, vehicles)	Acquired record table from authors. We used the authors recommendation to classify high human activity as the livestock grazing area and low human activity as the wildlife management area and game reserves.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Smith *et al.* 2019	-30.91, 30.28	South Africa	Woodland, Shrubland	Land use type (residential, agricultural, reserve)	Accquired data from Figure 4, Figure 5, and Figure 6. We used the author's classification of agricultural as high human activity sites, and reserve land as low human activity sites.	We used activity data from Table 2. We evenly divided dawn/dusk crepuscular activity into the nocturnal/diurnal categories.	We used the overlap coefficients in Figure 4, Figure 5, and Figure 6.
Wang *et al.* 2015	37.16, -122.08	USA	Forest	Human presence (on foot)	Acquired record table from authors. We used the author's classification of high and low human activity.	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the overlap coefficients in Table 3.
Wilkinson *et al.* 2021	-0.42, 36.09	Kenya	Savanna	Human presence (on foot, livestock)	Acquired record table from authors. We used the author's classification as sites of high human activity (>= 1 detection/trap night, n = 8 sites), and low human activity (< 1 detection/trap night, n = 11 sites)	We used the raw record table to filtered images for independence, classified day/night using sun-time for species, and calculate activity ratios.	We used the raw record table to estimate overlap coefficients with the 'overlap' package in R (delta 4, bootstrap 1000).
Xiao *et al.* 2018	42.85, 130.40	China	Forest	Human presence (on foot), Housing, Roads, Logging	Acquired data from Figure 8. We used the author's classification of the summer season as high human activity and the winter season as low human activity.	We used the authors breakdown of percent of daytime and night time images for each species.	We used the overlap coefficients in Figure 8.

Table S2. Summary of predator and prey activity ratios at high and low human use for 19 studies and 178 predator-prey dyads.

Reference	Predator		Prey		Predator diurnal activity		Prey diurnal activity	
					(low)	(high)	(low)	(high)
Baker & Leburg 2018	Bobcat	*(Lynx rufus)*	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.316	0.241	0.044	0.304
			Desert Cottontail	*(Sylvilagus audubonii)*			0.165	0.429
	Coyote	*(Canis latrans)*	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.1	0.216	0.044	0.304
			Desert Cottontail	*(Sylvilagus audubonii)*			0.165	0.429
			Deer spp.	*(Odocoileus hemionus, Odocoileus virginianus)*			0.578	0.463
			Javelina	*(Tayassu tajacu)*			0.538	0.25
	Gray Fox	*(Urocyon cinereoargenteus)*	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.035	0.04	0.044	0.304
			Desert Cottontail	*(Sylvilagus audubonii)*			0.165	0.429
Barrueto et al. 2014	Black Bear	*(Ursus americanus)*	Mule Deer	*(Odocoileus hemionus)*	0.744	0.646	0.631	0.599
	Brown Bear	*(Ursus arctos)*	Elk	*(Cervus elaphus)*	0.748	0.679	0.378	0.253
			Mule Deer	*(Odocoileus hemionus)*			0.631	0.599
			Moose	*(Alces alces)*			0.474	0.498
	Coyote	*(Canis latrans)*	Mule Deer	*(Odocoileus hemionus)*	0.563	0.481	0.631	0.599
	Puma	*(Puma concolor)*	Mule Deer	*(Odocoileus hemionus)*	0.411	0.262	0.631	0.599
	Wolf	*(Canis lupus)*	Elk	*(Cervus elaphus)*	0.463	0.44	0.378	0.253
			Mule Deer	*(Odocoileus hemionus)*			0.631	0.599
			Moose	*(Alces alces)*			0.474	0.498
Caldwell & Klip 2020	Bobcat	*(Lynx rufus)*	Lagomorph spp.	*(Sylvagus spp., Lepus californicus)*	0	0.337	0.762	0.631
	Coyote	*(Canis latrans)*	Lagomorph spp.	*(Sylvagus spp., Lepus californicus)*	0.5	0.534	0.762	0.631
			Mule Deer	*(Odocoileus hemionus)*			0.35	0.318
Carter et al. 2015	Leopard	*(Panthera pardus)*	Chital	*(Axis axis)*	0.535	0.157	0.743	0.527
			Muntjac	*(Muntiacus muntjak)*			0.648	0.527
			Wild Boar	*(Sus scrofa)*			0.624	0.254
			Sambar	*(Rusa unicolor)*			0.074	0.115
	Tiger	*(Panthera tigris)*	Chital	*(Axis axis)*	0.19	0.073	0.743	0.527
			Muntjac	*(Muntiacus muntjak)*			0.648	0.527
			Wild Boar	*(Sus scrofa)*			0.624	0.254
			Sambar	*(Rusa unicolor)*			0.074	0.115
Dawson et al. 2018	Dingo	*(Canis lupus dingo)*	Agile Wallaby	*(Macropus agilis)*	0.7	0.336	0.267	0.285
			Greater Bilby	*(Macrotis lagotis)*			0	0.022
	Domestic Cat	*(Felis catus)*	Agile Wallaby	*(Macropus agilis)*	0.302	0.142	0.267	0.285
			Greater Bilby	*(Macrotis lagotis)*			0	0.022
Díaz-Ruíz et al. 2016	Red Fox	*(Vulpes vulpes)*	European Rabbit	*(Oryctolagus cuniculus)*	0.26	0.181	0.889	0.937
Gallo et al. 2019	Coyote	*(Canis latrans)*	Eastern Cottontail	*(Sylvilagus floridanus)*	0.209	0.152	0.247	0.23
			White-Tailed Deer	*(Odocoileus virginianus)*			0.54	0.722
Gray et al. 2011	Leopard	*(Panthera pardus)*	Muntjac	*(Muntiacus muntjak)*	0.613	0.55	0.713	0.612
			Wild Boar	*(Sus scrofa)*			0.838	0.432

Kays et al. 2017	Black Bear	*(Ursus americanus)*	Eastern Fox Squirrel	*(Sciurus niger)*	0.781	0.663	0.986	0.971
			Virginia Opossum	*(Didelphis virginiana)*			0.013	0
			White-Tailed Deer	*(Odocoileus virginianus)*			0.577	0.502
			Wild Boar	*(Sus scrofa)*			0.444	0.333
			Eastern Cottontail	*(Sylvilagus floridanus)*			0.03	0.088
			Eastern Gray Squirrel	*(Sciurus carolinensis)*			0.962	0.982
			Raccoon	*(Procyon lotor)*			0.054	0.06
	Bobcat	*(Lynx rufus)*	Eastern Fox Squirrel	*(Sciurus niger)*	0.53	0.423	0.986	0.971
			Eastern Cottontail	*(Sylvilagus floridanus)*			0.03	0.088
			Eastern Gray Squirrel	*(Sciurus carolinensis)*			0.962	0.982
	Coyote	*(Canis latrans)*	Eastern Fox Squirrel	*(Sciurus niger)*	0.284	0.156	0.986	0.971
			Virginia Opossum	*(Didelphis virginiana)*			0.013	0
			White-Tailed Deer	*(Odocoileus virginianus)*			0.577	0.502
			Wild Boar	*(Sus scrofa)*			0.444	0.333
			Eastern Cottontail	*(Sylvilagus floridanus)*			0.03	0.088
			Eastern Gray Squirrel	*(Sciurus carolinensis)*			0.962	0.982
			Raccoon	*(Procyon lotor)*			0.054	0.06
	Gray Fox	*(Urocyon cinereoargenteus)*	Eastern Fox Squirrel	*(Sciurus niger)*	0.107	0.086	0.986	0.971
			Eastern Cottontail	*(Sylvilagus floridanus)*			0.03	0.088
			Eastern Gray Squirrel	*(Sciurus carolinensis)*			0.962	0.982
	Red Fox	*(Vulpes vulpes)*	Eastern Fox Squirrel	*(Sciurus niger)*	0.27	0.133	0.986	0.971
			Eastern Cottontail	*(Sylvilagus floridanus)*			0.03	0.088
			Eastern Gray Squirrel	*(Sciurus carolinensis)*			0.962	0.982
Lendrum et al. 2017	Bobcat	*(Lynx rufus)*	Lagomorph spp.	*(Sylvilagus nuttallii, Lepus californicus)*	0.346	0.232	0.25	0.163
	Coyote	*(Canis latrans)*	Lagomorph spp.	*(Sylvilagus nuttallii, Lepus californicus)*	0.588	0.382	0.25	0.163
			Mule Deer	*(Odocoileus hemionus)*			0.686	0.509
Mills & Harris 2020	Leopard	*(Panthera pardus)*	Aardvark	*(Orycteropus afer)*	0.366	0.412	0.165	0.314
			Hartebeest	*(Alcelaphus buselaphus)*			0.726	0.78
			Kob	*(Kobus kob)*			0.826	0.878
			Oribi	*(Ourebia ourebi)*			0.861	0.885
			Waterbuck	*(Kobus ellipsiprymnus)*			0.819	0.834
			Bushbuck	*(Tragelaphus sylvaticus)*			0.599	0.528
			Duiker spp.	*(Sylvicapra grimmia, Cephalophus rufilatus)*			0.625	0.551
			Reedbuck	*(Redunca redunca)*			0.592	0.468
			Roan Antelope	*(Hippotragus equinus)*			0.787	0.742
			Warthog	*(Phacochoerus africanus)*			0.987	0.941
	Lion	*(Panthera leo)*	Bushbuck	*(Tragelaphus sylvaticus)*	0.561	0.44	0.599	0.528
			Cape Buffalo	*(Syncerus caffer)*			0.578	0.534
			Duiker spp.	*(Sylvicapra grimmia, Cephalophus rufilatus)*			0.625	0.551
			Reedbuck	*(Redunca redunca)*			0.592	0.468
			Roan Antelope	*(Hippotragus equinus)*			0.787	0.742
			Warthog	*(Phacochoerus africanus)*			0.987	0.941
			Aardvark	*(Orycteropus afer)*			0.165	0.314
			Hartebeest	*(Alcelaphus buselaphus)*			0.726	0.78
			Kob	*(Kobus kob)*			0.826	0.878
			Oribi	*(Ourebia ourebi)*			0.861	0.885
			Waterbuck	*(Kobus ellipsiprymnus)*			0.819	0.834
	Spotted Hyena	*(Crocuta crocuta)*	Bushbuck	*(Tragelaphus sylvaticus)*	0.453	0.413	0.599	0.528
			Cape Buffalo	*(Syncerus caffer)*			0.578	0.534
			Duiker spp.	*(Sylvicapra grimmia, Cephalophus rufilatus)*			0.625	0.551
			Reedbuck	*(Redunca redunca)*			0.592	0.468
			Roan Antelope	*(Hippotragus equinus)*			0.787	0.742
			Warthog	*(Phacochoerus africanus)*			0.987	0.941
			Aardvark	*(Orycteropus afer)*			0.165	0.314
			Hartebeest	*(Alcelaphus buselaphus)*			0.726	0.78
			Kob	*(Kobus kob)*			0.826	0.878
			Oribi	*(Ourebia ourebi)*			0.861	0.885
			Waterbuck	*(Kobus ellipsiprymnus)*			0.819	0.834

Nix et al. 2018	Puma	*(Puma concolor)*	Beaver	*(Castor canadensis)*	0	0	0.047	0.061
			Raccoon	*(Procyon lotor)*			0.018	0.08
			Mule Deer	*(Odocoileus hemionus)*			0.578	0.561
Oberosler et al. 2017	Red Fox	*(Vulpes vulpes)*	Brown Hare	*(Lepus europaeus)*	0.239	0.248	0.6	0.239
Reilly et al. 2017	Bobcat	*(Lynx rufus)*	Brush Rabbit	*(Sylvilagus bachmani)*	0.38	0.314	0.494	0.729
	Coyote	*(Canis latrans)*	Mule Deer	*(Odocoileus hemionus)*	0.459	0.364	0.615	0.585
			Raccoon	*(Procyon lotor)*			0.097	0.078
			Brush Rabbit	*(Sylvilagus bachmani)*			0.494	0.729
	Gray Fox	*(Urocyon cinereoargenteus)*	Brush Rabbit	*(Sylvilagus bachmani)*	0.049	0.059	0.494	0.729
Rich et al. 2016	Black-backed Jackal	*(Canis mesomelas)*	Springhare	*(Pedetes capensis)*	0.162	0.372	0.097	0.118
			Cape Hare	*(Lepus capensis)*			0.213	0.177
	Caracal	*(Caracal caracal)*	Cape Hare	*(Lepus capensis)*	0.14	0.136	0.213	0.177
			Springhare	*(Pedetes capensis)*			0.097	0.118
	Leopard	*(Panthera pardus)*	Duiker	*(Sylvicapra grimmia)*	0.333	0.254	0.554	0.432
			Warthog	*(Phacochoerus africanus)*			0.887	0.819
			Greater Kudu	*(Tragelaphus strepsiceros)*			0.584	0.6
			Impala	*(Aepyceros melampus)*			0.735	0.746
	Lion	*(Panthera leo)*	Buffalo	*(Bubalus bubalis)*	0.417	0.322	0.313	0.275
			Duiker	*(Sylvicapra grimmia)*			0.554	0.432
			Warthog	*(Phacochoerus africanus)*			0.887	0.819
			Greater Kudu	*(Tragelaphus strepsiceros)*			0.584	0.6
			Impala	*(Aepyceros melampus)*			0.735	0.746
			Zebra	*(Equus burchelli)*			0.511	0.526
	Spotted Hyena	*(Crocuta crocuta)*	Buffalo	*(Bubalus bubalis)*	0.239	0.179	0.313	0.275
			Duiker	*(Sylvicapra grimmia)*			0.554	0.432
			Warthog	*(Phacochoerus africanus)*			0.887	0.819
			Greater Kudu	*(Tragelaphus strepsiceros)*			0.584	0.6
			Impala	*(Aepyceros melampus)*			0.735	0.746
			Zebra	*(Equus burchelli)*			0.511	0.526
	Wild Dog	*(Lycaon pictus)*	Greater Kudu	*(Tragelaphus strepsiceros)*	0.443	0.748	0.584	0.6
			Impala	*(Aepyceros melampus)*			0.735	0.746
			Springhare	*(Pedetes capensis)*			0.097	0.118
			Zebra	*(Equus burchelli)*			0.511	0.526
			Cape Hare	*(Lepus capensis)*			0.213	0.177
			Duiker	*(Sylvicapra grimmia)*			0.554	0.432
			Warthog	*(Phacochoerus africanus)*			0.887	0.819
Smith et al. 2019	Black-backed Jackal	*(Canis mesomelas)*	Blue Duiker	*(Philantomba monticola)*	0.52	0.49	0.665	0.65
			Duiker	*(Sylvicapra grimmia)*			0.71	0.735
	Caracal	*(Caracal caracal)*	Blue Duiker	*(Philantomba monticola)*	0.6	0.475	0.665	0.65
			Bushbuck	*(Tragelaphus sylvaticus)*			0.625	0.655
			Duiker	*(Sylvicapra grimmia)*			0.71	0.735
Wang et al. 2015	Coyote	*(Canis latrans)*	Mule Deer	*(Odocoileus hemionus)*	0.503	0.305	0.525	0.551
			Raccoon	*(Procyon lotor)*			0.067	0.17
			Virginia Opossum	*(Didelphus virginianus)*			0.028	0.043
	Puma	*(Puma concolor)*	Mule Deer	*(Odocoileus hemionus)*	0.263	0.136	0.525	0.551
			Raccoon	*(Procyon lotor)*			0.067	0.17
			Virginia Opossum	*(Didelphus virginianus)*			0.028	0.043

Wilkinson et al. 2021	Black-backed Jackal	*(Canis mesomelas)*	Kirk's Dikdik	*(Madoqua kirkii)*	0.382	0.063	0.789	0.359
			Thomsons Gazelle	*(Eudorcas thomsonii)*			1	0.712
			African Hare	*(Lepus victoriae)*			0	0.014
	Leopard	*(Panthera pardus)*	Common Zebra	*(Equus quagga)*	0.164	0.08	0.254	0.195
			Impala	*(Aepyceros melampus)*			0.927	0.568
			Kirk's Dikdik	*(Madoqua kirkii)*			0.789	0.359
			Olive Baboon	*(Papio anubis)*			1	0.999
			Vervet	*(Chlorocebus pygerythrus)*			1	0.999
			Aardvark	*(Orycteropus afer)*			0	0
			Warthog	*(Phacochoerus africanus)*			0.953	0.983
			Waterbuck	*(Kobus ellipsiprymnus)*			0.167	0.605
	Lion	*(Panthera leo)*	Warthog	*(Phacochoerus africanus)*	0	0.075	0.953	0.983
			Waterbuck	*(Kobus ellipsiprymnus)*			0.167	0.605
			Aardvark	*(Orycteropus afer)*			0	0
			Buffalo	*(Bubalus bubalis)*			0.058	0.03
			Common Zebra	*(Equus quagga)*			0.254	0.195
			Eland	*(Taurotragus oryx)*			0.651	0.477
			Impala	*(Aepyceros melampus)*			0.927	0.568
			Kirk's Dikdik	*(Madoqua kirkii)*			0.789	0.359
			Olive Baboon	*(Papio anubis)*			1	0.999
			Thomsons Gazelle	*(Eudorcas thomsonii)*			1	0.712
	Serval	*(Leptailurus serval)*	African Hare	*(Lepus victoriae)*	0.093	0.053	0	0.014
	Spotted Hyena	*(Crocuta crocuta)*	Buffalo	*(Bubalus bubalis)*	0.152	0.019	0.058	0.03
			Common Zebra	*(Equus quagga)*			0.254	0.195
			Eland	*(Taurotragus oryx)*			0.651	0.477
			Impala	*(Aepyceros melampus)*			0.927	0.568
			Kirk's Dikdik	*(Madoqua kirkii)*			0.789	0.359
			Olive Baboon	*(Papio anubis)*			1	0.999
			Thomsons Gazelle	*(Eudorcas thomsonii)*			1	0.712
			Aardvark	*(Orycteropus afer)*			0	0
			Warthog	*(Phacochoerus africanus)*			0.953	0.983
			Waterbuck	*(Kobus ellipsiprymnus)*			0.167	0.605
	Striped Hyena	*(Hyaena hyaena)*	Olive Baboon	*(Papio anubis)*	0.13	0.021	1	0.999
Xiao et al. 2018	Tiger	*(Panthera tigris)*	Roe Deer	*(Capreolus capreolus)*	0.295	0.392	0.577	0.665
			Sika Deer	*(Cervus nippon)*			0.635	0.655
			Wild Boar	*(Sus scrofa)*			0.633	0.584

Table S3. Summary of predator and prey overlap coefficients at high and low human use for 17 studies and 172 predator-prey dyads.

Reference	Predator	Prey		Overlap (low)	CI	Overlap (high)	CI	Change in overlap
Baker & Leburg 2018	Bobcat (*Lynx rufus*)	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.641	0.464-0.851	0.806	0.718-0.981	0.165
		Desert Cottontail	*(Sylvilagus audubonii)*	0.758	0.652-0.97	0.814	0.728-0.982	0.056
	Coyote (*Canis latrans*)	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.664	0.486-0.933	0.879	0.827-0.993	0.215
		Deer spp.	*(Odocoileus hemionus, Odocoileus virginianus)*	0.517	0.312-0.752	0.757	0.624-0.86	0.24
		Desert Cottontail	*(Sylvilagus audubonii)*	0.815	0.714-1.137	0.767	0.67-0.841	-0.048
		Javelina	*(Tayassu tajacu)*	0.451	0.217-0.761	0.82	0.736-0.949	0.369
	Gray Fox (*Urocyon cinereoargenteus*)	Black-Tailed Jackrabbit	*(Lepus californicus)*	0.768	0.669-0.911	0.772	0.692-0.85	0.004
		Desert Cottontail	*(Sylvilagus audubonii)*	0.857	0.79-0.963	0.655	0.557-0.703	-0.202
Barrueto et al. 2014	Black Bear (*Ursus americanus*)	Mule Deer	*(Odocoileus hemionus)*	0.742	0.66-0.793	0.821	0.778-0.857	0.079
	Brown Bear (*Ursus arctos*)	Elk	*(Cervus elaphus)*	0.647	0.592-0.679	0.58	0.54-0.605	-0.067
		Moose	*(Alces alces)*	0.724	0.656-0.771	0.797	0.737-0.85	0.073
		Mule Deer	*(Odocoileus hemionus)*	0.834	0.792-0.866	0.852	0.817-0.886	0.018
	Puma (*Puma concolor*)	Mule Deer	*(Odocoileus hemionus)*	0.672	0.579-0.738	0.668	0.601-0.7	-0.004
	Coyote (*Canis latrans*)	Mule Deer	*(Odocoileus hemionus)*	0.871	0.836-0.917	0.84	0.806-0.861	-0.031
	Wolf (*Canis lupus*)	Elk	*(Cervus elaphus)*	0.865	0.824-0.904	0.804	0.771-0.823	-0.061
		Moose	*(Alces alces)*	0.884	0.835-0.949	0.888	0.839-0.939	0.004
		Mule Deer	*(Odocoileus hemionus)*	0.821	0.776-0.857	0.852	0.823-0.876	0.031
Caldwell & Klip 2020	Bobcat (*Lynx rufus*)	Lagomorph spp.	*(Sylvagus spp., Lepus californicus)*	0.276	0.104-0.429	0.641	0.544-0.713	0.365
	Coyote (*Canis latrans*)	Lagomorph spp.	*(Sylvagus spp., Lepus californicus)*	0.75	0.645-1.018	0.741	0.679-0.819	-0.009
		Mule Deer	*(Odocoileus hemionus)*	0.742	0.607-0.962	0.807	0.728-0.872	0.065
Carter et al. 2015	Leopard (*Panthera pardus*)	Wild Boar	*(Sus scrofa)*	0.767	0.687-0.909	0.808	0.712-0.924	0.041
		Chital	*(Axis axis)*	0.656	0.533-0.742	0.557	0.402-0.612	-0.099
		Muntjac	*(Muntiacus muntjak)*	0.597	0.443-0.69	0.571	0.415-0.645	-0.026
		Sambar	*(Rusa unicolor)*	0.492	0.351-0.604	0.841	0.774-0.964	0.349
	Tiger (*Panthera tigris*)	Wild Boar	*(Sus scrofa)*	0.481	0.387-0.52	0.71	0.6-0.807	0.229
		Chital	*(Axis axis)*	0.453	0.379-0.479	0.495	0.369-0.553	0.042
		Muntjac	*(Muntiacus muntjak)*	0.545	0.447-0.59	0.511	0.375-0.593	-0.034
		Sambar	*(Rusa unicolor)*	0.735	0.625-0.781	0.756	0.652-0.856	0.021
Dawson et al. 2018	Dingo (*Canis lupus dingo*)	Agile Wallaby	*(Macropus agilis)*	0.62	0.42-0.83	0.86	0.79-0.93	0.24
		Greater Bilby	*(Macrotis lagotis)*	0.3	0.08-0.46	0.66	0.57-0.75	0.36
	Domestic Cat (*Felis catus*)	Agile Wallaby	*(Macropus agilis)*	0.75	0.64-0.85	0.81	0.74-0.85	0.06
		Greater Bilby	*(Macrotis lagotis)*	0.55	0.42-0.68	0.68	0.58-0.78	0.13
Díaz-Ruíz et al. 2016	Red Fox (*Vulpes vulpes*)	European Rabbit	*(Oryctolagus cuniculus)*	0.425	-	0.3975	-	-0.0275
Gallo et al. 2019	Coyote (*Canis latrans*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.828	0.771-0.857	0.853	0.814-0.858	0.025
		White-Tailed Deer	*(Odocoileus virginianus)*	0.726	0.693-0.734	0.48	0.427-0.488	-0.246
Gray et al. 2011	Leopard (*Panthera pardus*)	Red Muntjac	*(Muntiacus muntjak)*	-	-	-	-	-
		Wild Boar	*(Sus scrofa)*	-	-	-	-	-

Kays et al. 2017	Black Bear (*Ursus americanus*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.405	0.325-0.433	0.581	0.483-0.648	0.176
		Eastern Fox Squirrel	*(Sciurus niger)*	0.63	0.57-0.669	0.548	0.448-0.588	-0.082
		Eastern Gray Squirrel	*(Sciurus carolinensis)*	0.697	0.664-0.729	0.549	0.489-0.582	-0.148
		Raccoon	*(Procyon lotor)*	0.33	0.273-0.333	0.419	0.343-0.433	0.089
		Virginia Opossum	*(Didelphis virginiana)*	0.29	0.212-0.288	0.339	0.225-0.345	0.049
		White-Tailed Deer	*(Odocoileus virginianus)*	0.801	0.767-0.825	0.825	0.781-0.859	0.024
		Wild Boar	*(Sus scrofa)*	0.634	0.495-0.746	0.743	0.632-0.832	0.109
	Bobcat (*Lynx rufus*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.627	0.539-0.709	0.728	0.634-0.804	0.101
		Eastern Fox Squirrel	*(Sciurus niger)*	0.486	0.387-0.554	0.438	0.335-0.473	-0.048
		Eastern Gray Squirrel	*(Sciurus carolinensis)*	0.529	0.445-0.612	0.417	0.354-0.445	-0.112
	Coyote (*Canis latrans*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.757	0.677-0.803	0.856	0.774-0.923	0.099
		Eastern Fox Squirrel	*(Sciurus niger)*	0.309	0.225-0.32	0.241	0.144-0.249	-0.068
		Eastern Gray Squirrel	*(Sciurus carolinensis)*	0.348	0.292-0.375	0.216	0.175-0.219	-0.132
		Raccoon	*(Procyon lotor)*	0.748	0.702-0.789	0.854	0.82-0.882	0.106
		Virginia Opossum	*(Didelphis virginiana)*	0.697	0.633-0.733	0.763	0.682-0.797	0.066
		White-Tailed Deer	*(Odocoileus virginianus)*	0.748	0.699-0.779	0.705	0.66-0.713	-0.043
		Wild Boar	*(Sus scrofa)*	0.723	0.585-0.817	0.71	0.609-0.784	-0.013
	Gray Fox (*Urocyon cinereoargenteus*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.863	0.806-0.974	0.824	0.727-0.914	-0.039
		Eastern Fox Squirrel	*(Sciurus niger)*	0.151	0.02-0.173	0.173	0.041-0.176	0.022
		Eastern Gray Squirrel	*(Sciurus carolinensis)*	0.18	0.069-0.22	0.148	0.054-0.159	-0.032
	Red Fox (*Vulpes vulpes*)	Eastern Cottontail	*(Sylvilagus floridanus)*	0.757	0.7-0.851	0.835	0.765-0.918	0.078
		Eastern Fox Squirrel	*(Sciurus niger)*	0.302	0.199-0.328	0.232	0.131-0.244	-0.07
		Eastern Gray Squirrel	*(Sciurus carolinensis)*	0.345	0.268-0.387	0.209	0.16-0.219	-0.136
Lendrum et al. 2017	Bobcat (*Lynx rufus*)	Lagomorph spp.	*(Sylvilagus nuttallii, Lepus californicus)*	0.55	-	0.794	-	0.244
	Coyote (*Canis latrans*)	Lagomorph spp.	*(Sylvilagus nuttallii, Lepus californicus)*	0.748	-	0.915	-	0.167
		Mule Deer	*(Odocoileus hemionus)*	0.838	-	0.84	-	0.002
Mills & Harris 2020	Spotted Hyena (*Crocuta crocuta*)	Aardvark	*(Orycteropus afer)*	0.656	-	0.796	-	0.14
		Buffalo	*(Syncerus caffer)*	0.763	-	0.777	-	0.014
		Bushbuck	*(Tragelaphus sylvaticus)*	0.726	-	0.779	-	0.053
		Duiker	*(Sylvicapra grimmia, Cephalophus rufilatus)*	0.776	-	0.778	-	0.002
		Hartebeest	*(Alcelaphus buselaphus)*	0.584	-	0.466	-	-0.118
		Kob	*(Kobus kob)*	0.428	-	0.312	-	-0.116
		Oribi	*(Ourebia ourebi)*	0.386	-	0.333	-	-0.053
		Reedbuck	*(Redunca redunca)*	0.78	-	0.813	-	0.033
		Roan Antelope	*(Hippotragus equinus)*	0.541	-	0.509	-	-0.032
		Warthog	*(Phacochoerus africanus)*	0.289	-	0.291	-	0.002
		Waterbuck	*(Kobus ellipsiprymnus)*	0.523	-	0.394	-	-0.129
	Leopard (*Panthera pardus*)	Aardvark	*(Orycteropus afer)*	0.776	-	0.74	-	-0.036
		Bushbuck	*(Tragelaphus sylvaticus)*	0.613	-	0.839	-	0.226
		Duiker spp.	*(Sylvicapra grimmia, Cephalophus rufilatus)*	0.661	-	0.795	-	0.134
		Hartebeest	*(Alcelaphus buselaphus)*	0.524	-	0.538	-	0.014
		Kob	*(Kobus kob)*	0.385	-	0.385	-	0
		Oribi	*(Ourebia ourebi)*	0.349	-	0.402	-	0.053
		Reedbuck	*(Redunca redunca)*	0.69	-	0.812	-	0.122
		Roan Antelope	*(Hippotragus equinus)*	0.467	-	0.581	-	0.114
		Warthog	*(Phacochoerus africanus)*	0.234	-	0.363	-	0.129
		Waterbuck	*(Kobus ellipsiprymnus)*	0.414	-	0.466	-	0.052
	Lion (*Panthera leo*)	Aardvark	*(Orycteropus afer)*	0.547	-	0.78	-	0.233
		Buffalo	*(Syncerus caffer)*	0.906	-	0.862	-	-0.044
		Bushbuck	*(Tragelaphus sylvaticus)*	0.897	-	0.858	-	-0.039
		Duiker spp.	*(Sylvicapra grimmia, Cephalophus rufilatus)*	0.831	-	0.849	-	0.018
		Hartebeest	*(Alcelaphus buselaphus)*	0.769	-	0.549	-	-0.22
		Kob	*(Kobus kob)*	0.608	-	0.391	-	-0.217
		Oribi	*(Ourebia ourebi)*	0.562	-	0.41	-	-0.152
		Reedbuck	*(Redunca redunca)*	0.899	-	0.874	-	-0.025
		Roan Antelope	*(Hippotragus equinus)*	0.715	-	0.592	-	-0.123
		Warthog	*(Phacochoerus africanus)*	0.462	-	0.375	-	-0.087
		Waterbuck	*(Kobus ellipsiprymnus)*	0.682	-	0.478	-	-0.204

Nix et al. 2018	Puma (*Puma concolor*)	Beaver	*(Castor canadensis)*	0.843	0.72-0.973	0.758	0.62-0.886	-0.085
		Mule Deer	*(Odocoileus hemionus)*	0.361	0.194-0.423	0.291	0.143-0.33	-0.07
		Raccoon	*(Procyon lotor)*	0.825	0.697-0.943	0.731	0.617-0.859	-0.094
Oberosler et al. 2017	Red Fox (*Vulpes vulpes*)	Brown Hare	*(Lepus europaeus)*	-	-	-	-	-
Reilly et al. 2017	Bobcat (*Lynx rufus*)	Brush Rabbit	*(Sylvilagus bachmani)*	0.632	0.496-0.707	0.576	0.434-0.623	-0.056
	Coyote (*Canis latrans*)	Brush Rabbit	*(Sylvilagus bachmani)*	0.764	0.654-0.87	0.619	0.496-0.691	-0.145
		Mule Deer	*(Odocoileus hemionus)*	0.841	0.773-0.909	0.784	0.728-0.829	-0.057
		Raccoon	*(Procyon lotor)*	0.638	0.511-0.725	0.724	0.653-0.767	0.086
	Gray Fox (*Urocyon cinereoargenteus*)	Brush Rabbit	*(Sylvilagus bachmani)*	0.44	0.293-0.485	0.373	0.237-0.405	-0.067
Rich et al. 2016	Black-backed Jackal (*Canis mesomelas*)	Cape Hare	*(Lepus capensis)*	0.849	0.753-0.932	0.793	0.737-0.846	-0.056
		Springhare	*(Pedetes capensis)*	0.9	0.839-1.085	0.685	0.598-0.738	-0.215
	Caracal (*Caracal caracal*)	Cape Hare	*(Lepus capensis)*	0.868	0.788-0.967	0.851	0.774-0.944	-0.017
		Springhare	*(Pedetes capensis)*	0.826	0.719-0.997	0.767	0.648-0.855	-0.059
	Leopard (*Panthera pardus*)	Common Duiker	*(Sylvicapra grimmia)*	0.724	0.612-0.843	0.744	0.65-0.82	0.02
		Greater Kudu	*(Tragelaphus strepsiceros)*	0.665	0.552-0.748	0.584	0.46-0.647	-0.081
		Impala	*(Aepyceros melampus)*	0.533	0.45-0.59	0.388	0.298-0.407	-0.145
		Warthog	*(Phacochoerus africanus)*	0.292	0.158-0.316	0.259	0.146-0.272	-0.033
	Lion (*Panthera leo*)	Buffalo	*(Bubalus bubalis)*	0.556	0.412-0.665	0.742	0.636-0.833	0.186
		Common Duiker	*(Sylvicapra grimmia)*	0.536	0.37-0.623	0.754	0.638-0.819	0.218
		Greater Kudu	*(Tragelaphus strepsiceros)*	0.487	0.367-0.573	0.608	0.485-0.682	0.121
		Impala	*(Aepyceros melampus)*	0.357	0.246-0.408	0.401	0.309-0.438	0.044
		Warthog	*(Phacochoerus africanus)*	0.186	0.075-0.236	0.255	0.13-0.274	0.069
		Zebra	*(Equus burchelli)*	0.489	0.364-0.568	0.619	0.492-0.677	0.13
	Spotted Hyena (*Crocuta crocuta*)	Buffalo	*(Bubalus bubalis)*	0.735	0.642-0.828	0.581	0.465-0.65	-0.154
		Common Duiker	*(Sylvicapra grimmia)*	0.531	0.393-0.603	0.564	0.44-0.597	0.033
		Greater Kudu	*(Tragelaphus strepsiceros)*	0.479	0.361-0.534	0.425	0.301-0.467	-0.054
		Impala	*(Aepyceros melampus)*	0.324	0.245-0.333	0.225	0.159-0.224	-0.099
		Warthog	*(Phacochoerus africanus)*	0.135	0.035-0.14	0.108	0.024-0.103	-0.027
		Zebra	*(Equus burchelli)*	0.517	0.419-0.567	0.476	0.371-0.528	-0.041
	Wild Dog (*Lycaon pictus*)	Cape Hare	*(Lepus capensis)*	0.433	0.29-0.461	0.4	0.301-0.446	-0.033
		Common Duiker	*(Sylvicapra grimmia)*	0.662	0.498-0.682	0.617	0.538-0.721	-0.045
		Greater Kudu	*(Tragelaphus strepsiceros)*	0.57	0.408-0.58	0.486	0.39-0.568	-0.084
		Impala	*(Aepyceros melampus)*	0.553	0.418-0.53	0.419	0.339-0.455	-0.134
		Springhare	*(Pedetes capensis)*	0.336	0.148-0.37	0.307	0.192-0.345	-0.029
		Warthog	*(Phacochoerus africanus)*	0.318	0.146-0.308	0.23	0.118-0.252	-0.088
		Zebra	*(Equus burchelli)*	0.628	0.489-0.645	0.426	0.33-0.501	-0.202
Smith et al. 2019	Black-backed Jackal (*Canis mesomelas*)	Blue Duiker	*(Philantomba monticola)*	0.74	0.59-0.86	0.72	0.62-0.81	-0.02
		Gray Duiker	*(Sylvicapra grimmia)*	0.66	0.48-0.83	0.66	0.55-0.76	0
	Caracal (*Caracal caracal*)	Blue Duiker	*(Philantomba monticola)*	0.82	0.75-0.87	0.69	0.54-0.80	-0.13
		Bushbuck	*(Tragelaphus sylvaticus)*	0.86	0.64-0.89	0.72	0.64-0.89	-0.14
		Gray Duiker	*(Sylvicapra grimmia)*	0.76	0.61-0.89	0.67	0.50-0.83	-0.09
Wang et al. 2015	Puma (*Puma concolor*)	Mule Deer	*(Odocoileus hemionus)*	-	-	-	-	-
		Virginia Opossum	*(Didelphus virginianus)*	0.659	0.554-0.730	0.771	0.680-0.850	0.112
		Raccoon	*(Procyon lotor)*	0.793	0.774-0.937	0.844	0.725-0.866	0.051
	Coyote (*Canis latrans*)	Mule Deer	*(Odocoileus hemionus)*	-	-	-	-	-
		Virginia Opossum	*(Didelphus virginianus)*	0.454	0.389-0.464	0.653	0.594-0.672	0.199
		Raccoon	*(Procyon lotor)*	0.545	0.485-0.557	0.739	0.681-0.781	0.194

Wilkinson et al. 2021	Black-backed Jackal (*Canis mesomelas*)	Kirk's Dikdik	*(Madoqua kirkii)*	0.523	0.407-0.561	0.65	0.55-0.69	0.127
		African Hare	*(Lepus victoriae)*	0.606	0.483-0.669	0.836	0.771-0.86	0.23
		Thomsons Gazelle	*(Eudorcas thomsonii)*	0.213	0.031-0.258	0.338	0.152-0.366	0.125
	Leopard (*Panthera pardus*)	Aardvark	*(Orycteropus afer)*	0.78	0.683-0.914	0.601	0.388-0.719	-0.179
		Kirk's Dikdik	*(Madoqua kirkii)*	0.332	0.158-0.347	0.511	0.298-0.603	0.179
		Impala	*(Aepyceros melampus)*	0.221	0.093-0.265	0.473	0.333-0.582	0.252
		Olive Baboon	*(Papio anubis)*	0.151	0.051-0.206	0.085	-0.219	-0.066
		Vervet	*(Chlorocebus pygerythrus)*	0.137	0.026-0.184	0.108	-0.233	-0.029
		Warthog	*(Phacochoerus africanus)*	0.202	0.066-0.241	0.103	-0.222	-0.099
		Waterbuck	*(Kobus ellipsiprymnus)*	0.36	0.134-0.471	0.439	0.26-0.551	0.079
		Common Zebra	*(Equus quagga)*	0.675	0.519-0.69	0.867	0.794-1.024	0.192
	Lion (*Panthera leo*)	Aardvark	*(Orycteropus afer)*	0.617	0.362-0.795	0.742	0.622-0.826	0.125
		Buffalo	*(Bubalus bubalis)*	0.575	0.308-0.635	0.579	0.517-0.632	0.004
		Kirk's Dikdik	*(Madoqua kirkii)*	0.12	-0.175	0.611	0.475-0.648	0.491
		Eland	*(Taurotragus oryx)*	0.256	0.112-0.307	0.381	0.22-0.365	0.125
		Impala	*(Aepyceros melampus)*	0.028	-0.101	0.406	0.293-0.39	0.378
		Olive Baboon	*(Papio anubis)*	0.001	-0.06	0.069	0.022-0.089	0.068
		Thomsons Gazelle	*(Eudorcas thomsonii)*	0.01	-0.151	0.295	0.097-0.305	0.285
		Warthog	*(Phacochoerus africanus)*	0.015	-0.1	0.09	0.029-0.103	0.075
		Waterbuck	*(Kobus ellipsiprymnus)*	0.27	0.052-0.445	0.464	0.327-0.546	0.194
		Common Zebra	*(Equus quagga)*	0.508	0.272-0.6	0.733	0.608-0.728	0.225
	Serval (*Leptailurus serval*)	African Hare	*(Lepus victoriae)*	0.736	0.622-0.793	0.756	0.614-0.881	0.02
	Spotted Hyena (*Crocuta crocuta*)	Aardvark	*(Orycteropus afer)*	0.674	0.526-0.73	0.664	0.529-0.731	-0.01
		Buffalo	*(Bubalus bubalis)*	0.799	0.757-0.799	0.671	0.649-0.675	-0.128
		Kirk's Dikdik	*(Madoqua kirkii)*	0.432	0.363-0.459	0.631	0.549-0.689	0.199
		Eland	*(Taurotragus oryx)*	0.481	0.403-0.516	0.304	0.203-0.307	-0.177
		Impala	*(Aepyceros melampus)*	0.254	0.205-0.246	0.381	0.325-0.351	0.127
		Olive Baboon	*(Papio anubis)*	0.129	0.076-0.109	0.018	0.004-0.011	-0.111
		Thomsons Gazelle	*(Eudorcas thomsonii)*	0.158	0.036-0.198	0.33	0.176-0.407	0.172
		Warthog	*(Phacochoerus africanus)*	0.247	0.19-0.243	0.045	0.019-0.036	-0.202
		Waterbuck	*(Kobus ellipsiprymnus)*	0.38	0.199-0.478	0.41	0.279-0.504	0.03
		Common Zebra	*(Equus quagga)*	0.821	0.77-0.829	0.81	0.776-0.809	-0.011
	Striped Hyena (*Hyaena hyaena*)	Olive Baboon	*(Papio anubis)*	0.117	0.048-0.101	0.02	-0.019	-0.097
Xiao et al. 2018	Tiger (*Panthera tigris*)	Roe Deer	*(Capreolus capreolus)*	0.84	0.76-0.92	0.75	0.68-0.82	-0.09
		Sika Deer	*(Cervus nippon)*	0.69	0.61-0.76	0.71	0.63-0.78	0.02
		Wild Boar	*(Sus scrofa)*	0.78	0.70-0.86	0.75	0.68-0.83	-0.03

References

1. Baker, A. D., & Leberg, P. L. (2018). Impacts of human recreation on carnivores in protected areas. PloS ONE, 13(4), e0195436.
2. Barrueto, M., Ford, A. T., & Clevenger, A. P. (2014). Anthropogenic effects on activity patterns of wildlife at crossing structures. Ecosphere, 5(3), 1–19.
3. Caldwell, M. R., & Klip, J. M. K. (2020). Wildlife interactions within highway underpasses. J. Wildl. Manag., 84(2), 227–236.
4. Carter, N., Jasny, M., Gurung, B., & Liu, J. (2015). Impacts of people and tigers on leopard spatiotemporal activity patterns in a global biodiversity hotspot. Glob. Ecol. Conserv., 3, 149–162.
5. Dawson, S. J., Adams, P. J., Moseby, K. E., Waddington, K. I., Kobryn, H. T., Bateman, P. W. et al. (2018). Peak hour in the bush: linear anthropogenic clearings funnel predator and prey species. Aust. Ecol., 43(2), 159–171.
6. Díaz‐Ruiz, F., Caro, J., Delibes‐Mateos, M., Arroyo, B., & Ferreras, P. (2016). Drivers of red fox (Vulpes vulpes) daily activity: prey availability, human disturbance or habitat structure? J. Zool., 298(2), 128–138.
7. Frid, A., & Dill, L. (2002). Human-caused Disturbance Stimuli as a Form of Predation Risk. Cons. Ecol., 6(1), 11–27.
8. Gallo, T., Fidino, M., Lehrer, E. W., & Magle, S. (2019). Urbanization alters predator‐avoidance behaviours. J. Anim. Ecol., 88(5), 793–803.
9. Gray, T. N., & Phan, C. (2011). Habitat preferences and activity patterns of the larger mammal community in Phnom Prich Wildlife Sanctuary, Cambodia. Raffles Bull. Zool., 59(2), 311– 318.
10. Kays, R., Parsons, A. W., Baker, M. C., Kalies, E. L., Forrester, T., Costello, R. et al. (2017). Does hunting or hiking affect wildlife communities in protected areas? J. Appl. Ecol., 54(1), 242–252.
11. Lendrum, P. E., Crooks, K. R., & Wittemyer, G. (2017). Changes in circadian activity patterns of a wildlife community post high-intensity energy development. J. Mammal., 98(5), 1265–1271.
12. Mills, K. L., & Harris, N. C. (2020). Humans induce differential access to prey for large African carnivores. bioRxiv. 10.1007/s11252-018-0758-6

13. Moll, R. J., Redilla, K. M., Mudumba, T., Muneza, A. B., Gray, S. M., Abade, L. et al. (2017). The many faces of fear: a synthesis of the methodological variation in characterizing predation risk. J. Anim. Ecol., 86(4), 749–765.
14. Nix, J. H., Howell, R. G., Hall, L. K., & McMillan, B. R. (2018). The influence of periodic increases of human activity on crepuscular and nocturnal mammals: Testing the weekend effect. Behav. Processes, 146, 16–21.
15. Nouvellet, P., Rasmussen, G. S. A., Macdonald, D. W., & Courchamp, F. (2012). Noisy clocks and silent sunrises: measurement methods of daily activity pattern. J. Zool., 286(3), 179–184.
16. Oberosler, V., Groff, C., Iemma, A., Pedrini, P., & Rovero, F. (2017). The influence of human disturbance on occupancy and activity patterns of mammals in the Italian Alps from systematic camera trapping. Mammal. Biol., 87(1), 50–61.
17. Parr, C. S.,Wilson N., Leary P., Schulz, K. S., Lans, K., Walley, L. et al. (2014). The Encyclopedia of Life v2: Providing Global Access to Knowledge About Life on Earth. Biodivers. Data J., 2, e1079.
18. Reilly, M. L., Tobler, M. W., Sonderegger, D. L., & Beier, P. (2017). Spatial and temporal response of wildlife to recreational activities in the San Francisco Bay ecoregion. Biol. Conserv., 207, 117–126.
19. Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera trap data. J. Agric. Biol. Environ. Stat., 14(3), 322–337.
20. Rich, L. N., Miller, D. A., Robinson, H. S., McNutt, J. W., & Kelly, M. J. (2016). Using camera trapping and hierarchical occupancy modeling to evaluate the spatial ecology of an African mammal community. J. Appl. Ecol., 53(4), 1225–1235.
21. Smith, Y. C. E., Smith, D. A. E., Ramesh, T., & Downs, C. T. (2019). Novel predators and anthropogenic disturbance influence spatio-temporal distribution of forest antelope species. Behav. Processes, 159, 9–22.
22. Sollmann, R. (2018). A gentle introduction to camera‐trap data analysis. Afr. J. Ecol., 56(4), 740-749.
23. Wang, Y., Allen, M. L., & Wilmers, C. C. (2015). Mesopredator spatial and temporal responses to large predators and human development in the Santa Cruz Mountains of California. Biol. Conserv., 190, 23–33.

24. Wilkinson, C. E., McInturff, A., Kelly, M., & Brashares, J. S. (2021). Quantifying wildlife responses to conservation fencing in East Africa. Biol. Conserv., 256, 109071.
25. Xiao, W., Hebblewhite, M., Robinson, H., Feng, L., Zhou, B., Mou, P. et al. (2018). Relationships between humans and ungulate prey shape Amur tiger occurrence in a core protected area along the Sino‐Russian border. Ecol. Evol., 8(23), 11677–11693.