

R401: Statistical and Mathematical Foundations

Lecture 19: Infinite-Horizon Deterministic Optimal Control in Discrete Time

Andrey Vassilev

2016/2017

Lecture Contents

- 1 Introduction
- 2 Problems in variational form
- 3 Problems with explicit controls

Introduction

Switching to discrete time

- You are already familiar with a number of dynamic optimization problems in which time is continuous.
- In many applications, however, it is natural to work in discrete time.
- This provides a bridge to validating/calibrating models with data or estimating them, among others.
- We therefore need to develop the counterpart of the continuous-time optimal control framework for the case of discrete time.

Switching to discrete time

- Some of the details of such a transition are easily predictable:
 - Differential equations will be replaced by difference equations.
 - The objective functional will involve a series instead of an integral.
- Other details need to be specified further. In particular, there exist two broad classes of dynamic optimization problems in discrete time:
 - Problems in variational form
 - Problems with explicit controls
- Both classes can be used to address a wide variety of problems.
- However, problems with explicit controls are a bit more transparent in terms of their structure.

Specific assumptions

- We will work in an infinite-horizon setup. Finite-horizon formulations for discrete-time problems exist but are less common in economic applications.
- We sacrifice some generality from the outset by assuming a specific structure of the problems:
 - Special (time) discounting in the objective functionals.
 - Autonomous difference equations describing the evolution of the system that is being modelled.

Problems in variational form

Formulation

The Euler equation

Sufficiency

Example: NCs for a problem in variational form

Compute the NCs for the problem:

$$\max_{\{c_t\}} \sum_{t=0}^{\infty} \beta^t \ln c_t$$

$$k_{t+1} = (1 - \delta)k_t + y_t - c_t, \quad k_0 > 0 \text{ - given}$$

$$y_t = Ak_t^\alpha, \quad 0 < \underline{c} \leq c_t \leq \bar{c}, \quad \alpha, \beta \in (0, 1), \quad A > 0$$

NCs:

$$\sum_{t=0}^{\infty} \beta^t \underbrace{\ln((1 - \delta)k_t + Ak_t^\alpha - k_{t+1})}_{=F(x_t, x_{t+1})}.$$

Euler equation:

$$F_y(x_t, x_{t+1}) + \beta F_x(x_{t+1}, x_{t+2}) = 0$$

Example: NCs for a problem in variational form

Applying the Euler equation, we get:

$$\frac{-1}{(1-\delta)k_t + Ak_t^\alpha - k_{t+1}} + \beta \frac{(1-\delta) + \alpha Ak_{t+1}^{\alpha-1}}{(1-\delta)k_{t+1} + Ak_{t+1}^\alpha - k_{t+2}} = 0$$

The last result has the form

$$G(x_{t+2}, x_{t+1}, x_t) = 0,$$

i.e. a nonlinear second-order difference equation.

Example: NCs for a problem in variational form

Applying the Euler equation, we get:

$$\frac{-1}{(1-\delta)k_t + Ak_t^\alpha - k_{t+1}} + \beta \frac{(1-\delta) + \alpha Ak_{t+1}^{\alpha-1}}{(1-\delta)k_{t+1} + Ak_{t+1}^\alpha - k_{t+2}} = 0$$

The last result has the form

$$G(x_{t+2}, x_{t+1}, x_t) = 0,$$

i.e. a nonlinear second-order difference equation.

Question: Can you compute the solution of the equation for $\beta = 0.95$, $A = 1$, $\delta = 0.05$, $\alpha = 0.5$ and $k_0 = 2$?

Example: NCs for a problem in variational form

Applying the Euler equation, we get:

$$\frac{-1}{(1-\delta)k_t + Ak_t^\alpha - k_{t+1}} + \beta \frac{(1-\delta) + \alpha Ak_{t+1}^{\alpha-1}}{(1-\delta)k_{t+1} + Ak_{t+1}^\alpha - k_{t+2}} = 0$$

The last result has the form

$$G(x_{t+2}, x_{t+1}, x_t) = 0,$$

i.e. a nonlinear second-order difference equation.

Question: Can you compute the solution of the equation for $\beta = 0.95$, $A = 1$, $\delta = 0.05$, $\alpha = 0.5$ and $k_0 = 2$? If not, what else do you need?

Problems with explicit controls

Formulation of the problem with explicit controls

$X \subset \mathbb{R}^n$ – *state space*, the state variables are $x = (x^1, \dots, x^n)$.

We assume that $\forall x \in X, \exists U(x) \subset \mathbb{R}^m, U(x) \neq \emptyset$. The elements of $u = (u^1, \dots, u^m)$ are our *controls*.

Objective function (instantaneous): $f^0(x, u) \quad x \in X, u \in U(x)$

State equations:

$$x_{t+1} = f(x_t, u_t), \quad x_0 - \text{given} \quad (1)$$

where $f(x, u)$ is a vector function taking values in X , for $x \in X, u \in U(x)$.

Formulation of the problem with explicit controls

We need to find a sequence of admissible controls $\mathbf{u} = \{u_t\}, t = 0, 1, 2, \dots$, which determine a sequence of state variables $\{x_{t+1}\}, t = 0, 1, 2, \dots$ via (1) for which

$$j(x_0, \mathbf{u}) = \sum_{t=0}^{\infty} \beta^t f^0(x_t, u_t) \quad (2)$$

attains a maximum

$$v(x_0) = \sup_{\mathbf{u}} j(x_0, \mathbf{u}). \quad (3)$$

We refer to the above problem as *Problem A*.

Formulation of the problem with explicit controls

The number $\beta \in (0, 1)$ is the *discount factor* in the model.

Denote by $\text{FC}(x_0)$ the set of all feasible control sequences $\{u_t\}_{t=0}^{\infty}$ for initial $x_0 \in X$, i.e. x_{t+1} satisfies (1) for $u_t \in U(x_t)$, $t = 0, 1, 2, \dots$, and a given x_0 .

We denote the optimal sequence of pairs of state variables and controls for Problem A by $\{x_{t+1}^*, u_t^*\}$, $t = 0, 1, 2, \dots$, i.e. $\{u_t^*\} \in \text{FC}(x_0)$, and

$$v(x_0) = j(x_0, \mathbf{u}^*), \quad \mathbf{u}^* = \{u_t^*\}.$$

Necessary conditions for optimality

A variation of the familiar approach for problems of this type is the following:

- 1 Construct the Lagrangian

$$\mathcal{L}(x_1, x_2, \dots, u_0, u_1, \dots) = \sum_{t=0}^{\infty} \beta^t \left[f^0(x_t, u_t) + \lambda'_t \cdot [f(x_t, u_t) - x_{t+1}] \right],$$

where $\lambda_t = (\lambda_t^1, \dots, \lambda_t^n)$, $t = 0, 1, 2, \dots$, are the Lagrange multipliers and the dot (\cdot) denotes a matrix product or a scalar product:

$$\lambda'_t \cdot [f(x_t, u_t) - x_{t+1}] = \sum_{i=1}^n \lambda_t^i [f^i(x_t, u_t) - x_{t+1}^i].$$

Necessary conditions for optimality

- ② Differentiate \mathcal{L} w.r.t. x_t and u_t , set the resulting expressions equal to zero and obtain first-order necessary conditions for optimality:

$$\beta \left[f_{x_t^k}^0(x_t, u_t) + \sum_{i=1}^n \lambda_t^i f_{x_t^k}^i(x_t, u_t) \right] = \lambda_{t-1}^k, \quad k = 1, \dots, n, \quad (4)$$

$$f_{u_t}^0(x_t, u_t) + \sum_{i=1}^n \lambda_t^i f_{u_t}^i(x_t, u_t) = 0, \quad j = 1, \dots, m.$$

- ③ Equations (1) and (4) are solved as a system and we obtain a candidate solution $\{u_t\}_{t=0}^\infty$ or, more precisely, a sequence $\{x_{t+1}, u_t\}_{t=0}^\infty$. (It is common to find a stationary point of the system (1) and (4), and work with a linearised version of the system around that point.)

Necessary conditions for optimality

In matrix notation the above takes that form:

$$\begin{aligned}\mathcal{L}_x &= \beta^t f_x^0(x_t, u_t) + \beta^t f'_x(x_t, u_t) \cdot \lambda_t - \beta^{t-1} \lambda_{t-1} = 0 \Rightarrow \\ &\beta(f_x^0(x_t, u_t) + f'_x(x_t, u_t) \cdot \lambda_t) = \lambda_{t-1}.\end{aligned}\tag{5}$$

$$\begin{aligned}\mathcal{L}_u &= \beta^t f_u^0(x_t, u_t) + \beta^t f'_u(x_t, u_t) \cdot \lambda_t = 0 \Rightarrow \\ &f_u^0(x_t, u_t) + f'_u(x_t, u_t) \cdot \lambda_t = 0\end{aligned}\tag{6}$$

Necessary conditions for optimality

Why is this algorithm valid?

The value function for Problem A satisfies a version of the Bellman equation:

$$v(x) = \sup_{u \in U(x)} \left[f^0(x, u) + \beta v(f(x, u)) \right]. \quad (7)$$

Let the supremum in (7) be attained on the interior of the set $U(x)$. Denote this point by $u = v(x)$ and assume that all objects used below are differentiable.

Necessary conditions for optimality

We have

$$v(x) = f^0(x, v(x)) + \beta v(f(x, v(x))). \quad (8)$$

Also the extremum condition is

$$f_u^0(x, v(x)) + \beta f_u'(x, v(x)) \cdot v_x(f(x, v(x))) = 0. \quad (9)$$

Differentiating (8) w.r.t. x , we obtain

$$\begin{aligned} v_x(x) &= f_x^0(x, v(x)) + v'_x(x) \cdot f_u^0(x, v(x)) + \\ &\quad \beta [f'_x(x, v(x)) + v'_x(x) \cdot f'_u(x, v(x))] \cdot v_x(f(x, v(x))) \\ &= f_x^0(x, v(x)) + \beta f'_x(x, v(x)) \cdot v_x(f(x, v(x))) + \\ &\quad \underbrace{v'_x(x) \cdot f_u^0(x, v(x)) + \beta v'_x(x) \cdot f'_u(x, v(x)) \cdot v_x(f(x, v(x)))}_{=0 \text{ in view of (9)}}. \end{aligned}$$

Necessary conditions for optimality

We thus get

$$v_x(x) = f_x^0(x, v(x)) + \beta f'_x(x, v(x)) \cdot v_x(f(x, v(x))). \quad (10)$$

For $x = x_t^*$ and $u_t^* = v(x_t^*)$, equations (9) and (10) take the form

$$f_u^0(x_t^*, u_t^*) + \beta f'_u(x_t^*, u_t^*) \cdot v_x(x_{t+1}^*) = 0, \quad (11)$$

$$v_x(x_t^*) = f_x^0(x_t^*, u_t^*) + \beta f'_x(x_t^*, u_t^*) \cdot v_x(x_{t+1}^*). \quad (12)$$

Set $\lambda_t := \beta v_x(x_{t+1}^*)$ in (11) and (12), to obtain precisely (5) and (6).

Sufficient conditions

Fact 1

Let $\{\lambda_t\}$ and $\{x_{t+1}^*, u_t^*\}$, $t = 0, 1, 2, \dots$, be obtained by using (1) and (4). If

- ① The functions $f^0(x, u)$ and $f(x, u)$ are concave in (x, u) ,
- ② The Lagrange multipliers $\lambda_t^1, \dots, \lambda_t^n$, $t = 0, 1, 2, \dots$ are nonnegative,
- ③ The state space X is a subset of \mathbb{R}_+^n and the following transversality condition is valid

$$\lim_{T \rightarrow \infty} \beta^T \lambda'_T \cdot x_{T+1}^* = 0,$$

then the sequence $\{x_{t+1}^*, u_t^*\}$ (for a given x_0) is optimal for Problem A.

Sufficient conditions

We shall verify the validity of Fact 1.

Recall that (4) in matrix terms is given by (5) and (6).
Consider

$$\mathcal{L}_T(x_t, u_t) = \sum_{t=0}^T \beta^t \left\{ f^0(x_t, u_t) + \lambda'_t \cdot [f(x_t, u_t) - x_{t+1}] \right\}.$$

We have

$$\begin{aligned} D := \mathcal{L}_T(x_t, u_t) - \mathcal{L}_T(x_t^*, u_t^*) &= \sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \\ &\sum_{t=0}^T \beta^t [f^0(x_t, u_t) + \lambda'_t \cdot f(x_t, u_t) - f^0(x_t^*, u_t^*) - \lambda'_t \cdot f(x_t^*, u_t^*)]. \end{aligned} \quad (13)$$

Sufficient conditions

Then, in view of concavity, we get

$$\begin{aligned}
 \mathcal{L}_T(x_t, u_t) - \mathcal{L}_T(x_t^*, u_t^*) \quad (= D) &\leq \sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \\
 &\sum_{t=0}^T \beta^t [f_x^{0'}(x_t^*, u_t^*) \cdot (x_t - x_t^*) + f_u^{0'}(x_t^*, u_t^*) \cdot (u_t - u_t^*) + \\
 &\lambda'_t \cdot [f_x(x_t^*, u_t^*) \cdot (x_t - x_t^*) + f_u(x_t^*, u_t^*) \cdot (u_t - u_t^*)]] = \\
 &\sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \sum_{t=0}^T \beta^t \underbrace{[f_x^{0'}(x_t^*, u_t^*) + \lambda'_t \cdot f_x(x_t^*, u_t^*)]}_{=\frac{\lambda'_{t-1}}{\beta} \text{ in view of (5)}} \cdot (x_t - x_t^*) \\
 &+ \sum_{t=0}^T \beta^t \underbrace{[f_u^{0'}(x_t^*, u_t^*) + \lambda'_t \cdot f_u(x_t^*, u_t^*)]}_{=0' \text{ in view of (6)}} \cdot (u_t - u_t^*).
 \end{aligned}$$

Sufficient conditions

i.e.

$$\begin{aligned}
 D &\leq \sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \sum_{t=0}^T \beta^t \frac{\lambda'_{t-1}}{\beta} \cdot \underbrace{(x_t - x_t^*)}_{\text{N.B.: } x_0 = x_0^*} = \\
 &\sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \sum_{t=1}^T \beta^{t-1} \frac{\lambda'_{t-1}}{\beta} \cdot (x_t - x_t^*) = \\
 &\sum_{t=0}^T \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \sum_{t=0}^{T-1} \beta^t \frac{\lambda'_t}{\beta} \cdot (x_{t+1} - x_{t+1}^*) = \\
 &\underbrace{\sum_{t=0}^{T-1} \beta^t \lambda'_t \cdot (x_{t+1}^* - x_{t+1}) + \sum_{t=0}^{T-1} \beta^t \frac{\lambda'_t}{\beta} \cdot (x_{t+1} - x_{t+1}^*)}_{=0} + \\
 &\beta^T \lambda'_T \cdot (x_{T+1}^* - x_{T+1}) \underbrace{\leq}_{\text{since } \lambda_t, x_t \geq 0} \beta^T \lambda'_T \cdot x_{T+1}^*.
 \end{aligned}$$

Sufficient conditions

In view of the transversality condition, we have:

$$D \leq \beta^T \lambda'_T \cdot x_{T+1}^* \xrightarrow{T \rightarrow \infty} 0,$$

i.e.

$$\mathcal{L}_T(x_t^*, u_t^*) - \mathcal{L}_T(x_t, u_t) \geq 0,$$

which proves the optimality of the sequence $\{x_{t+1}^*, u_t^*\}$.

Homework

Readings

Main references: Additional readings:

Sydsæter et al. [SHSS] *Further mathematics for economic analysis*. Chapter 12.