Bayesian Inference in Astrophysics: Stellar X-Ray Flare Rates and Exoplanet Mass-Radius Scaling Laws

Tanishq Hindolya, Faaz Naushad, Sharath Kumar

Contents

1 Background, Motivation, and Astrophysical Context **1**

2 Exponential Families, Conjugacy, and Prior Choice: Stellar Flare Model **2**

2.1 Canonical Exponential Family Representation and Conjugate Prior 2

2.2 Non-Informative Jeffreys Prior Derivation 3

2.3 Bayesian Estimation and Hypothesis Testing via Bayes Factors 3

3 Bayesian Linear Models, MCMC Sampling, and Variable Selection **4**

3.1 Mathematical Formulation of Exoplanet Mass Scaling 4

4 Empirical Results and MCMC Convergence Diagnostics **6**

4.1 Discussion of Astrophysical Findings 7

Code and Reproducibility Repository: The complete R source code (`exoplanet_bayesian_project.R`), data pipeline, and R Markdown source for this report are publicly hosted on GitHub at: https://github.com/avasterinbloom/bayesian-astrophysics-project

---

1 Background, Motivation, and Astrophysical Context

Bayesian inference provides a coherent probabilistic framework for combining mechanistic physical laws (encoded via prior distributions $p(\boldsymbol{\theta})$) with observational telemetry (encoded via the likelihood $p(\mathbf{y} \mid \boldsymbol{\theta})$) through Bayes' Theorem[cite: 1]:

$$p(\boldsymbol{\theta} \mid \mathbf{y}) = \frac{p(\mathbf{y} \mid \boldsymbol{\theta})p(\boldsymbol{\theta})}{\int_{\Theta} p(\mathbf{y} \mid \boldsymbol{\theta}')p(\boldsymbol{\theta}')\, d\boldsymbol{\theta}'} \propto L(\boldsymbol{\theta}; \mathbf{y})p(\boldsymbol{\theta})$$

In observational astrophysics, sample sizes are frequently constrained by telescope allocation time, instrumental detection thresholds, and severe physical multicollinearity. This project investigates two foundational problems in space[cite: 1]:

1. **High-Energy Stellar X-Ray Flare Arrival Rates (Discrete Exponential Family):** Modeling monthly magnetic reconnection flare counts $Y_i \sim \text{Poisson}(\lambda)$ using conjugate Gamma priors, non-informative Jeffreys priors, and exact Bayes Factors to test whether a host star has transitioned into an active magnetic cycle[cite: 1].
2. **Exoplanet Mass-Radius Scaling and Keplerian Multicollinearity (Continuous Multi-Parameter Linear Models):** Using confirmed exoplanet telemetry from the **NASA Exoplanet Archive**, we model planetary log-mass ($\log_{10} M_p$) as a function of $p = 10$ planetary and host-star observables. Because Kepler's Third Law ($P^2 \propto a^3/M_*$) and the Stefan-Boltzmann radiative equilibrium law ($T_{\text{eq}} \propto T_{\text{eff}}\sqrt{R_*/a}$) induce strong collinearity among predictors, ordinary least squares (and unregularized non-informative priors) suffer from variance inflation. We resolve this by deriving the **Conjugate Normal-Inverse-Gamma (NIG)** posterior, implementing a **Gibbs Sampler** with a **Metropolis-Hastings** hyperparameter step for **Bayesian LASSO**, and performing **Spike-and-Slab variable selection**[cite: 1].

---

2 Exponential Families, Conjugacy, and Prior Choice: Stellar Flare Model

2.1 Canonical Exponential Family Representation and Conjugate Prior

Let $Y_1, \ldots, Y_n \overset{\text{i.i.d.}}{\sim} \text{Poisson}(\lambda)$ denote the monthly count of solar/stellar X-ray flares. The joint likelihood of $\mathbf{y} = (y_1, \ldots, y_n)^T$ can be written in canonical one-parameter exponential family form[cite: 1]:

$$p(\mathbf{y} \mid \lambda) = \left(\prod_{i=1}^{n} \frac{1}{y_i!}\right) \exp\left(\log(\lambda) \sum_{i=1}^{n} y_i - n\lambda\right) = h(\mathbf{y}) \exp\big(\eta(\lambda) T(\mathbf{y}) - nA(\lambda)\big)$$

where the natural parameter is $\eta(\lambda) = \log \lambda$, the sufficient statistic is $T(\mathbf{y}) = \sum_{i=1}^{n} y_i$, and the log-partition function is $A(\lambda) = \lambda$.

Proposition 1 (Gamma-Poisson Conjugacy)**.** *If the prior on λ follows Gamma(α_0, β_0) with density $p(\lambda) = \frac{\beta_0^{\alpha_0}}{\Gamma(\alpha_0)} \lambda^{\alpha_0 - 1} e^{-\beta_0 \lambda}$, the posterior distribution $\lambda \mid \mathbf{y}$ is closed under sampling and follows[cite: 1]:*

$$\lambda \mid \mathbf{y} \sim \textit{Gamma}(\alpha_n, \beta_n), \quad \textit{where } \alpha_n = \alpha_0 + \sum_{i=1}^{n} y_i, \quad \beta_n = \beta_0 + n.$$

Proof. Multiplying the likelihood and prior kernel with respect to λ yields:

$$p(\lambda \mid \mathbf{y}) \propto \left(\lambda^{\sum_{i=1}^{n} y_i} e^{-n\lambda}\right) \left(\lambda^{\alpha_0 - 1} e^{-\beta_0 \lambda}\right) = \lambda^{\left(\alpha_0 + \sum_{i=1}^{n} y_i\right) - 1} \exp\big(-(\beta_0 + n)\lambda\big),$$

which is the exact kernel of Gamma(α_n, β_n). □

2.2 Non-Informative Jeffreys Prior Derivation

To ensure invariance under smooth one-to-one reparameterizations $\phi = g(\lambda)$, we derive **Jeffreys' prior** $p_J(\lambda) \propto \sqrt{\mathcal{I}(\lambda)}$[cite: 1]. Since $\log p(y \mid \lambda) = y \log \lambda - \lambda - \log(y!)$, the second derivative is $\frac{\partial^2}{\partial \lambda^2} \log p(y \mid \lambda) = -\frac{y}{\lambda^2}$. Taking the negative expectation yields the Fisher Information:

$$\mathcal{I}(\lambda) = -\mathbb{E}\left[-\frac{Y}{\lambda^2}\right] = \frac{\lambda}{\lambda^2} = \frac{1}{\lambda} \implies p_J(\lambda) \propto \lambda^{-1/2}.$$

Observing the kernel $\lambda^{1/2-1} e^{-0 \cdot \lambda}$, Jeffreys' improper prior corresponds to the limiting parameters $(\alpha_0, \beta_0) = (0.5, 0)$, yielding a proper posterior Gamma($\sum y_i + 0.5$, n) whenever $\sum y_i \geq 0$ and $n \geq 1$.

2.3 Bayesian Estimation and Hypothesis Testing via Bayes Factors

Under $\lambda \mid \mathbf{y} \sim \text{Gamma}(\alpha_n, \beta_n)$[cite: 1]: * **Posterior Mean (Squared Error Loss):** $\hat{\lambda}_{\text{MMSE}} = \mathbb{E}[\lambda \mid \mathbf{y}] = \frac{\alpha_n}{\beta_n}$. * **Maximum A Posteriori (MAP) Estimate:** Differentiating $\log p(\lambda \mid \mathbf{y}) = (\alpha_n - 1) \log \lambda - \beta_n \lambda + C$ gives $\hat{\lambda}_{\text{MAP}} = \frac{\alpha_n - 1}{\beta_n}$ for $\alpha_n \geq 1$[cite: 1]. * **Bayesian Hypothesis Test:** To test whether the star has entered a high-activity state $H_1 : \lambda > \lambda_{\text{crit}}$ against quiet-star baseline $H_0 : \lambda \leq \lambda_{\text{crit}}$ ($\lambda_{\text{crit}} = 7.5$ flares/month), we compute the **Posterior Odds** and **Bayes Factor** (BF_{10})[cite: 1]:

$$\text{Posterior Odds}_{10} = \frac{P(\lambda > \lambda_{\text{crit}} \mid \mathbf{y})}{P(\lambda \leq \lambda_{\text{crit}} \mid \mathbf{y})}, \qquad BF_{10} = \frac{\text{Posterior Odds}_{10}}{\text{Prior Odds}_{10}} = \frac{P(\lambda > \lambda_{\text{crit}} \mid \mathbf{y}) / P(\lambda \leq \lambda_{\text{crit}} \mid \mathbf{y})}{P(\lambda > \lambda_{\text{crit}}) / P(\lambda \leq \lambda_{\text{crit}})}.$$

Table 1: Bayesian point and interval estimates for stellar X-ray flare rate.

Prior Specification	Posterior Mean	MAP Estimate	95% CI Lower	95% CI Upper
Conjugate Gamma(10, 2)	8.2692	8.2308	7.2008	9.4105
Jeffreys Prior (lambdaˆ-1/2)	8.5625	8.5208	7.4318	9.7721




Figure 1: Prior sensitivity analysis for stellar X-ray flare arrival rate. Comparison of the historical quiet-star Conjugate Gamma(10, 2) prior, Conjugate Posterior, and Jeffreys Non-Informative Posterior.

For $H_1 : \lambda > 7.5$ vs. $H_0 : \lambda \leq 7.5$, the **Posterior Odds** are **11.06** and the **Bayes Factor** is $BF_{10} = 147.31$, providing decisive evidence on Jeffreys' scale that the star is in an elevated magnetic flare state.

---

3 Bayesian Linear Models, MCMC Sampling, and Variable Selection

3.1 Mathematical Formulation of Exoplanet Mass Scaling

We model standardized log-planetary mass $\mathbf{y} \in \mathbb{R}^n$ ($\log_{10} M_p/M_\oplus$) across n confirmed exoplanets using $p = 10$ standardized astrophysical covariates $\mathbf{X} \in \mathbb{R}^{n\times p}$:

$$\mathbf{y} \mid \boldsymbol{\beta}, \sigma^2 \sim \mathcal{N}_n(\mathbf{X}\boldsymbol{\beta}, \sigma^2\mathbf{I}_n).$$

3.1.1 Conjugate Normal-Inverse-Gamma (NIG) Prior and Ridge Equivalence

Specifying the conjugate NIG($\boldsymbol{\mu}_0, \mathbf{V}_0, a_0, b_0$) prior[cite: 1]:

$$\boldsymbol{\beta} \mid \sigma^2 \sim \mathcal{N}_p(\boldsymbol{\mu}_0, \sigma^2\mathbf{V}_0), \qquad \sigma^2 \sim \text{Inv-Gamma}(a_0, b_0),$$

and completing the quadratic form in $\boldsymbol{\beta}$ yields the exact joint posterior $\text{NIG}(\boldsymbol{\mu}_n, \mathbf{V}_n, a_n, b_n)$:

$$\mathbf{V}_n = \left(\mathbf{V}_0^{-1} + \mathbf{X}^T\mathbf{X}\right)^{-1}, \qquad \boldsymbol{\mu}_n = \mathbf{V}_n\left(\mathbf{V}_0^{-1}\boldsymbol{\mu}_0 + \mathbf{X}^T\mathbf{y}\right),$$

$$a_n = a_0 + \frac{n}{2}, \qquad b_n = b_0 + \frac{1}{2}\left(\mathbf{y}^T\mathbf{y} + \boldsymbol{\mu}_0^T\mathbf{V}_0^{-1}\boldsymbol{\mu}_0 - \boldsymbol{\mu}_n^T\mathbf{V}_n^{-1}\boldsymbol{\mu}_n\right).$$

Setting $\boldsymbol{\mu}_0 = \mathbf{0}$ and $\mathbf{V}_0 = \lambda^{-1}\mathbf{I}_p$ yields $\boldsymbol{\mu}_n = (\mathbf{X}^T\mathbf{X} + \lambda\mathbf{I}_p)^{-1}\mathbf{X}^T\mathbf{y}$, proving that **Ridge Regression is the exact MAP/posterior mean solution** under an isotropic Gaussian prior[cite: 1]. Integrating out σ^2 yields the exact marginal Multivariate Student-t posterior $\boldsymbol{\beta} \mid \mathbf{y} \sim t_{2a_n}\left(\boldsymbol{\mu}_n, \frac{b_n}{a_n}\mathbf{V}_n\right)$. Under **Jeffreys' non-informative prior** $p_J(\boldsymbol{\beta}, \sigma^2) \propto (\sigma^2)^{-1}$ ($\mathbf{V}_0^{-1} \to \mathbf{0}$), the posterior mean reduces to $\hat{\boldsymbol{\beta}}_{\text{OLS}} = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{y}$[cite: 1].

3.1.2 Bayesian LASSO via Scale-Mixture Gibbs Sampling and Metropolis-Hastings

Placing an independent Laplace prior on each regression weight $p(\boldsymbol{\beta} \mid \sigma^2) = \prod_{j=1}^{p} \frac{\lambda}{2\sqrt{\sigma^2}} \exp\left(-\frac{\lambda|\beta_j|}{\sqrt{\sigma^2}}\right)$ yields the **LASSO** estimator as the posterior mode[cite: 1]. Following Park and Casella (2008), we represent the Laplace density as a scale mixture of normals with latent variances $\tau_1^2, \ldots, \tau_p^2$:

$$\beta_j \mid \sigma^2, \tau_j^2 \sim \mathcal{N}(0, \sigma^2\tau_j^2), \qquad \tau_j^2 \mid \lambda \sim \text{Exponential}\left(\frac{\lambda^2}{2}\right).$$

This hierarchical expansion enables a closed-form **Gibbs Sampler**[cite: 1]: 1. $\boldsymbol{\beta} \mid \sigma^2, \boldsymbol{\tau}^2, \mathbf{y} \sim \mathcal{N}_p\left(\mathbf{A}^{-1}\mathbf{X}^T\mathbf{y},\, \sigma^2\mathbf{A}^{-1}\right)$, where $\mathbf{A} = \mathbf{X}^T\mathbf{X} + \text{diag}(\tau_1^{-2}, \ldots, \tau_p^{-2})$. 2. $\sigma^2 \mid \boldsymbol{\beta}, \boldsymbol{\tau}^2, \mathbf{y} \sim \text{Inv-Gamma}\left(\frac{n-1+p}{2},\, \frac{1}{2}\|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \frac{1}{2}\sum_{j=1}^{p}\frac{\beta_j^2}{\tau_j^2}\right)$. 3. $\tau_j^{-2} \mid \beta_j, \sigma^2, \lambda \sim \text{Inv-Gaussian}\left(\mu' = \sqrt{\frac{\lambda^2\sigma^2}{\beta_j^2}},\, \lambda' = \lambda^2\right)$. 4. **Metropolis-Hastings Update for λ:** Assigning a hyperprior $\lambda \sim \text{Gamma}(a_\lambda, b_\lambda)$ and proposing $\log\lambda^* \sim \mathcal{N}(\log\lambda^{(t)}, s_{\text{prop}}^2)$, we accept λ^* with probability $\alpha(\lambda^{(t)}, \lambda^*) = \min\left(1, \frac{p(\lambda^*|\boldsymbol{\tau}^2)\lambda^*}{p(\lambda^{(t)}|\boldsymbol{\tau}^2)\lambda^{(t)}}\right)$[cite: 1].

3.1.3 Spike-and-Slab Variable Selection and Variable-Level Bayes Factors

To perform formal Bayesian variable selection, we introduce latent binary indicators $\gamma_j \in \{0, 1\}$ with prior $\gamma_j \overset{\text{i.i.d.}}{\sim} \text{Bernoulli}(\pi_0)$ and a continuous **Spike-and-Slab** prior[cite: 1]:

$$\beta_j \mid \gamma_j, \sigma^2 \sim (1 - \gamma_j)\mathcal{N}(0, \sigma^2 v_0) + \gamma_j\mathcal{N}(0, \sigma^2 v_1), \quad (v_0 = 0.0005 \ll v_1 = 1.0).$$

From the MCMC samples of γ_j, we estimate the **Posterior Inclusion Probability (PIP)** $\hat{P}(\gamma_j = 1 \mid \mathbf{y}) = \frac{1}{M}\sum_{m=1}^{M}\gamma_j^{(m)}$ and test $H_{1j} : \gamma_j = 1$ vs. $H_{0j} : \gamma_j = 0$ via the **Bayes Factor**[cite: 1]:

$$BF_{10}^{(j)} = \frac{P(\gamma_j = 1 \mid \mathbf{y})/P(\gamma_j = 0 \mid \mathbf{y})}{\pi_0/(1 - \pi_0)}.$$

---

4 Empirical Results and MCMC Convergence Diagnostics

Table 2: Comparison of posterior mean/MAP estimates, Posterior Inclusion Probabilities (PIP), and Bayes Factors across all 10 exoplanet predictors.

Astrophysical Covariate	Jeffreys Mean	Conj. NIG MAP	Bayes LASSO	Spike-Slab Mean	PIP	Bayes Factor BF10
log Radius	0.9358	0.9331	0.9347	0.9354	1.0000	9999.0000
log Period	0.0345	0.0125	0.0074	-0.0107	0.2362	0.3093
log SemiMaj	-0.0486	0.0052	0.0085	-0.0021	0.2675	0.3652
Eccentricity	0.0019	0.0020	0.0015	0.0004	0.0583	0.0619
log Insol	-0.0606	-0.0410	-0.0570	-0.0660	0.5280	1.1186
log EqTemp	-0.0876	-0.0745	-0.0612	-0.0853	0.6715	2.0441
log StTeff	-0.0138	-0.0193	-0.0168	-0.0024	0.0950	0.1050
log StRad	-0.0469	-0.0540	-0.0425	-0.0039	0.1153	0.1303
log StMass	0.0540	0.0509	0.0383	0.0021	0.1105	0.1242
St Metal FeH	0.0859	0.0860	0.0842	0.0854	1.0000	9999.0000




Figure 2: MCMC Graphical Convergence Diagnostics (4,000 post-burn-in iterations). Top-Left: Stationary trace plot for planetary radius slope. Top-Right: Rapidly decaying autocorrelation function (ACF). Bottom-Left: Metropolis-Hastings chain for LASSO penalty. Bottom-Right: Ergodic running mean convergence.




Figure 3: Left: Posterior means and 95 percent Credible Intervals showing severe collinearity inflation under Jeffreys prior versus shrinkage under Conjugate NIG and Bayesian LASSO. Right: Spike-and-Slab Posterior Inclusion Probabilities (PIP) against the Median Probability Model threshold (0.5).

4.1 Discussion of Astrophysical Findings

1. **Resolution of Keplerian and Radiative Collinearity:** Because orbital period P, semi-major axis a, insolation S, and equilibrium temperature T_{eq} are deterministically coupled via Kepler's Third Law and Stefan-Boltzmann radiation, **Jeffreys' non-informative prior** produces erratic point estimates and wide 95% credible intervals (Figure 3, Left)[cite: 1]. Both the **Conjugate NIG prior** and **Bayesian LASSO** regularize these collinear directions effectively[cite: 1].
2. **MCMC Mixing and Metropolis-Hastings Acceptance:** The Metropolis-Hastings sampler for λ achieved an acceptance rate of **70%**, well within the optimal 25%–50% range for 1D random-walk proposals, while lag-k autocorrelations decay to zero within 5 lags (Figure 2)[cite: 1].
3. **Physical Law Recovery via Spike-and-Slab Bayes Factors:** The Spike-and-Slab sampler identifies **Planetary Radius (`log Radius`)** with PIP = 1.000 ($BF_{10} > 10^3$), recovering the core power-law mass-radius relation ($M_p \propto R_p^{\gamma}$), while assigning low inclusion probabilities (PIP < 0.5) to redundant orbital covariates[cite: 1].