

## Dot Product Properties and Applications Learning Objectives

- Sketch simple surfaces in space
- Determine when a point lies on a specified surface.

## Dot Product examples

**Theorem.** *The angle between two vectors  $\mathbf{a}$  and  $\mathbf{b}$  is given by*

$$\theta = \cos^{-1} \left( \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|} \right)$$

*or equivalently*

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos(\theta)$$

**Theorem** (Properties of the dot product). *Let  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  be vectors and let  $c$  be a scalar:*

(a)  $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$

(b)  $(c\mathbf{a}) \cdot \mathbf{b} = c(\mathbf{a} \cdot \mathbf{b})$

(c)  $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$

(d)  $0\mathbf{a} = 0$

(e)  $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$

1. Find the angle between the following vectors

(a)  $\langle 4, 1, 1/4 \rangle, \langle 6, -3, -8 \rangle$

**Solution:**

$$|\langle 4, 1, 1/4 \rangle| = \sqrt{16 + 1 + 1/16} = \sqrt{273/16} = \sqrt{273}/4$$

$$|\langle 6, -3, -8 \rangle| = \sqrt{36 + 9 + 64} = \sqrt{109}$$

$$\langle 4, 1, 1/4 \rangle \cdot \langle 6, -3, -8 \rangle = 19$$

So the angle is

$$\cos^{-1} \left( \frac{19}{\sqrt{109}\sqrt{273}/4} \right) \approx 1.1146$$

(b)  $\mathbf{i} + \mathbf{j}, \mathbf{k}$ 

$$\cos^{-1} \left( \frac{\langle 1, 1, 0 \rangle \cdot \langle 0, 0, 1 \rangle}{|\langle 1, 1, 0 \rangle| |\langle 0, 0, 1 \rangle|} \right) = \cos^{-1}(0) = \frac{\pi}{2}$$

(c)  $\langle p, -p, 2p \rangle, \langle 2q, q, -q \rangle$  where  $p$  and  $q$  are any two non-zero real numbers.**Solution:**

$$|\langle p, -p, 2p \rangle| = \sqrt{p^2 + p^2 + 4p^2} = \sqrt{6p^2} = \sqrt{6}|p|$$

$$|\langle 2q, q, -q \rangle| = \sqrt{4q^2 + q^2 + q^2} = \sqrt{6q^2} = \sqrt{6}|q|$$

$$\langle p, -p, 2p \rangle \cdot \langle 2q, q, -q \rangle = 2pq - pq - 2pq = -pq$$

So

$$\theta = \cos^{-1} \left( \frac{-pq}{6|pq|} \right)$$

2. Let  $\mathbf{a} = \langle -2, 2, 1 \rangle$ ,  $\mathbf{b} = \langle 1, 2, 0 \rangle$  and  $\mathbf{c} = \langle 0, -1, -1 \rangle$ .(a) Find a vector  $\mathbf{v}$  so that  $\mathbf{b} \neq \mathbf{v}$  but  $\mathbf{a} \cdot \mathbf{b} = \mathbf{a} \cdot \mathbf{v}$ **Solution:** Many vectors work. Since

$$\langle -2, 2, 1 \rangle \cdot \langle 1, 2, 0 \rangle = 2$$

, if  $\mathbf{v} = \langle x, y, z \rangle$ 

$$\mathbf{a} \cdot \mathbf{v} = -2x + 2y + z$$

Any values for  $x, y, z$  that make  $-2x + 2y + z = 2$  give a vector that works. E.g.  $\langle -1, 0, 0 \rangle$ .(b) Verify part (c) above by computing  $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c})$  and  $\mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$  and showing they are the same.

# Projections

**Definition.** The *scalar projection* of  $\mathbf{b}$  onto  $\mathbf{a}$  is

$$\text{comp}_{\mathbf{a}}(\mathbf{b}) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|}$$


The *vector projection* of  $\mathbf{b}$  onto  $\mathbf{a}$  is

$$\text{proj}_{\mathbf{a}}(\mathbf{b}) = \left( \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|} \right) \left( \frac{\mathbf{a}}{|\mathbf{a}|} \right) = \left( \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{a} \cdot \mathbf{a}} \right) \mathbf{a}$$

The *orthogonal projection* of  $\mathbf{b}$  onto  $\mathbf{a}$  is

$$\text{orth}_{\mathbf{a}}(\mathbf{b}) = \mathbf{b} - \text{proj}_{\mathbf{a}}(\mathbf{b})$$

Let  $\mathbf{a} = \langle 6, 4 \rangle$  and  $\mathbf{b} = \langle 4, 0 \rangle$ , shown below



- Compute the following values and sketch them alongside the vectors  $\mathbf{a}$  and  $\mathbf{b}$  above
  - $\text{proj}_{\mathbf{a}}(\mathbf{b})$  and  $\text{orth}_{\mathbf{a}}(\mathbf{b})$
  - $\text{proj}_{\mathbf{b}}(\mathbf{a})$  and  $\text{orth}_{\mathbf{b}}(\mathbf{a})$
- What vector is equal to  $\text{proj}_{\mathbf{a}}(\mathbf{b}) + \text{orth}_{\mathbf{a}}(\mathbf{b})$ ?

3. Let  $\mathbf{a} = -5\mathbf{i} + 5\mathbf{j} + 2\mathbf{k}$  and  $\mathbf{b} = -\mathbf{i} + 8\mathbf{j} - \mathbf{k}$

(a) Compute  $\text{proj}_{\mathbf{a}}(\mathbf{b})$

(b) Compute  $\text{orth}_{\mathbf{a}}(\mathbf{b})$